

## LEAD THE RACE

Comprehensive **flow cytometry** solutions designed to accelerate your work



Learn More

MILLIPORE  
SIGMA



The Journal of  
**Immunology**

### Curcumin Inhibits Experimental Allergic Encephalomyelitis by Blocking IL-12 Signaling Through Janus Kinase-STAT Pathway in T Lymphocytes

This information is current as of January 19, 2018.

Chandramohan Natarajan and John J. Bright

*J Immunol* 2002; 168:6506-6513; ;  
doi: 10.4049/jimmunol.168.12.6506  
<http://www.jimmunol.org/content/168/12/6506>

#### Why *The JI*?

- **Rapid Reviews!** 30 days\* from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Speedy Publication!** 4 weeks from acceptance to publication

\*average

**References** This article **cites 71 articles**, 22 of which you can access for free at:  
<http://www.jimmunol.org/content/168/12/6506.full#ref-list-1>

**Subscription** Information about subscribing to *The Journal of Immunology* is online at:  
<http://jimmunol.org/subscription>

**Permissions** Submit copyright permission requests at:  
<http://www.aai.org/About/Publications/JI/copyright.html>

**Email Alerts** Receive free email-alerts when new articles cite this article. Sign up at:  
<http://jimmunol.org/alerts>



# Curcumin Inhibits Experimental Allergic Encephalomyelitis by Blocking IL-12 Signaling Through Janus Kinase-STAT Pathway in T Lymphocytes<sup>1</sup>

Chandramohan Natarajan and John J. Bright<sup>2</sup>

Experimental allergic encephalomyelitis (EAE) is a CD4<sup>+</sup> Th1 cell-mediated inflammatory demyelinating autoimmune disease of the CNS that serves as an animal model for multiple sclerosis (MS). IL-12 is a proinflammatory cytokine that plays a crucial role in the induction of neural Ag-specific Th1 differentiation and pathogenesis of CNS demyelination in EAE and MS. Curcumin (1,7-Bis(4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione) is a naturally occurring polyphenolic phytochemical isolated from the rhizome of the medicinal plant *Curcuma longa*. It has profound anti-inflammatory activity and been traditionally used to treat inflammatory disorders. In this study we have examined the effect and mechanism of action of curcumin on the pathogenesis of CNS demyelination in EAE. In vivo treatment of SJL/J mice with curcumin significantly reduced the duration and clinical severity of active immunization and adoptive transfer EAE. Curcumin inhibited EAE in association with a decrease in IL-12 production from macrophage/microglial cells and differentiation of neural Ag-specific Th1 cells. In vitro treatment of activated T cells with curcumin inhibited IL-12-induced tyrosine phosphorylation of Janus kinase 2, tyrosine kinase 2, and STAT3 and STAT4 transcription factors. The inhibition of Janus kinase-STAT pathway by curcumin resulted in a decrease in IL-12-induced T cell proliferation and Th1 differentiation. These findings highlight the fact that curcumin inhibits EAE by blocking IL-12 signaling in T cells and suggest its use in the treatment of MS and other Th1 cell-mediated inflammatory diseases. *The Journal of Immunology*, 2002, 169: 6506–6513.

Multiple sclerosis (MS)<sup>3</sup> is an inflammatory demyelinating disease of the CNS that afflicts more than one million people worldwide (1). A substantial percentage of MS patients develop clinical paralysis and become wheelchair bound or generally disabled during their lives. There is no medical treatment available so far that can cure MS (2, 3). The destruction of oligodendrocyte and myelin sheath in the CNS is the pathological hallmark of MS (4). Although the etiology is not known, MS is generally viewed as an inflammatory autoimmune disease of the CNS resulting from activation, expansion, and homing of myelin Ag-sensitized T cells in the CNS (5–7). Experimental allergic encephalomyelitis (EAE) is a CD4<sup>+</sup> Th1 cell-mediated inflammatory autoimmune demyelinating disease of the CNS (8). EAE can be induced in many susceptible strains of rodents and primates by immunizing with whole brain homogenate or purified neural Ags such as myelin basic protein (MBP), proteolipid protein, and myelin oligodendrocyte glycoprotein (9). The antigenic epitopes of neural Ags have been mapped, and adoptive transfer of T cells reactive to these epitopes is sufficient to induce the disease (10).

Many clinical and pathological features of EAE show close similarity to MS; therefore EAE has been commonly used as a model system to study the mechanism of MS pathogenesis and to test the efficacy of potential therapeutic agents for the treatment of MS.

The pathogenesis of CNS demyelination in EAE/MS is a complex process that involves activation of macrophage/microglial cells, differentiation of neural Ag-specific Th1 cells, and secretion of inflammatory cytokines in the CNS (11). IL-12 is a 70-kDa heterodimeric proinflammatory cytokine, produced mainly by macrophage/microglial cells in the CNS, that plays a critical role in the differentiation of encephalitogenic Th1 cells and pathogenesis of EAE and MS (12–14). MS patients showed increased expression of IL-12 in brain lesions, cerebrospinal fluid, and circulation in association with clinical relapses (15–20). The expression of IL-12 in the CNS and lymphoid organs of mice with EAE was also associated with the pathogenesis of CNS inflammation and demyelination (21–23). LPS and CD40 ligand (CD40L) are potent inducers of IL-12 gene expression that involves the activation of NF- $\kappa$ B signaling pathway (24, 25), whereas IL-12 signaling through Janus kinase (JAK)-STAT pathway is crucial in the induction of Th1 differentiation (26–28). We and others (29–31) have shown earlier that the inhibition of IL-12 production or IL-12 signaling prevents the differentiation of Th1 cells and pathogenesis of CNS demyelination in EAE.

Curcumin (diferuloylmethane) is a naturally occurring yellow pigment isolated from the rhizomes of the plant *Curcuma longa* (Linn), found in south Asia (32, 33). It has commonly been used as a coloring and flavoring spice in food products. Curcumin has also been traditionally used to treat many inflammatory disorders and for wound healing for centuries. The medicinal value of curcumin has been well recognized with its antioxidant, antitumor, and anti-inflammatory activities and is under preclinical trial for the treatment of cancer and inflammation (34–36). Recent studies have shown that curcumin inhibits inflammation in the animal

Division of Neuroimmunology, Department of Neurology, Vanderbilt University Medical Center, Nashville, TN 37212

Received for publication February 13, 2002. Accepted for publication April 22, 2002.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup> This work was supported in part by National Multiple Sclerosis Society Grant RG3069-A-2 (to J.J.B.).

<sup>2</sup> Address correspondence and reprint requests to Dr. John J. Bright, Department of Neurology, Vanderbilt University Medical Center, 1222F Vanderbilt Stallworth Rehabilitation Hospital, 2201 Capers Avenue, Nashville, TN 37212. E-mail address: brightj@ctrvax.vanderbilt.edu

<sup>3</sup> Abbreviations used in this paper: MS, multiple sclerosis; EAE, experimental allergic encephalomyelitis; MBP, myelin basic protein; JAK, Janus kinase; MMCS, mean maximum clinical severity; MSCH, mouse spinal cord homogenate; TYK, tyrosine kinase; CD40L, CD40 ligand.

models of atherosclerosis, Alzheimer's disease, and arthritis (37–40). The anti-inflammatory activity of curcumin was associated with its ability to inhibit the production of proinflammatory cytokines such as TNF- $\alpha$ , IL-1, IL-8, and inducible NO synthase (41–44). Although the exact mechanisms involved in the anti-inflammatory activity of curcumin is not fully defined, it prevents the activation of NF- $\kappa$ B, AP-1, and c-Jun kinase (45, 46). Recent studies have also shown that curcumin inhibits IL-12 production from macrophages and thereby prevents the differentiation of Th1 cells *in vitro* (47, 48). However, no previous study has examined the use of curcumin in the treatment of MS or other Th1 cell-mediated inflammatory diseases of the CNS.

In this study, we have examined the effects and mechanism of action of curcumin on the pathogenesis of CNS inflammation and demyelination in EAE. Our results showed that curcumin inhibits CNS demyelination by blocking IL-12 production, IL-12 signaling, and differentiation of neural Ag-specific Th1 cells in EAE and suggest its use in the treatment of MS and other Th1 cell-mediated inflammatory diseases.

## Materials and Methods

### Animals and cells

SJL/J mice were purchased from C. Reader (National Institutes of Health, Bethesda, MD) and maintained in the animal care facility at Vanderbilt University Medical Center (Nashville, TN). Activated T cells were prepared by stimulation of spleen cells from SJL/J mice ( $2 \times 10^6$ /ml) with 5  $\mu$ g/ml Con A (Pharmacia Biotech, Uppsala, Sweden) in RPMI medium supplemented with 10% FBS (Life Technologies, Rockville, MD) at 37°C and 5% CO<sub>2</sub>. After 3 days of culture, cells were harvested and cultured in medium containing 0.5% FBS for an additional 24 h to synchronize to G<sub>1</sub> phase of the cell cycle. The T cell blasts were isolated by centrifugation over Histopaque (Sigma-Aldrich, St. Louis, MO) at 1200  $\times$  g for 15 min and used for experiments (49, 50). This population of cells normally contains >98% T cell blasts as measured by flow cytometry. The peritoneal macrophage was isolated from thioglycolate-stimulated SJL/J mice as described elsewhere (51, 52). The EOC-20 mouse microglial cell line (53) was a kind gift of W. Walker (St. Jude Children's Research Hospital, Memphis, TN).

### Reagents

The curcumin (1,7-Bis(4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione) was purchased from Calbiochem (La Jolla, CA). Recombinant murine IL-12 and IFN- $\gamma$  were purchased from R&D Systems (Minneapolis, MN). The anti-IFN- $\gamma$  mAb R4-6A2 was purified from ascitic fluid collected from nude mice following transplantation of R4-6A2 hybridoma cells (no. HB 170; American Type Culture Collection, Manassas, VA). The anti-IFN- $\gamma$  mAb MM700 was obtained from Endogen (Woburn, MA) and conjugated with biotin according to standard protocol. Anti-IL-12 mAb C17.8 (anti-p40) was prepared from hybridoma cells kindly provided by G. Trinchieri (Wistar Institute, Philadelphia, PA). Mouse spinal cord homogenate (MSCH) and Guinea pig MBP were prepared according to standard protocols (21, 29–31). Anti-JAK2 Ab and anti-phosphotyrosine mAb 4G10 were purchased from Upstate Biotechnology (Lake Placid, NY). Anti-tyrosine kinase (TYK)2, anti-STAT3, and anti-STAT4 Abs were from Santa Cruz Biotechnology (Santa Cruz, CA).

### Induction and treatment of EAE

To induce active EAE, 4- to 6-wk-old female SJL/J mice were immunized (s.c.) with 800  $\mu$ g of MSCH in 150  $\mu$ l emulsion of IFA containing 50  $\mu$ g/ml H37Ra in the lower dorsum on days 0 and 7. To induce adoptive transfer EAE, 4- to 6-wk-old female SJL/J mice (donor) were immunized with 350  $\mu$ g MBP in CFA on days 0 and 7. On day 14, the lymph node and spleen cells were isolated and cultured in RPMI medium ( $5 \times 10^6$  or  $2.5 \times 10^6$  cells/ml, respectively) with 25  $\mu$ g/ml MBP. After 4 days of culture, the T cell blasts were harvested and  $1 \times 10^7$  cells were injected (i.p.) into naive female SJL/J mice. Mice in the test groups were treated (i.p.) with 50 or 100  $\mu$ g curcumin in 25  $\mu$ l DMSO on every other day from 0 to 25 days following induction of active or passive EAE. Mice in the control group received 25  $\mu$ l DMSO. The clinical paralysis in active and passive EAE was graded as follows; 0, normal; 0.5, stiff tail; 1, limp tail; 1.5, limp tail

with inability to right; 2, paralysis of one limb; 2.5, paralysis of one limb and weakness of one other limb; 3, complete paralysis of both hind limbs; 4, moribund; 5, death (21, 30).

### Histological analysis

To assess the degree of CNS inflammation and demyelination, SJL/J mice treated with curcumin following induction of active EAE were euthanized on day 15 (at the peak of the disease) by CO<sub>2</sub> asphyxiation and perfused by intracardiac injection of PBS containing 4% paraformaldehyde and 1% glutaraldehyde. Five-micrometer thick transverse sections were taken from cervical, upper thoracic, lower thoracic, and lumbar regions of the spinal cord (four sections per mouse). The sections were stained with Luxol Fast Blue (E. M. Science, Cherry Hill, NJ) to assess demyelination and with H&E to assess leukocyte infiltration and inflammation. The signs of inflammation and demyelination in the anterior, posterior, and two lateral columns (four quadrants) of the spinal cord sections were scored under microscope. Each quadrant displaying the infiltration of mononuclear cells was assigned a score of one inflammation and the quadrants that showed perivascular lesion and loss of myelin staining were assigned a score of one demyelination. Thus, each animal had a potential maximum score of 16 inflammation and/or 16 demyelination, and this study represents the analysis of 15 representative mice from three different groups. The pathologic score (inflammation or demyelination) for each group was expressed as the percentage positive over the total number of quadrants examined (29–31).

### T cell proliferation assay

The effect of curcumin on neural Ag- or IL-12-induced T cell proliferation was measured by [<sup>3</sup>H]thymidine incorporation assay. MBP-immune spleen cells were cultured in 96-well tissue culture plates in RPMI medium ( $2 \times 10^5$ /200  $\mu$ l/well) in the presence of 0, 25, or 100  $\mu$ g/ml MBP with different concentrations (0–20  $\mu$ g/ml) of curcumin. [<sup>3</sup>H]Thymidine (0.5  $\mu$ Ci/ml) was added at 72 h and the uptake of radiolabel was measured after 96 h by a Wallac beta plate scintillation counter (Wallac, Turku, Finland) (30, 31). MBP-immune spleen cells were also cultured with 0 or 25  $\mu$ g/ml MBP and 2 ng/ml rIL-12 with different concentrations (0–20  $\mu$ g/ml) of curcumin. [<sup>3</sup>H]Thymidine (0.5  $\mu$ Ci/ml) was added at 72 h and the uptake of radiolabel was measured after 96 h by a Wallac beta plate scintillation counter. Con A-activated T cells were cultured in RPMI medium in 96-well tissue culture plates ( $1 \times 10^5$ /well) with 2 ng/ml rIL-12 in the absence or presence of different concentrations of curcumin for 48 h. [<sup>3</sup>H]Thymidine was added for the last 12 h and the radiolabel was measured as above (29–31).

### Immunoprecipitation and Western blot analysis

The immunoprecipitation and Western blot analyses of JAK and STAT proteins were performed as described earlier (31, 49, 50). Briefly, Con A-activated T cells ( $2.5 \times 10^7$ /lane) were pretreated with different concentrations of curcumin for 15 min and then stimulated with 2 ng/ml IL-12 at 37°C for 15 min. Cell lysates were prepared and JAK2, TYK2, STAT3, and STAT4 proteins were immunoprecipitated using specific Abs and protein A-Sepharose. The phosphoproteins in the immune complex were analyzed by 7.5% SDS-PAGE and Western blot using anti-phosphotyrosine mAb 4G10 and visualized by an ECL detection system. The blots were stripped and reprobed with specific Ab to ensure equal protein loading.

### Culture for IL-12 and IFN- $\gamma$ assay

MBP-immune spleen cells were cultured in 24-well plates in RPMI medium ( $5 \times 10^5$ /ml) with 25  $\mu$ g/ml MBP in the presence of different concentrations of curcumin and the culture supernatants were collected after 24 h. Peritoneal macrophage and EOC-20 microglial cells were cultured in DMEM containing 10% FBS with 50 ng/ml IFN- $\gamma$  plus 1  $\mu$ g/ml LPS or 2  $\mu$ g/ml anti-CD40 Ab plus second Ab in the absence or presence of curcumin, and the culture supernatants were collected after 48 h (31, 52). Naive spleen T cells from SJL/J mice were enriched (>95% purity) by passing through a nylon wool column and plastic adherence. Anti-CD3 mAb (2C11, 5  $\mu$ g/ml) was immobilized onto six-well tissue culture plates by incubation for 1 h at 37°C. After washing the plates with PBS, T cells were added into the wells ( $1 \times 10^6$  cells/ml) and cultured in the presence or absence of 2 ng/ml rIL-12 and different concentrations of curcumin. After 5 days of culture, an equal number of viable cells ( $5 \times 10^5$  cells/ml) were restimulated with soluble anti-CD3 mAb (2C11, 5  $\mu$ g/ml), and the supernatants were collected after 36 h (30).

### ELISA for IL-12 and IFN- $\gamma$

The levels of IL-12 and IFN- $\gamma$  in the culture supernatants were measured by ELISA as described earlier (21, 29–31, 52). Briefly, ELISA plates were coated with 2  $\mu$ g/ml anti-IL-12 mAb C17.15 or anti-IFN- $\gamma$  mAb R4-6A2

capture Ab, and the residual binding sites were blocked with 3% BSA. The test samples (culture supernatants) and standards (rIL-12 or rIFN- $\gamma$ ) were added and incubated overnight at 4°C. The plates were washed and 0.2  $\mu$ g/ml biotin-conjugated anti-IL-12 mAb, C15.6, or anti-IFN- $\gamma$  mAb MM700 was added. After a 1-h incubation at room temperature, avidin-alkaline phosphatase and *p*-nitrophenyl phosphate were added and the absorbance was read at 405 nm. The levels of IL-12 and IFN- $\gamma$  in the culture supernatants were calculated from the standard curve.

## Results

### *Curcumin inhibits active immunization EAE in SJL/J mice*


To test the use of curcumin in the treatment of MS and other inflammatory demyelinating diseases of the CNS, we first examined the protective effect of curcumin on the pathogenesis of active immunization EAE. SJL/J mice were treated with curcumin (i.p., 50 and 100  $\mu$ g every other day) from day 0 to 25 following induction of active EAE by immunization with MSCH. All 15 mice in the DMSO-treated control group developed clinical paralysis for a mean duration of 16 days with a mean maximum clinical severity (MMCS) of 3.1 on day 14 (Fig. 1A). Conversely, treatment with curcumin decreased the clinical severity and duration of active EAE. The mice treated with 50  $\mu$ g curcumin developed paralysis only for a mean duration of 10 days (37.5% reduction) with a MMCS of 0.8 (74.2% reduction). In contrast, treatment with 100  $\mu$ g curcumin decreased the mean duration of disease to 8 days (50% reduction) with a MMCS of 0.3 (90.32% inhibition;  $p < 0.001$ ). These results suggest that curcumin inhibits the severity and duration of clinical paralysis in active EAE.

### *Curcumin inhibits adoptive transfer EAE in SJL/J mice*

To further study the effect of curcumin on the pathogenesis of CNS inflammation and demyelination, we examined the *in vivo* effect of curcumin on adoptive transfer EAE. SJL/J mice were treated with curcumin (i.p., 50 and 100  $\mu$ g every other day) from days 0 to 25 following induction of passive EAE by adoptive transfer of MBP-specific T cells. All 15 mice in the DMSO-treated control group developed clinical paralysis for a duration of 16 days with a MMCS of 2.8 on day 10 (Fig. 1B). Treatment of mice with curcumin decreased the clinical severity of adoptive transfer EAE. Treatment with 50  $\mu$ g curcumin decreased the development of paralysis only for a duration of 12 days (25% reduction) with a MMCS of 1.2 (64.3% inhibition), and that decreased further to a duration of 8 days (50% inhibition) with a MMCS of 0.3 (89.3% inhibition;  $p < 0.001$ ) after treatment with 100  $\mu$ g curcumin (Fig. 1B). These results suggest that curcumin inhibits the clinical severity and duration of adoptive transfer EAE.

### *Curcumin decreases CNS inflammation and demyelination in EAE*

We have further examined the effect of curcumin on the pathogenesis of inflammation and demyelination in the CNS with EAE. Spinal cord sections from mice treated with curcumin following induction of active EAE were analyzed for the infiltration of mononuclear cells (inflammation) and myelin loss (demyelination). As shown in Fig. 1C, the DMSO-treated control EAE mice showed profound inflammation and demyelination in the CNS that decreased following treatment with curcumin. In control group of EAE mice, 52% of spinal cord quadrants showed positive for inflammation and 48% of spinal cord quadrants showed positive for demyelination. Conversely, treatment with curcumin decreased inflammation and demyelination in the CNS. The treatment of mice with 50 and 100  $\mu$ g curcumin resulted in 75.9 and 95% decreases in inflammation, respectively (Fig. 1C). Treatment with 50 and 100  $\mu$ g curcumin also resulted in 67.7 and 90.6%




**FIGURE 1.** Inhibition of clinical and pathological symptoms of EAE by curcumin. *A*, Active EAE was induced in SJL/J mice by immunization with MSCH in CFA. *B*, Passive EAE was induced by adoptive transfer of MBP-sensitized T cells into recipient SJL/J mice. The mice (five per group) were treated (i.p.) with 50 or 100  $\mu$ g curcumin on every other day from days 0 to 25 following induction of EAE. The clinical symptoms were scored every day and the mean clinical score of 15 mice from three different experiments is shown. Data are representatives of five different experiments. *C*, The spinal cord sections from cervical, upper thoracic, lower thoracic, and lumbar regions were prepared (four sections per mouse) and the four quadrants of every spinal cord section were scored for the presence of inflammation or demyelination. The pathological score (one for each positive quadrant) for each treatment group was expressed as a percentage over total number of quadrants examined. The average number of quadrants examined per mouse was 16; therefore, this study included the analysis of 240 spinal cord quadrants ( $n = 3 \times 5$  mice).

decreases in CNS demyelination. These results suggest that curcumin inhibits CNS inflammation and demyelination in EAE.

### *Curcumin inhibits neural Ag-specific T cell responses*

To define the mechanisms involved in the regulation of CNS demyelination by curcumin, we examined its effect on neural Ag-specific T cell responses *in vitro*. As shown in Fig. 2A, MBP-immune T cells displayed a strong proliferative response to Ag *in vitro* and treatment with curcumin resulted in a dose-dependent




**FIGURE 2.** Inhibition of MBP-specific T cell responses by curcumin in vitro. *A*, MBP-immune spleen cells were stimulated in vitro with Ag in the presence of different concentrations of curcumin. [<sup>3</sup>H]Thymidine was added at 72 h and the uptake of radioactivity measured after 96 h. *B*, MBP-immune spleen cells were stimulated with Ag in the presence of curcumin for 48 h. The levels of IFN- $\gamma$  secreted in the culture supernatants was measured by ELISA. The values are means of triplicates at each point and the error bars represent SD. Data are representatives of five independent experiments.

decrease in proliferation. Stimulation of T cells with 25  $\mu$ g/ml MBP increased the [<sup>3</sup>H]thymidine uptake from  $649 \pm 89$  cpm in the control background to  $28,573 \pm 1,795$  cpm, and that decreased to 1,215 cpm (95.75% inhibition) following treatment with 20  $\mu$ g/ml curcumin. Stimulation of cells with 100  $\mu$ g/ml MBP induced the uptake of  $48,213 \pm 595$  cpm [<sup>3</sup>H]thymidine, and that decreased to 3,423 cpm (91.99% inhibition) following treatment with 20  $\mu$ g/ml curcumin (Fig. 2*A*). We have further examined the Ag-induced secretion of IFN- $\gamma$  from MBP-immune spleen cells in vitro. The spleen cells cultured with 100  $\mu$ g/ml MBP produced  $7.20 \pm 0.27$  ng/ml IFN- $\gamma$  in 48 h. Treatment of cells with curcumin resulted in a dose-dependent decrease in IFN- $\gamma$  production, reaching  $0.25 \pm 0.01$  ng/ml at 20  $\mu$ g/ml curcumin (96.53% inhibition, Fig. 2*B*). These results suggest that curcumin inhibits EAE by inhibiting the neural Ag-specific Th1 cell proliferation and IFN- $\gamma$  production in EAE.

#### Curcumin inhibits IL-12 production in spleen cells, macrophages, and microglia

To determine the mechanisms involved in the curcumin regulation of neural Ag-specific Th1 responses in EAE, we examined the effect of curcumin on IL-12 production in spleen cells. As shown in Fig. 3*A*, stimulation of MBP-immune spleen cells with 100  $\mu$ g/ml MBP resulted in an increase in the production of IL-12 in vitro. Treatment with curcumin decreased the neural Ag-induced IL-12 production in a dose-dependent manner with a maximum of 93.27% inhibition at 20  $\mu$ g/ml (Fig. 3*A*). Because macrophages are the major producers of IL-12 in spleen, we isolated peritoneal mac-




**FIGURE 3.** Inhibition of IL-12 production by curcumin. *A*, MBP-immune spleen cells were stimulated with 25  $\mu$ g/ml MBP in the presence of curcumin for 48 h. Peritoneal macrophage (*B*) and EOC-20 microglial cells (*C*) were stimulated with LPS plus IFN- $\gamma$  in the presence of curcumin for 48 h. The secretion of IL-12 in the culture supernatant was measured by ELISA. The values are means of triplicates at each point and the error bars represent SD. Data are representative of four different experiments.

rophages from thioglycolate-stimulated SJL/J mice and stimulated them with 50 ng/ml IFN- $\gamma$  and 1  $\mu$ g/ml LPS or anti-CD40 Ab in the absence or presence of curcumin in vitro. As shown in Fig. 3*B*, treatment of macrophage with 10  $\mu$ g/ml curcumin resulted in 95.44 and 97.02% inhibition of IL-12 production induced, respectively, by IFN- $\gamma$  plus LPS and CD40L. Because microglial cells are the major producers of IL-12 in the CNS, we have further examined the effect of curcumin on IL-12 production in microglia. Treatment of EOC-20 mouse microglial cells with 10  $\mu$ g/ml curcumin also resulted in 96.56 and 94.33% inhibition of IL-12 production induced, respectively, by IFN- $\gamma$  plus LPS and CD40L in microglia (Fig. 3*C*). These results suggest that curcumin inhibits neural Ag-specific Th1 cells by blocking IL-12 production from splenic macrophages and microglia in EAE.

### Curcumin inhibits IL-12-induced T cell responses

To determine the effect of curcumin on IL-12-induced responses in T cells, we first examined the effect of curcumin on the proliferative response of MBP-immune T cells in the presence of exogenous IL-12 in culture. As shown in Fig. 2A, curcumin inhibited the proliferation of T cells in response to MBP in vitro. However, the addition of rIL-12 failed to restore the MBP-induced proliferation of T cells from inhibition by curcumin (data not shown), suggesting that curcumin may also regulate IL-12-induced responses in T cells. To test this possibility, we have further examined the effect of curcumin on IL-12-induced proliferation of activated T cells. Stimulation of Con A-activated T cells with IL-12 resulted in a dose-dependent increase in proliferation as measured by [<sup>3</sup>H]thymidine uptake assay. The cells cultured in medium alone showed a background count of  $1,840 \pm 126$  cpm, which increased to  $43,978 \pm 2,186$  following the addition of 2 ng/ml rIL-12. Treatment of cells with curcumin resulted in a dose-dependent decrease in IL-12-induced proliferation of T cells (Fig. 4A). While the cells treated with 1  $\mu$ g/ml curcumin showed 50.02% inhibition, treatment with 10  $\mu$ g/ml curcumin resulted in 94.41% inhibition of proliferation, suggesting that curcumin inhibits IL-12-induced proliferation of activated T cells in vitro. Finally, we have also examined the effect of curcumin on IL-12-induced Th1 differentiation in vitro. When compared with T cells stimulated with anti-CD3 alone, splenic T cells activated in the presence of IL-12 displayed a dramatic increase in Th1 differentiation, as measured by




**FIGURE 4.** Inhibition of IL-12-induced responses by curcumin in T cells. *A*, Con A-activated T cells were stimulated with 2 ng/ml IL-12 in the presence of different concentrations of curcumin. [<sup>3</sup>H]Thymidine was added at 36 h and the radioactivity was measured after 48 h. The values (cpm) are mean of triplicates at each point and the error bars represent SD. Data are representative of five experiments. *B*, Naive T cells were stimulated with anti-CD3 mAb in the presence of 2 ng/ml IL-12 and different concentrations of curcumin. After 5 days, the cells were restimulated with anti-CD3 for 24 h and the IFN- $\gamma$  in the culture supernatant was measured by ELISA.

IFN- $\gamma$  production upon secondary stimulation with anti-CD3 mAb. Strikingly, addition of curcumin resulted in a dose-dependent decrease in IL-12/anti-CD3-induced IFN- $\gamma$  production with a maximum of 95.24% inhibition at 20  $\mu$ g/ml levels (Fig. 4B), suggesting that curcumin inhibits IL-12-induced differentiation of Th1 cells in vitro.

### Curcumin inhibits IL-12-induced tyrosine phosphorylation of STAT3 and STAT4 in T cells

To define the mechanisms involved in the regulation of IL-12-induced T cell responses by curcumin, we have examined the effect of curcumin on the activation of STAT transcription factors in the IL-12 signaling pathway. Stimulation of Con A-activated T cells with 2 ng/ml IL-12 induced the tyrosine phosphorylation of STAT3 and STAT4 in 15 min (Fig. 5). Pretreatment of T cells with curcumin for 15 min inhibited the IL-12-induced tyrosine phosphorylation of STAT3 and STAT4. While 10  $\mu$ g/ml curcumin induced partial inhibition, addition of 25  $\mu$ g/ml curcumin resulted in a complete inhibition in IL-12-induced tyrosine phosphorylation of STAT3 in T cells (Fig. 5A). Similarly, treatment with 10  $\mu$ g/ml curcumin induced partial inhibition of IL-12-induced STAT4



**FIGURE 5.** Inhibition of IL-12-induced tyrosine phosphorylation of STAT transcription factors by curcumin in T cells. Shown is Western blot analysis of STAT3 (*A*) and STAT4 (*B*) proteins immunoprecipitated from Con A-activated T cells after incubation in medium alone and with 2 ng/ml IL-12 in the absence or presence of different concentrations of curcumin at 37°C for 15 min. The cells were pretreated with curcumin for 15 min before the addition of IL-12. The STAT3 and STAT4 immune complexes were resolved on 7.5% SDS-PAGE, transferred onto nylon membrane, and probed with anti-phosphotyrosine mAb 4G10. The blots were stripped and reprobbed with anti-STAT3 (*A*) or STAT4 (*B*) Ab. Arrows point to the 92-kDa STAT3 (*A*) and 89-kDa STAT4 (*B*) protein bands. Data are representative of six experiments.

phosphorylation, which was also completely inhibited by the addition of 25  $\mu\text{g/ml}$  curcumin (Fig. 5B). These results suggest that curcumin inhibits IL-12-induced differentiation of Th1 cell by blocking the tyrosine phosphorylation of STAT3 and STAT4 transcription factors in T cells.

*Curcumin inhibits IL-12-induced tyrosine phosphorylation and activation of JAK2 and TYK2 in T cells*

To determine whether the inhibition of STAT proteins by curcumin was a direct effect or consequence of inhibition of upstream JAK, we examined the effect of curcumin on IL-12-induced tyrosine phosphorylation of JAK2 and TYK2 in T cells. Immunoprecipitation and Western blot analyses showed that the stimulation of Con A-activated T cells with 2 ng/ml IL-12 induced the tyrosine phosphorylation of JAK2 and TYK2 in 15 min (Fig. 6). Pretreatment of cells with curcumin reduced IL-12-induced tyrosine phosphorylation of JAK2 and TYK2 in a dose-dependent manner. While the addition of 10  $\mu\text{g/ml}$  curcumin induced partial inhibition, treatment with 25  $\mu\text{g/ml}$  curcumin resulted in a complete inhibition of JAK2 phosphorylation (Fig. 6A). Similarly, treatment with 10  $\mu\text{g/ml}$  curcumin induced partial inhibition of TYK2 phosphorylation, which was also completely inhibited following addition of 25  $\mu\text{g/ml}$  curcumin (Fig. 6B). These results

suggest that curcumin inhibits IL-12-induced STAT3 and STAT4 phosphorylation by blocking tyrosine phosphorylation and activation of the upstream JAK2 and TYK2 in T cells.


## Discussion

In this study, we have shown that curcumin is a potent regulator of IL-12 production, IL-12 signaling, Th1 differentiation, and pathogenesis of EAE in SJL/J mice. In vivo treatment of mice with curcumin decreased the clinical and pathological symptoms of active and passive EAE. The inhibition of EAE by curcumin was associated with a decrease in IL-12 production and differentiation of neural Ag-specific Th1 cells in vivo. In vitro treatment of activated T cells with curcumin inhibited IL-12-induced activation of JAK-STAT pathway, suggesting the blockade of IL-12 signaling as the molecular mechanism in the regulation of CNS demyelination by curcumin.

The pathogenesis of EAE/MS is a complex process involving the activation of macrophage/microglial cells and differentiation of neural Ag-specific Th1 cells. The proinflammatory cytokines produced by immune cells determine the final outcome of the disease (11–14). Using the EAE model of MS, we have shown earlier that in vivo treatment with lisofylline or tyrphostin inhibits the pathogenesis of CNS inflammation and demyelination in active immunization and adoptive transfer models of EAE in SJL/J mice (30, 31). In this study, we have used the EAE model to test the potential therapeutic use of curcumin in the treatment of MS. The in vivo treatment of SJL/J mice with curcumin reduced the duration and clinical severity of active immunization and adoptive transfer models of EAE. The inhibition of clinical paralysis by curcumin was associated with a decrease in inflammation and demyelination in the CNS. Earlier studies have shown that curcumin inhibits inflammatory disease models of atherosclerosis, arthritis, and Alzheimer's disease (54–58). Those studies attributed the blockade of macrophage activation and secretion of proinflammatory cytokines and chemokines as the mechanism of inhibition of the inflammatory diseases by curcumin.

Among the many proinflammatory cytokines, IL-12 is the critical one involved in the pathogenesis of CNS demyelination in EAE and MS. Earlier studies have shown that the inhibition of IL-12 production or IL-12 signaling was effective in preventing the clinical and pathological symptoms of EAE (30, 31). In this study, we have confirmed the recent reports that curcumin inhibits LPS-induced production of IL-12 in macrophage cells in vitro (47, 48). The inhibition of IL-12 production observed in this study in MBP-immune spleen cells in response to neural Ags suggests that the inhibition of IL-12 is a mechanism of regulation of EAE by curcumin. Although the exact mechanisms involved in the regulation of IL-12 gene expression by curcumin in EAE is not known, in view of the pivotal role played by NF- $\kappa\text{B}$  in IL-12 gene expression (24, 25), the blockade of NF- $\kappa\text{B}$  signaling pathway may be a molecular mechanism in the regulation of IL-12 by curcumin in EAE.

To further define the mechanisms involved in the curcumin regulation of CNS demyelination, we examined the direct effect of curcumin on neural Ag-specific T cells. In vitro treatment of MBP-immune spleen cells with curcumin resulted in a significant decrease in the neural Ag-specific T cell proliferation and IFN- $\gamma$  production. The induction of T cell proliferation is a multistep process that requires two independent signals (59). The primary signal (signal 1) is initiated through TCR that drives resting naive T cells from  $G_0$  to  $G_1$  phase of the cell cycle. In contrast, the secondary signal (signal 2) can be delivered through IL-2 and/or IL-12 receptors that are required for the transition of activated T cells from  $G_1$  to S/ $G_2$ /M phase of the cell cycle (60–63). IL-12 plays a critical role in the induction of cell cycle progression in




**FIGURE 6.** Inhibition of IL-12-induced tyrosine phosphorylation of JAK by curcumin in T cells. Shown is Western blot analysis of Con A-activated T cells incubated in medium alone and with IL-12 in the absence or presence of different concentrations of curcumin at 37°C for 15 min. The cells were pretreated with curcumin for 15 min before the induction of IL-12 signaling. The JAK2 (A) and TYK2 (B) immunoprecipitates were resolved on 7.5% SDS-PAGE, transferred onto a nylon membrane, and probed with anti-phosphotyrosine mAb 4G10. The blots were stripped and reprobbed with anti-JAK2 (A) or anti-TYK2 (B) Ab. Arrows point to 130-kDa JAK2 and 135-kDa TYK2 protein bands. Data are representative of six experiments.

activated T cells and, in particular, Th1 cells (62–64). Earlier studies have shown that curcumin inhibits Ag (signal 1)-induced T cell proliferation by inhibiting the activation of AP-1 and CD28 signaling (65). The inhibition of T cell proliferation was also attributed to a decrease in IL-12 production from Ag-presenting macrophages after pretreatment with curcumin (24, 25). This finding was further supported by restoration of Ag-induced T cell proliferation by the addition of exogenous rIL-12 to curcumin-pretreated macrophages (24, 25). However, in this study we observed that the addition of rIL-12 in the presence of curcumin in spleen cell cultures failed to restore the proliferation of neural Ag-specific T cells in vitro, suggesting that curcumin also induce a direct inhibitory effect on IL-12-induced responses in T cells.

To further determine the mechanisms involved in the inhibition of IL-12-induced T cell responses, we examined the effect of curcumin on IL-12 signaling and its consequence to T cell proliferation and Th1 differentiation in vitro. Signaling through its receptor, IL-12 induces tyrosine phosphorylation and activation of JAK2 and TYK2, leading to activation of STAT3 and STAT4 proteins (27, 28). Interestingly, we report here for the first time that treatment of activated T cells with curcumin blocked IL-12-induced tyrosine phosphorylation of STAT3 and STAT4 transcription factors in T cells. Curcumin also inhibited IL-12-induced activation of JAK2 and TYK2 in T cells, suggesting that the inhibition of STAT3 and STAT4 may be consequent to the blockade of upstream JAK. This is also consistent with our earlier report showing the inhibition of STAT3 and STAT4 phosphorylation following the blockade of IL-12-induced tyrosine phosphorylation of JAK2 and TYK2 by TGF- $\beta$  (49). However, the recent observation on the inhibition of oncostatin M-induced phosphorylation of STAT1 in chondrocytes without affecting the activation of JAK by curcumin suggested the possible direct effect of curcumin on STAT protein in T cells (66). The blockade of JAK-STAT pathway by curcumin resulted in a decrease in IL-12-induced proliferation and Th1 differentiation in activated T cells. Earlier studies

have shown that the targeted disruption or pharmacological inhibition of STAT4 activity was sufficient to block the IL-12-induced differentiation of Th1 cells in vitro and in vivo (27–31, 67). The inhibition of JAK and STAT proteins by curcumin observed in this study suggests that JAK-STAT pathway be the molecular target for curcumin regulation of neural Ag-specific Th1 cells and pathogenesis of inflammation and demyelination in EAE (Fig. 7).

The intracellular targets involved in the anti-inflammatory actions of curcumin are not well defined. Earlier studies have shown that curcumin inhibits proinflammatory molecules such as IL-12, inducible NO synthase, and cyclooxygenase-2 by blocking the activation of NF- $\kappa$ B signaling pathway in macrophages (37–48). The inhibition of IL-12 signaling through JAK-STAT pathway in T cells and IL-12-induced differentiation of Th1 cells by curcumin observed in this study imply that the blockade of JAK-STAT signaling is an important mechanism by which curcumin regulates inflammation. Although the exact mechanisms involved in the curcumin regulation of JAK-STAT pathway in EAE are not known, it is likely that curcumin may directly interact with and block the tyrosine phosphorylation and activation of JAK and STAT transcription factors and interfere with the nuclear translocation and DNA binding activity of STAT proteins in T cells. Alternatively, curcumin may activate a phosphatase or other unknown proteins that in turn inhibit IL-12 signaling through JAK-STAT pathway in T cells. At this point we do not know whether the blockade of JAK-STAT pathway by curcumin is specific to IL-12 signaling alone or common to other cytokine signaling pathways as well. Our future experiments are directed to addressing these issues. IFN- $\beta$ , which has been considered a therapeutic agent for MS, inhibits the induction of CD40-mediated IL-12 production from APC and differentiation of encephalitogenic Th1 cells (68–72). The inhibition of IL-12 production and IL-12 signaling in EAE by curcumin observed in this study places curcumin as a potent therapeutic agent with a complementary and perhaps synergistic effect with IFN- $\beta$  in the treatment of MS. In conclusion, this study highlights the fact that curcumin inhibits EAE by blocking IL-12 signaling leading to the differentiation of neural Ag-specific Th1 cells and suggests its use in the treatment of MS and other Th1 cell-mediated inflammatory diseases.



**FIGURE 7.** Curcumin regulation of IL-12 signaling in CNS demyelination. IL-12 induces tyrosine phosphorylation and activation of JAK-2, TYK2, STAT3, and STAT4, which are critical in the differentiation of neural Ag-specific Th1 cells and pathogenesis of CNS inflammation and demyelination. Treatment of cells with curcumin blocks IL-12-induced activation of JAK2, TYK2, STAT3, and STAT4 in T cells. The blockade of the JAK-STAT pathway by curcumin results in an inhibition of IL-12-induced differentiation of neural Ag-specific Th1 cells and pathogenesis of CNS inflammation and demyelination.

## References

- Dean, G. 1994. How many people in the world have MS? *Neuroepidemiology* 13:1.
- Clegg, A., and J. Bryant. 2001. Immunomodulatory drugs for multiple sclerosis: a systematic review of clinical and cost effectiveness. *Expert Opin. Pharmacother.* 2:623.
- Donoghue, S., and C. Greenlees. 2000. Drugs in development for the treatment of multiple sclerosis: antigen non-specific therapies, an update. *Expert Opin. Investig. Drugs* 9:167.
- Wingerchuk, D. M., C. F. Lucchinetti, and J. H. Noseworthy. 2001. Multiple sclerosis: current pathophysiological concepts. *Lab. Invest.* 81:263.
- Martin, R., H. F. McFarland, and D. E. McFarlin. 1992. Immunology of demyelinating disease. *Annu. Rev. Immunol.* 10:153.
- O'Connor, K. C., A. Bar-Or, D. A. Hafler. 2001. The neuroimmunology of multiple sclerosis: possible roles of T and B lymphocytes in immunopathogenesis. *J. Clin. Immunol.* 21:81.
- Gironi, M., A. Bergami, and E. Brambilla. 2000. Immunological markers in multiple sclerosis. *Neurol. Sci.* 21:5871.
- Gold, R., H. P. Hartung, and K. V. Toyka. 2000. Animal models for autoimmune demyelinating disorders of the nervous system. *Mol. Med. Today* 6:88.
- Owens, T., and S. Sriram. 1995. The immunology of multiple sclerosis and its animal model, experimental allergic encephalomyelitis. *Neurol. Clin.* 13:51.
- Xiao, B. G., and H. Link. 1995. Antigen-specific T cells in autoimmune diseases with a focus on multiple sclerosis and experimental allergic encephalomyelitis. *Cell. Mol. Life Sci.* 56:5.
- Betelli, E., and E. Nicholson. 2000. The role of cytokines in experimental autoimmune encephalomyelitis. *Arch. Immunol. Ther. Exp.* 48:389.
- Trinchieri, G. 1995. Interleukin 12: a pro-inflammatory cytokine with immunoregulatory functions that bridge innate resistance and antigen specific adaptive immunity. *Annu. Rev. Immunol.* 13:247.
- Trembleau, S. T., M. Germann, K. Gately, and L. Adorini. 1995. The role of IL-12 in the induction of organ-specific autoimmune diseases. *Immunol. Today* 16:383.
- Karp, C. L. 1999. Interleukin-12: amiss in MS. *Ann. Neurol.* 45:689.



15. Nicoletti, F., F. Patti, C. Cocuzza, P. Zaccone, A. Nicoletti, R. Di Marco, and A. Reggio. 1996. Elevated serum levels of interleukin-12 in chronic progressive multiple sclerosis. *J. Neuroimmunol.* 70:87.
16. Heesen, C., F. Sieverding, B. G. Schoser, B. Hadji, and K. Kunze. 1999. IL-12 is detectable in sera of patients with multiple sclerosis: association with chronic progressive disease course? *Eur. J. Neurol.* 6:591.
17. Fassbender, K., A. Ragoschke, S. Rossol, A. Schwartz, O. Mielke, A. Paulig, and M. Hennerici. 1998. Increased release of interleukin-12p40 in MS: association with intracerebral inflammation. *Neurology* 51:753.
18. Drulovic, J., M. Mostarica-Stojkovic, Z. Levic, S. Mesaros, N. Stojisavljevic, D. Popadi, and V. Pravica. 1998. Serum interleukin-12 levels in patients with multiple sclerosis. *Neurosci. Lett.* 251:129.
19. Comabella, M., K. Balashov, S. Issazadeh, D. Smith, H. L. Weiner, and S. J. Khoury. 1998. Elevated interleukin-12 in progressive multiple sclerosis correlates with disease activity and is normalized by pulse cyclophosphamide therapy. *J. Clin. Invest.* 102:671.
20. Balashov, K. E., D. R. Smith, S. J. Khoury, D. A. Hafler, and H. L. Weiner. 1997. Increased interleukin 12 production in progressive multiple sclerosis: induction by activated CD4<sup>+</sup> T cells via CD40 ligand. *Proc. Natl. Acad. Sci. USA* 94:599.
21. Bright, J. J., B. F. Musuro, C. Du, and S. Sriram. 1998. Expression of IL-12 in CNS and lymphoid organs of mice with experimental allergic encephalomyelitis. *J. Neuroimmunol.* 82:22.
22. Smith, T., A. K. Hewson, C. I. Kingsley, J. P. Leonard, and M. L. Cuzner. 1997. Interleukin-12 induces relapse in experimental allergic encephalomyelitis in the Lewis rat. *Am. J. Pathol.* 150:1909.
23. Leonard, J. P., K. E. Waldburger, and S. J. Goldman. 1995. Prevention of experimental autoimmune encephalomyelitis by antibodies against interleukin-12. *J. Exp. Med.* 181:381.
24. Zhang, G., and S. Ghosh. 2000. Molecular mechanisms of NF- $\kappa$ B activation induced by bacterial lipopolysaccharide through Toll-like receptors. *J. Endotoxin Res.* 6:453.
25. Murphy, T. L., M. G. Cleveland, P. Kulesza, J. Magram, and K. M. Murphy. 1995. Regulation of interleukin 12 p40 expression through an NF- $\kappa$ B half-site. *Mol. Cell. Biol.* 15:5258.
26. Darnell, J. E., Jr., I. M. Kerr, and G. R. Stark. 1994. Jak-STAT pathways and transcriptional activation in response to IFNs and other extracellular signaling proteins. *Science* 264:1415.
27. Jacobson, N. G., S. J. Szabo, R. M. Weber-Nordt, Z. Zhong, R. D. Schreiber, J. E. Darnell, Jr., and K. M. Murphy. 1995. Interleukin-12 signaling in T helper type 1 (Th1) cells involves tyrosine phosphorylation of signal transducer and activator of transcription (Stat)3 and Stat4. *J. Exp. Med.* 181:1755.
28. Bacon, C. M., E. F. Petricoin, J. R. Ortaldo, R. C. Rees, A. C. Larner, J. A. Johnston, and J. J. O'Shea. 1995. Interleukin 12 induces tyrosine phosphorylation and activation of STAT4 in human lymphocytes. *Proc. Natl. Acad. Sci. USA* 92:7307.
29. Bright J. J., M. Rodriguez, and S. Sriram. 1999. Differential influence of interleukin-12 in the pathogenesis of autoimmune and virus-induced CNS demyelination. *J. Virol.* 73:1637.
30. Bright, J. J., C. Du, M. Coon, S. Sriram, and S. J. Klaus. 1998. Prevention of experimental allergic encephalomyelitis via inhibition of IL-12 signaling and IL-12-mediated Th1 differentiation: an effect of the novel anti-inflammatory drug lisofylline. *J. Immunol.* 161:7015.
31. Bright, J. J., C. Du, and S. Sriram. 1999. Tyroprostin B42 inhibits IL-12-induced tyrosine phosphorylation and activation of Janus kinase-2 and prevents experimental allergic encephalomyelitis. *J. Immunol.* 162:6255.
32. Lodha, R., and A. Bagga. 2000. Traditional Indian systems of medicine. *Ann. Acad. Med. Singapore* 29:37.
33. Srimal, R. C., and B. N. Dhawan. 1973. Pharmacology of diferuloyl methane (curcumin), a non-steroidal anti-inflammatory agent. *J. Pharm. Pharmacol.* 25:447.
34. Araujo, C. C., and L. L. Leon. 2001. Biological activities of *Curcuma longa* L. *Mem. Inst. Oswaldo Cruz* 96:723.
35. Talalay, P. 2001. The importance of using scientific principles in the development of medicinal agents from plants. *Acad. Med.* 76:238.
36. Claeson, P., A. Panthong, P. Tuchinda, V. Reutrakul, D. Kanjanapothi, W. C. Taylor, and T. Santisuk. 1993. Three non-phenolic diarylheptanoids with anti-inflammatory activity from *Curcuma xanthorrhiza*. *Planta Med.* 59:451.
37. Mukhopadhyay, A., N. Basu, N. Ghatak, and P. K. Gujral. 1982. Anti-inflammatory and irritant activities of curcumin analogues in rats. *Agents Actions* 12:508.
38. Arora, R. B., N. Basu, V. Kapoor, and A. P. Jain. 1971. Anti-inflammatory studies on *Curcuma longa* (turmeric). *Indian J. Med. Res.* 59:1289.
39. Chandra, D., and S. S. Gupta. 1972. Anti-inflammatory and anti-arthritis activity of volatile oil of *Curcuma longa* (Haldi). *Indian J. Med. Res.* 60:138.
40. Ghatak, N., and N. Basu. 1972. Sodium curcumin as an effective anti-inflammatory agent. *Indian J. Exp. Biol.* 10:235.
41. Surh, Y. 1999. Molecular mechanisms of chemopreventive effects of selected dietary and medicinal phenolic substances. *Mutat. Res.* 428:305.
42. Brouet, I., and H. Ohshima. 1995. Curcumin, an anti-tumour promoter and anti-inflammatory agent, inhibits induction of nitric oxide synthase in activated macrophages. *Biochem. Biophys. Res. Commun.* 206:533.
43. Sreejayan, N., and M. N. Rao. 1994. Curcuminoids as potent inhibitors of lipid peroxidation. *J. Pharm. Pharmacol.* 46:1013.
44. Chan, M. M., H. I. Huang, M. R. Fenton, and D. Fong. 1998. In vivo inhibition of nitric oxide synthase gene expression by curcumin, a cancer preventive natural product with anti-inflammatory properties. *Biochem. Pharmacol.* 55:955.
45. Kakar, S. S., and D. Roy. 1994. Curcumin inhibits TPA induced expression of *c-fos*, *c-jun* and *c-myc* proto-oncogenes messenger RNAs in mouse skin. *Cancer Lett.* 87:85.
46. Surh, Y., K. Chun, H. Cha, S. S. Han, Y. Keum, K. Park, and S. S. Lee. 2001. Molecular mechanisms underlying chemopreventive activities of anti-inflammatory phytochemicals: down-regulation of COX-2 and iNOS through suppression of NF- $\kappa$ B activation. *Mutat. Res.* 480:243.
47. Kang, B. Y., S. W. Chung, W. Chung, S. Im, S. Y. Hwang, and T. S. Kim. 1999. Inhibition of interleukin-12 production in lipopolysaccharide-activated macrophages by curcumin. *Eur. J. Pharmacol.* 19:191.
48. Kang, B. Y., Y. J. Song, K. M. Kim, Y. K. Choe, S. Y. Hwang, and T. S. Kim. 1999. Curcumin inhibits Th1 cytokine profile in CD4<sup>+</sup> T cells by suppressing interleukin-12 production in macrophages. *Br. J. Pharmacol.* 128:380.
49. Bright, J. J., and S. Sriram. 1998. TGF- $\beta$  inhibits IL-12-induced activation of Jak-STAT pathway in T lymphocytes. *J. Immunol.* 161:1772.
50. Bright, J. J., L. D. Kerr, and S. Sriram. 1997. TGF- $\beta$  inhibits IL-2-induced tyrosine phosphorylation and activation of Jak1 and Stat5 in T lymphocytes. *J. Immunol.* 159:175.
51. Mishell, B. B., and S. M. Shiigi. 1980. *Selected Methods in Cellular Immunology*. Freeman, San Francisco.
52. Du, C., J. J. Bright, and S. Sriram. 2001. Inhibition of CD40 signaling pathway by tyroprostin A1 reduces secretion of IL-12 in macrophage Th1 cell development and experimental allergic encephalomyelitis in SJL/J mice. *J. Neuroimmunol.* 114:69.
53. Walker, W. S., J. Gatewood, E. Olivas, D. Askew, and C. E. Havenith. 1995. Mouse microglial cell lines differing in constitutive and interferon- $\gamma$ -inducible antigen-presenting activities for naive and memory CD4<sup>+</sup> and CD8<sup>+</sup> T cells. *J. Neuroimmunol.* 63:163.
54. Ramirez-Tortosa, M. C., M. D. Mesa, M. C. Aguilera, J. L. Quiles, L. Baro, C. L. Ramirez-Tortosa, E. Martinez-Victoria, and A. Gil. 1999. Oral administration of a turmeric extract inhibits LDL oxidation and has hypocholesterolemic effects in rabbits with experimental atherosclerosis. *Atherosclerosis* 147:371.
55. Lim, G. P., T. Chu, F. Yang, W. Beech, S. A. Frautschy, and G. M. Cole. 2001. The curry spice curcumin reduces oxidative damage and amyloid pathology in an Alzheimer transgenic mouse. *J. Neurosci.* 21:8370.
56. Kim, D. S., S. Y. Park, and J. K. Kim. 2001. Curcuminoids from *Curcuma longa* L. (Zingiberaceae) that protect PC12 rat pheochromocytoma and normal human umbilical vein endothelial cells from  $\beta$ A<sub>1-42</sub> insult. *Neurosci. Lett.* 303:57.
57. Joe, B., U. J. Rao, and B. R. Lokesh. 1997. Presence of an acidic glycoprotein in the serum of arthritic rats: modulation by capsaicin and curcumin. *Mol. Cell. Biochem.* 169:125.
58. Deodhar, S. D., R. Sethi, and R. C. Srimal. 1980. Preliminary study on antirheumatic activity of curcumin (diferuloyl methane). *Indian J. Med. Res.* 71:632.
59. Crabtree, G. R. 1989. Contingent genetic regulatory events in T lymphocyte activation. *Science* 243:355.
60. Gately, M. K., B. B. Desai, A. G. Wolitzky, P. M. Quinn, C. M. Dwyer, F. J. Podlaski, P. C. Familletti, F. Sinigaglia, R. Chizzonite, and U. Gubler. 1991. Regulation of human lymphocyte proliferation by a heterodimeric cytokine, IL-12 (cytotoxic lymphocyte maturation factor). *J. Immunol.* 147:874.
61. Bertagnolli, M. M., B. Y. Lin, D. Young, and S. H. Herrmann. 1992. IL-12 augments antigen-dependent proliferation of activated T lymphocytes. *J. Immunol.* 149:3778.
62. Yanagida, T., T. Kato, O. Igarashi, T. Inoue, and H. Nariuchi. 1994. Second signal activity of IL-12 on the proliferation and IL-2R expression of T helper cell-1 clone. *J. Immunol.* 152:4919.
63. Nguyen, T., R. Wang, and J. H. Russell. 2000. IL-12 enhances IL-2 function by inducing CD25 expression through a p38 mitogen-activated protein kinase pathway. *Eur. J. Immunol.* 30:1445.
64. Kennedy, M. K., K. S. Picha, K. D. Shanebeck, D. M. Anderson, and K. H. Grabstein. 1994. Interleukin-12 regulates the proliferation of Th1, but not Th2 or Th0, clones. *Eur. J. Immunol.* 24:2271.
65. Ranjan, D., T. D. Johnston, G. Wu, L. Elliott, S. Bondada, and M. Nagabhushan. 1998. Curcumin blocks cyclosporine A-resistant CD28 costimulatory pathway of human T-cell proliferation. *J. Surg. Res.* 77:174.
66. Li, W. Q., F. Dehnade, and M. Zafarullah. 2001. Oncostatin M-induced matrix metalloproteinase and tissue inhibitor of metalloproteinase-3 genes expression in chondrocytes requires Janus kinase/STAT signaling pathway. *J. Immunol.* 166:3491.
67. Kaplan, M. H., A. L. Wurster, and M. J. Grusby. 1998. A signal transducer and activator of transcription (Stat)4-independent pathway for the development of T helper type 1 cells. *J. Exp. Med.* 188:1191.
68. McRae, B. L., R. T. Semnani, M. P. Hayes, and G. A. van Seventer. 1998. Type I IFNs inhibit human dendritic cell IL-12 production and Th1 cell development. *J. Immunol.* 160:4298.
69. McRae, B. L., B. A. Beilfuss, and G. A. van Seventer. 2000. IFN- $\beta$  differentially regulates CD40-induced cytokine secretion by human dendritic cells. *J. Immunol.* 164:23.
70. Segal, B. M., B. K. Dwyer, and E. M. Shevach. 1998. An interleukin (IL)-10/IL-12 immunoregulatory circuit controls susceptibility to autoimmune disease. *J. Exp. Med.* 187:537.
71. Tuohy, V. K., M. Yu, L. Yin, P. M. Mathisen, J. M. Johnson, and J. A. Kawczak. 2000. Modulation of the IL-10/IL-12 cytokine circuit by interferon- $\beta$  inhibits the development of epitope spreading and disease progression in murine autoimmune encephalomyelitis. *J. Neuroimmunol.* 111:55.
72. Karp, C. L., C. A. Biron, and D. N. Irani. 2000. Interferon  $\beta$  in multiple sclerosis: is IL-12 suppression the key? *Immunol. Today* 21:24.