



The Attune™ NxT Flow Cytometer system

Let's get to the science

Find out more

invitrogen  
by Thermo Fisher Scientific



## Cutting Edge: Tissue-Resident Memory CTL Down-Regulate Cytolytic Molecule Expression following Virus Clearance

This information is current as of April 22, 2019.

Justine D. Mintern, Carole Guillonneau, Francis R. Carbone, Peter C. Doherty and Stephen J. Turner

*J Immunol* 2007; 179:7220-7224; ;  
doi: 10.4049/jimmunol.179.11.7220  
<http://www.jimmunol.org/content/179/11/7220>

**References** This article **cites 24 articles**, 16 of which you can access for free at:  
<http://www.jimmunol.org/content/179/11/7220.full#ref-list-1>

Why *The JI*? [Submit online.](#)

- **Rapid Reviews! 30 days\*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

*\*average*

**Subscription** Information about subscribing to *The Journal of Immunology* is online at:  
<http://jimmunol.org/subscription>

**Permissions** Submit copyright permission requests at:  
<http://www.aai.org/About/Publications/JI/copyright.html>

**Email Alerts** Receive free email-alerts when new articles cite this article. Sign up at:  
<http://jimmunol.org/alerts>

*The Journal of Immunology* is published twice each month by  
The American Association of Immunologists, Inc.,  
1451 Rockville Pike, Suite 650, Rockville, MD 20852  
Copyright © 2007 by The American Association of  
Immunologists All rights reserved.  
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



## Cutting Edge: Tissue-Resident Memory CTL Down-Regulate Cytolytic Molecule Expression following Virus Clearance<sup>1</sup>

Justine D. Mintern,<sup>\*</sup> Carole Guillonneau,<sup>\*</sup> Francis R. Carbone,<sup>\*</sup> Peter C. Doherty,<sup>\*†</sup> and Stephen J. Turner<sup>2\*</sup>

*CTL express lytic proteins that mediate the cytolysis of virus-infected cells. In this study, cytolitic transcriptional profiles were determined for individual CTL responding to influenza A virus and HSV-1. During acute infection, influenza-specific CTL in the spleen and respiratory airways displayed highly activated cytolitic profiles, as did HSV-1-specific CTL localized in the spleen, skin, and dorsal root ganglia (DRG). In contrast, memory CTL dramatically down-regulated cytolitic molecule transcription. This occurred for both lymphoid (spleen) and tissue-resident (skin and/or lung) memory CTL. In contrast, HSV-1-specific CTL localized in the dorsal root ganglia in the presence latent HSV-1 Ag did not down-regulate cytolitic molecule transcription. Therefore, both lymphoid and tissue-resident memory CTL down-regulate cytolitic molecule transcription following virus clearance unless localized Ag is present. The Journal of Immunology, 2007, 179: 7220–7224.*

Cytotoxic T lymphocytes are critical participants in viral immunity. One of their major roles is the destruction of infected cells via cytolitic granules containing lytic proteins, including perforin (Pfp)<sup>3</sup> and granzyme (Gzm) serine protease family members. Pfp is critical for protease delivery, whereas Gzms initiate proteolysis and trigger apoptotic pathways (1). Individual virus-specific CTL express heterogeneous patterns of Pfp and specific Gzm family members (2, 3). This unique cytolitic profile is the outcome of specific transcriptional regulation of the different lytic proteins (3, 4).

In response to virus, CD8<sup>+</sup> T cells acquire effector function, including the expression of cytotoxic molecules. Once virus is cleared, memory CTL reside in lymphoid organs and peripheral tissue(s) associated with infection. Memory CTL are considered to maintain cytolitic molecule transcription as a means of Ag preparedness. Lymphoid-resident memory CTL express some

cytolitic transcripts, although at reduced levels compared with peak immunity (2, 5, 6). Little to no cytolitic protein is expressed (5, 7, 8), which avoids the dangerous scenario of improper cytolitic molecule release. Cytolitic transcription by tissue-resident memory CTL has not been examined. These cells display potent cytolitic function (8–10) and are predicted to display activated cytolitic transcriptional profiles. In this study, we investigated cytolitic transcriptional profiles using two models of anti-viral CTL immunity; infection of the respiratory tract with influenza A virus and infection of the skin epidermis with HSV-1. Together, these models provide lymphoid-resident (spleen) and tissue-resident (lung, skin, and dorsal root ganglia (DRG) virus-specific CTL. Cytolitic transcriptional profiling of single CTL showed that both lymphoid and tissue-resident memory CTL expressed reduced cytolitic transcription unless localized Ag was present.

### Materials and Methods

#### Mice

C57BL6/J (B6), C57BL6/J-TgN(TCR-HSV)gBT-1.1Frc (gBT-1), and gBT-1 × B6.SJL-Ptprc<sup>a</sup>Pep3<sup>b</sup>/BoyJ (gBT-1 × B6.Ly5.1) mice (gBT is gB-specific CTL) were maintained at the Department of Microbiology and Immunology, University of Melbourne (Parkville, Victoria, Australia). The gBT-1 TCR transgenic mice are specific for the H-2K<sup>b</sup>-restricted HSV-1 epitope gB<sub>498–505</sub>-SS-IEFARL (gB) (11).

#### Virus infection

B6 mice were adoptively transferred with  $5 \times 10^4$  gBT lymph node cells i.v. before infection. Mice were inoculated with  $1 \times 10^6$  PFU of HSV-1 (KOS strain) in a region of scarified flank skin or with intranasal administration of 50 PFU of flu-gB. At days 7–8 (acute) and days 20–21 (late) following infection, spleen, lung, skin, and DRG were harvested. Lymphocytes were isolated by mechanical disruption or collagenase (Roche) digestion.

#### Flow cytometry

Cells were stained with anti-CD8 $\alpha$ -allophycocyanin and anti-Ly5.1-PE (BD Pharmingen). CD8<sup>+</sup>Ly5.1<sup>+</sup> cells were sorted into individual wells of a 96-well plate (Eppendorf) by using a FACSAria flow cytometer with a single cell deposition unit (BD Biosciences).

<sup>\*</sup>Department of Microbiology and Immunology, University of Melbourne, Parkville, Victoria, Australia; and <sup>†</sup>Department of Immunology, St. Jude Children's Research Hospital, Memphis, TN 38105

Received for publication July 17, 2007. Accepted for publication September 26, 2007.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.


<sup>1</sup>J.D.M. was supported by a National Health and Medical Research Council (NHMRC; Australia) C. J. Martin Fellowship; C.G. was supported by a Marie Curie Outgoing

International Fellowship; F.R.C. was supported by NHMRC (Australia); P.C.D. was supported by a NHMRC (Australia) Burnet Grant; and S.J.T. was supported by a NHMRC (Australia) R. D. Wright Fellowship.

<sup>2</sup>Address correspondence and reprint requests to Dr. Stephen J. Turner, Department of Microbiology and Immunology, University of Melbourne, Parkville, Victoria 3010, Australia. E-mail address: stjturn@unimelb.edu.au

<sup>3</sup>Abbreviations used in this paper: Pfp, perforin; Gzm, granzyme; DRG, dorsal root ganglia; gBT, gB-specific CTL.

Copyright © 2007 by The American Association of Immunologists, Inc. 0022-1767/07/\$2.00



**FIGURE 1.** HSV-1 vs flu-gB-specific CTL do not exhibit unique Gzm preferences during acute immunity. gBT were isolated from pooled spleens following HSV-1 or flu-gB infection at acute time points. Patterns of cytolytic transcripts expressed per single CD8<sup>+</sup> CTL are shown (mean). A filled circle represents an expressed transcript. For HSV-1 infection,  $n = 144$  CD8<sup>+</sup> T cells harvested from six mice for three individual experiments. For flu-gB infection,  $n = 76$  CD8<sup>+</sup> T cells harvested from five mice for two individual experiments.

#### Cytolytic profiling


cDNA was generated and two rounds of nested PCR were performed (2, 3). Samples that were not CD8 $\alpha$ -positive were excluded from the analysis; the data was analyzed with Student's  $t$  test.

## Results and Discussion

### *HSV-1 and flu-gB-specific gBT CTL do not exhibit unique Gzm preferences during acute immunity*

Cytolytic transcriptional profiling of CTL responding to HSV-1 and/or influenza was undertaken by single cell RT-PCR (2, 3). The frequency, number, and pattern of Pfp and Gzms A, B, C and K were assessed. To allow direct comparison between the two viruses, we used an engineered influenza virus expressing the HSV-1 gB epitope, gB<sub>498–505</sub> (flu-gB) (12). gB-specific CTL were studied upon the adoptive transfer of low numbers ( $5 \times 10^4$ ) of Ly5.1<sup>+</sup> gBT (gBT) before infection. This enabled CTL isolation without MHC class I tetramer reagents that might trigger the TCR and inadvertently alter cytolytic granule composition. Naive gBT did not express cytolytic transcripts (data not shown).

We determined whether cytolytic transcription was intrinsic or modulated by different viral infection contexts. In support of the latter, Gzm A (13) and Gzm B-deficient mice (14) display increases in HSV-1 susceptibility, whereas influenza elicits robust CTL immunity in the absence of Gzm A and/or Gzm B (15). Given this, we predicted an increase in CTL using Gzm A during HSV-1 infection compared with influenza. gBT were isolated from spleens following infection with either flu-gB or HSV-1 at acute (days 7–8) time points. For both viruses, this is the peak in CTL magnitude. Flu-gB and HSV-1-specific gBT expressed the same range of cytolytic phenotypes; 14 different patterns were observed (Fig. 1). For both infections, the prevalent phenotype was a combination of Pfp and Gzm A, B, and K. Few virus-specific CTL expressed Gzm C, confirming a limited role for Gzm C in viral immunity *in vivo* (2, 3) despite it being highly expressed in activated CTL *in vitro* (4). Some differences were noted. Flu-gB elicited a higher frequency of gBT expressing either Gzm B or Gzm K alone, however these cells repre-



**FIGURE 2.** Spleen-resident flu-gB-specific CTL display reduced cytolytic transcriptional profiles at late time points compared with acute immunity. gBT were isolated from pooled spleens following flu-gB infection at acute or late time points. Frequency (A) and number (B) of cytolytic transcripts expressed per CD8<sup>+</sup> CTL is shown (mean). For acute time points,  $n = 76$  CD8<sup>+</sup> T cells harvested from five mice for two individual experiments (same CTL as that analyzed in Fig. 1). For late time points,  $n = 90$  CD8<sup>+</sup> T cells harvested from five mice for two individual experiments.

sented only minor populations overall. In response to both viruses, gBT expressed similar numbers of cytolytic transcripts per cell, where the mean number expressed per flu-gB-specific gBT was  $2.9 \pm 1.3$  ( $n = 144$ ) compared with  $2.5 \pm 1.4$  ( $n = 76$ ) for HSV-1-specific gBT. Therefore, cytolytic transcriptional profiles elicited by the distinct viruses were similar, heterogeneous, and resembled those described previously for endogenous influenza-specific CTL (2, 3). Specifically, there was no increase in the prevalence of Gzm A-expressing CTL in response to HSV-1 vs flu-gB infection. This suggests that CTL cytolytic transcription is determined intrinsically rather than being modulated by the specific viral infection context. This does not rule out, however, the possibility that cytolytic transcriptional profiles may differ when examining CTL responses to other pathogens or, indeed, other foreign Ags (e.g., tumor cells).

### *Spleen-resident flu-gB-specific CTL display reduced cytolytic transcriptional profiles at late time points compared with acute immunity*

Previous studies by our laboratory showed that endogenous influenza-specific memory CTL in the spleen down-regulate cytolytic transcription compared with transcription during peak immunity (2). We determined whether this was also the case for spleen-resident memory CTL following flu-gB infection. Following acute infection, a high frequency of gBT expressed cytolytic transcripts was detected in Gzm A- and Gzm K-expressing CTL (Fig. 2A). In contrast, at late time points a reduced frequency of flu-gB-specific gBT expressing cytolytic transcripts was detected in Gzm A- and Gzm K-expressing CTL (Fig. 2A). The failure to maintain Gzm A expression supports previous studies (2, 5) but contrasts with human studies where memory CTL acquire Gzm A (16). This provides further evidence that mouse and human Gzms are structurally



and functionally divergent (17). At acute time points 31.5% of flu-gB-specific gBT expressed four cytolytic transcripts per cell, whereas at late time points no flu-gB-specific CTL were detected that expressed four cytolytic transcripts per cell (Fig. 2B). This was reflected in the significant reduction ( $p < 0.05$ ) in the mean number of cytolytic transcripts expressed per CTL at late time points:  $1.0 \pm 1.1$  ( $n = 90$ ) compared with  $2.5 \pm 1.4$  ( $n = 76$ ) at acute time points. Therefore, on a per cell basis spleen-resident flu-gB-specific memory CTL expressed reduced cytolytic molecule transcripts at late time points compared with peak immunity. It should be noted that memory CTL do not resemble naive  $CD8^+$  T cell precursors that lack cytolytic molecule expression altogether. In all cases, a considerable frequency expressed Gzm B and/or Gzm K (Fig. 2). These may simply represent longer lived mRNA transcripts or, alternatively, these Gzms may possess functions important for memory CTL other than the cytolysis of infected cells. This requires further investigation.


*Respiratory tract-resident flu-gB specific CTL display reduced cytolytic transcriptional profiles at late time points compared with acute immunity*

Of interest was the cytolytic transcriptional profiles of influenza-specific memory CTL resident in the respiratory airways. To examine this, flu-gB-specific gBT were harvested from the respiratory tissue at acute (days 7–8) vs late (days 20–21) time points. At late time points following flu-gB infection, a reduced frequency of flu-gB-specific gBT expressing cytolytic transcripts was detected in Pfp-expressing CTL compared with peak immunity (Fig. 3A). More than 50% of CTL examined at late time points did not express any cytolytic transcripts at all, with fewer CTL expressing three cytolytic transcripts per CTL compared with acute time points (Fig. 3B). A significant reduction ( $p < 0.05$ ) in the mean number of cytolytic transcripts expressed per CTL at late time points was observed:  $1.3 \pm 0.9$  ( $n = 57$ ) compared with  $2.5 \pm 1.2$  ( $n = 66$ ) at acute time points. Therefore, on a per cell basis lung-resident flu-gB-specific memory CTL expressed reduced cytolytic molecule transcripts at late time points compared with peak immunity.

Despite expressing an activated phenotype (18), memory CTL residing in the lung parenchyma down-regulate their cytolytic molecule transcription. This implies that influenza A virus Ag is no longer present and provides no support for the recent notion that influenza Ag persists in long-lived depots (19). More importantly, this data suggests that tissue-resident memory CTL are similar to lymphoid-resident memory CTL in that they do not express high levels of cytolytic transcripts.

*Skin-resident HSV-1-specific CTL display reduced cytolytic molecule transcription at late time points compared with acute immunity*

HSV-1 infection of skin epidermis (20, 21) provided a distinct anatomical location (skin) for examining tissue-resident memory CTL. This was important, as the down-regulation of cytolytic transcription by lung-resident memory CTL may result from the environmental milieu present in respiratory airways. In brief, HSV-1 infection involves skin inoculation, virus access to the sensory neurons, and virus re-emergence to promote a skin zosteriform lesion. The virus is cleared from the skin by day 7 (20) but establishes a persistent reservoir of latent virus in the DRG (22).




**FIGURE 3.** Respiratory tract-resident flu-gB specific CTL display reduced cytolytic transcriptional profiles at late time points compared with acute immunity. gBT were isolated from respiratory airways following flu-gB infection at acute or late time points. Frequency (A) and number (B) of cytolytic transcripts expressed per  $CD8^+$  CTL is shown (mean). For acute time points,  $n = 66$   $CD8^+$  T cells harvested from six mice for two individual experiments. For late time points  $n = 57$   $CD8^+$  T cells harvested from eight mice for two individual experiments.

To investigate cytolytic transcription in skin-resident HSV-1-specific CTL, gBT were isolated at acute (days 7–8) vs late (days 20–21) time points following HSV-1 infection. Pfp and Gzm A, B, and K were expressed by a high frequency of skin HSV-1-specific gBT CTL during acute infection. In contrast, all cytolytic transcripts were represented at significantly reduced ( $p < 0.05$ ) frequencies at late time points (Fig. 4A). As such, a dramatic down-regulation in the number of cytolytic transcripts expressed per cell was detected for skin HSV-1-specific gBT (Fig. 4B). At late time points, gBT expressed  $1.2 \pm 1.2$  ( $n = 153$ ) cytolytic transcripts per cell, significantly reduced ( $p < 0.05$ ) from the acute response where gBT expressed  $3.0 \pm 1.1$  ( $n = 146$ ) cytolytic transcripts per cell. Therefore, similar to the lung, memory CTL that reside in the skin following virus clearance display reduced cytolytic transcription compared with peak CTL immunity.

*DRG-resident HSV-1-specific CTL display similar cytolytic molecule transcription at late time points compared with acute immunity*

A unique feature of HSV-1 infection is that it provides a source of tissue-resident CTL localized to DRG. In this site, latent HSV-1 provides continuous stimulation of resident CTL by infected parenchymal cells (22). Therefore, a direct comparison of tissue-resident CTL that persist in the presence (DRG) or absence (skin) of Ag could be undertaken. CTL resident in neuronal cell bodies were also of interest, given that although they are reported to display an activated phenotype, including expression of Gzm B protein (22, 23), they are not considered to use cytotoxic mechanisms to combat virus infection (24).


HSV-1-specific gBT were isolated from DRG innervating the HSV-1 inoculation site during both acute (days 7–8) and late (days 20–21) CTL immunity. A high frequency of DRG HSV-1-specific gBT isolated at both time points expressed Pfp



**FIGURE 4.** Skin-resident HSV-1-specific CTL display reduced cytolysis transcriptional profiles at late time points compared with acute immunity. gBT were isolated from pooled skin samples following HSV-1 infection at acute or late time points. Frequency (*A*) and number (*B*) of cytolysis transcripts expressed per CD8<sup>+</sup> CTL is shown (mean  $\pm$  SEM). For acute time points,  $n = 146$  CD8<sup>+</sup> T cells harvested from six mice for three individual experiments. For late time points,  $n = 153$  CD8<sup>+</sup> T cells harvested from 15 mice for three individual experiments. \*,  $p < 0.05$ ; significantly different.

and Gzm A, B, and K (Fig. 5*A*). At both time points, the percentage of gBT expressing one, two, three, and four cytolysis transcripts per CTL was similar (Fig. 5*B*). This was also the case for the mean number of cytolysis transcripts expressed per cell: DRG-resident gBT expressed  $2.7 \pm 1.5$  ( $n = 151$ ) transcripts at acute time points compared with  $2.8 \pm 1.3$  at late time points. Therefore, for DRG-resident HSV-1-specific gBT, similar cytolysis transcriptional profiles were observed at both acute and late time points following infection. This has a number of implications. First, the inhibition of cytotoxic function reported for DRG-resident CTL (24) does not involve the termination of transcription of Pfp, Gzm A, Gzm B, and/or Gzm K. Second, latent HSV-1 is capable of eliciting a similar cytolysis transcriptional program to acute HSV-1 infection. Third, when DRG-resident CTL are compared with skin-resident CTL at the same time points (Fig. 4), it is apparent that cytolysis transcription is maintained only in the presence of ongoing localized Ag expression.

In summary, our assessment of CTL cytolysis transcriptional profiles over the course of viral immunity suggests that, unless persisting Ag is present, anti-viral memory CTL decrease their cytolysis transcription regardless of whether the CTL remain localized at the site of former infection or the lymphoid tissue. Therefore, tissue-resident memory CTL are no more poised for cytotoxic effector function than their lymphoid counterparts despite possessing the capacity to initiate rapid cytotoxicity (7–10). The inability to sustain the transcription of cytolysis molecules in the absence of localized Ag suggests this is tightly regulated and requires an Ag-specific signal via the TCR. This differs from the expression of activation markers such as CD69, which can be elicited in response to inflammatory stimuli in the



**FIGURE 5.** DRG-resident HSV-1-specific CTL display similar cytolysis transcriptional profiles at late time points compared with acute immunity. gBT were isolated from pooled DRG following HSV-1 infection at acute or late time points. Frequency (*A*) and number (*B*) of cytolysis transcripts expressed per CD8<sup>+</sup> CTL is shown. For acute time points,  $n = 151$  CD8<sup>+</sup> T cells, harvested from six mice for three individual experiments. At late time points,  $n = 92$  CD8<sup>+</sup> T cells harvested from 15 mice for three individual experiments.

absence of Ag (22). Overall, this analysis suggests that the selective advantage memory CTL possess, when compared with their naive CD8 T cell precursors, is due to increased frequency and/or anatomical distribution in sites of potential reinfection but is not due to enhanced cytotoxic capacity.

## Acknowledgments

We thank Nicole La Gruta, Katherine Kedzierska, and Thomas Gebhardt for critical review of the manuscript, Dina Stockwell and Lauren Kleinhert for technical assistance, and Ken Field for excellent flow cytometry.

## Disclosures

The authors have no financial conflict of interest.

## References

- Voskoboinik, I., M. J. Smyth, and J. A. Trapani. 2006. Perforin-mediated target-cell death and immune homeostasis. *Nat Rev Immunol.* 6: 940–952.
- Jenkins, M. R., K. Kedzierska, P. C. Doherty, and S. J. Turner. 2007. Heterogeneity of effector phenotype for acute phase and memory influenza A virus-specific CTL. *J. Immunol.* 179: 64–70.
- Johnson, B. J., E. O. Costelloe, D. R. Fitzpatrick, J. B. Haanen, T. N. Schumacher, L. E. Brown, and A. Kelso. 2003. Single-cell perforin and granzyme expression reveals the anatomical localization of effector CD8<sup>+</sup> T cells in influenza virus-infected mice. *Proc. Natl. Acad. Sci. USA* 100: 2657–2662.
- Kelso, A., E. O. Costelloe, B. J. Johnson, P. Groves, K. Buttigieg, and D. R. Fitzpatrick. 2002. The genes for perforin, granzymes A-C and IFN- $\gamma$  are differentially expressed in single CD8<sup>+</sup> T cells during primary activation. *Int. Immunol.* 14: 605–613.
- Kaech, S. M., S. Hemby, E. Kersh, and R. Ahmed. 2002. Molecular and functional profiling of memory CD8 T cell differentiation. *Cell* 111: 837–851.
- Peixoto, A., C. Evaristo, I. Munitic, M. Monteiro, A. Charbit, B. Rocha, and H. Veiga-Fernandes. 2007. CD8 single-cell gene coexpression reveals three different effector types present at distinct phases of the immune response. *J. Exp. Med.* 204: 1193–1205.
- Wherry, E. J., V. Teichgraber, T. C. Becker, D. Masopust, S. M. Kaech, R. Antia, U. H. von Andrian, and R. Ahmed. 2003. Lineage relationship and protective immunity of memory CD8 T cell subsets. *Nat. Immunol.* 4: 225–234.
- Byers, A. M., C. C. Kamball, J. M. Moser, and A. E. Lukacher. 2003. Cutting edge: rapid in vivo CTL activity by polyoma virus-specific effector and memory CD8<sup>+</sup> T cells. *J. Immunol.* 171: 17–21.
- Masopust, D., V. Vezys, A. L. Marzo, and L. Lefrançois. 2001. Preferential localization of effector memory cells in nonlymphoid tissue. *Science* 291: 2413–2417.

10. Barber, D. L., E. J. Wherry, and R. Ahmed. 2003. Cutting edge: rapid in vivo killing by memory CD8 T cells. *J. Immunol.* 171: 27–31.
11. Mueller, S. N., W. Heath, J. D. McLain, F. R. Carbone, and C. M. Jones. 2002. Characterization of two TCR transgenic mouse lines specific for herpes simplex virus. *Immunol. Cell Biol.* 80: 156–163.
12. Stock, A. T., C. M. Jones, W. R. Heath, and F. R. Carbone. 2006. CTL response compensation for the loss of an immunodominant class I-restricted HSV-1 determinant. *Immunol. Cell Biol.* 84: 543–550.
13. Pereira, R. A., M. M. Simon, and A. Simmons. 2000. Granzyme A, a noncytolytic component of CD8<sup>+</sup> cell granules, restricts the spread of herpes simplex virus in the peripheral nervous systems of experimentally infected mice. *J. Virol.* 74: 1029–1032.
14. Heusel, J. W., R. L. Wesselschmidt, S. Shresta, J. H. Russell, and T. J. Ley. 1994. Cytotoxic lymphocytes require granzyme B for the rapid induction of DNA fragmentation and apoptosis in allogeneic target cells. *Cell* 76: 977–987.
15. Mullbacher, A., P. Waring, R. Tha Hla, T. Tran, S. Chin, T. Stehle, C. Museteanu, and M. M. Simon. 1999. Granzymes are the essential downstream effector molecules for the control of primary virus infections by cytolytic leukocytes. *Proc. Natl. Acad. Sci. USA* 96: 13950–13955.
16. Takata, H., and M. Takiguchi. 2006. Three memory subsets of human CD8<sup>+</sup> T cells differently expressing three cytolytic effector molecules. *J. Immunol.* 177: 4330–4340.
17. Kaiserman, D., C. H. Bird, J. Sun, A. Matthews, K. Ung, J. C. Whisstock, P. E. Thompson, J. A. Trapani, and P. I. Bird. 2006. The major human and mouse granzymes are structurally and functionally divergent. *J. Cell Biol.* 175: 619–630.
18. Hogan, R. J., E. J. Usherwood, W. Zhong, A. A. Roberts, R. W. Dutton, A. G. Harmsen, and D. L. Woodland. 2001. Activated antigen-specific CD8<sup>+</sup> T cells persist in the lungs following recovery from respiratory virus infections. *J. Immunol.* 166: 1813–1822.
19. Zammit, D. J., D. L. Turner, K. D. Klonowski, L. Lefrancois, and L. S. Cauley. 2006. Residual antigen presentation after influenza virus infection affects CD8 T cell activation and migration. *Immunity* 24: 439–449.
20. van Lint, A., M. Ayers, A. G. Brooks, R. M. Coles, W. R. Heath, and F. R. Carbone. 2004. Herpes simplex virus-specific CD8<sup>+</sup> T cells can clear established lytic infections from skin and nerves and can partially limit the early spread of virus after cutaneous inoculation. *J. Immunol.* 172: 392–397.
21. Simmons, A., and A. A. Nash. 1984. Zosteriform spread of herpes simplex virus as a model of recrudescence and its use to investigate the role of immune cells in prevention of recurrent disease. *J. Virol.* 52: 816–821.
22. van Lint, A. L., L. Kleinert, S. R. Clarke, A. Stock, W. R. Heath, and F. R. Carbone. 2005. Latent infection with herpes simplex virus is associated with ongoing CD8<sup>+</sup> T-cell stimulation by parenchymal cells within sensory ganglia. *J. Virol.* 79: 14843–14851.
23. Verjans, G. M., R. Q. Hintzen, J. M. van Dun, A. Poot, J. C. Milikan, J. D. Laman, A. W. Langerak, P. R. Kinchington, and A. D. Osterhaus. 2007. Selective retention of herpes simplex virus-specific T cells in latently infected human trigeminal ganglia. *Proc. Natl. Acad. Sci. USA* 104: 3496–3501.
24. Suvas, S., A. K. Azkur, and B. T. Rouse. 2006. Qa-1b and CD94-NKG2a interaction regulate cytolytic activity of herpes simplex virus-specific memory CD8<sup>+</sup> T cells in the latently infected trigeminal ganglia. *J. Immunol.* 176: 1703–1711.