

ELISA KITS you can trust.

ISO Certified / Made in the USA / 100% Guaranteed

 RayBiotech
Empowering your proteomics

[Browse ELISAs](#)

 *The Journal of*
Immunology

Pillars Article: CD19: Lowering the Threshold for Antigen Receptor Stimulation of B Lymphocytes. *Science*, 1992. 256: 105–107

This information is current as of December 18, 2017.

Robert H. Carter and Douglas T. Fearon

J Immunol 2010; 184:2233-2235; ;
<http://www.jimmunol.org/content/184/5/2233.citation>

Why *The JI*?

- **Rapid Reviews! 30 days*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Speedy Publication!** 4 weeks from acceptance to publication

**average*

Subscription Information about subscribing to *The Journal of Immunology* is online at:
<http://jimmunol.org/subscription>

Permissions Submit copyright permission requests at:
<http://www.aai.org/About/Publications/JI/copyright.html>

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://jimmunol.org/alerts>

The Journal of Immunology is published twice each month by
The American Association of Immunologists, Inc.,
1451 Rockville Pike, Suite 650, Rockville, MD 20852
Copyright © 2010 by The American Association of
Immunologists, Inc. All rights reserved.
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



CD19: Lowering the Threshold for Antigen Receptor Stimulation of B Lymphocytes

Robert H. Carter* and Douglas T. Fearon

Lymphocytes must proliferate and differentiate in response to low concentrations of a vast array of antigens. The requirements of broad specificity and sensitivity conflict because the former is met by low-affinity antigen receptors, which precludes achieving the latter with high-affinity receptors. Coligation of the membrane protein CD19 with the antigen receptor of B lymphocytes decreased the threshold for antigen receptor–dependent stimulation by two orders of magnitude. B lymphocytes proliferated when approximately 100 antigen receptors per cell, 0.03 percent of the total, were coligated with CD19. The B cell resolves its dilemma by having an accessory protein that enables activation when few antigen receptors are occupied.

The immune system must respond to low concentrations of antigen for the efficient elimination of infections. The growth and differentiation of lymphocytes are mediated by antigen receptors that have low affinity for their ligands because they are products of recombinatorial gene rearrangement that takes place in the absence of selection by antigen. Therefore, lymphocytes must have mechanisms that enable them to be stimulated when relatively few receptors have bound antigen. The T lymphocyte has the accessory membrane proteins CD4 and CD8, which, when coligated with the T cell antigen receptor (TCR) by the major histocompatibility complex (MHC)–peptide complex, decrease the number of TCRs that must be ligated (1). No membrane protein of the B lymphocyte that is analogous to CD4 and CD8 has been identified.


The CD19 membrane protein, a member of the immunoglobulin (Ig) superfamily, is B cell–specific and is expressed at each developmental stage except that of the terminally differentiated plasma cell (2). It is a component of a complex that contains at least two other membrane proteins, complement receptor type 2 (CR2, also called CD21) and TAPA-1 (3). The CR2 protein mediates the capacity of the complement system to enhance the production of antibody in response to low concentrations of

Division of Molecular and Clinical Rheumatology, Department of Medicine, and the Graduate Program in Immunology, Johns Hopkins University School of Medicine, Baltimore, MD 21205.

*To whom correspondence should be addressed.

antigen *in vivo* (4). Both CD19 and CR2 augment activation of phospholipase C in B cells when coligated with membrane Ig (mIg), but independent ligation of CD19 suppresses B cell activation (5, 6).

To determine whether ligating CD19 alters the capacity of mIgM to induce DNA synthesis, replicate samples of human B cells and mitomycin-treated murine L cells that express human Fc_γRII (Fc_γRII–L cells) (7) were cultured for 2.5 days in the presence of incremental concentrations of a monoclonal antibody (MAb) to IgM, a saturating concentration of MAb to CD19, or an equal concentration of an irrelevant MAb, MOPC-21, and interleukin-4 (IL-4). In the absence of anti-CD19, the lowest concentration of anti-IgM that induced incorpo-



ration of [³H]thymidine by B cells above the background incorporation was 6.7×10^{-11} M. In the presence of anti-CD19, this was reduced to 6.7×10^{-13} M (Fig. 1) (8, 9). At higher concentrations of anti-IgM, anti-CD19 also increased the magnitude of [³H]thymidine incorporation to twice that induced by anti-IgM alone. Anti-CD19 alone did not induce proliferation. The ligation of CD19 both lowered the threshold for B cell activation by mIgM and increased the magnitude of the B cell response to optimal levels of mIgM stimulation.

In a parallel experiment to assess the binding characteristics of the monoclonal anti-IgM, replicate samples of B cells and Fc_γRII–L cells were incubated with incremental concentrations of [¹²⁵I]-labeled anti-IgM in the presence or absence of 2×10^{-7} M unlabeled anti-IgM. Cell-bound and free antibody were separated, and specific binding was calculated. The L cells alone did not specifically bind the radiolabeled antibody. Scatchard analysis showed that the anti-IgM bound to the mixture of B cells and L cells with a dissociation constant (K_d) of 1.8×10^{-9} M, and to 2.7×10^5 sites per cell at saturation. The presence of anti-CD19 did not alter the K_d of the anti-IgM for B cells in the presence of Fc_γRII–L cells (10). Therefore, in Fig. 1, the threshold concentration of anti-IgM that induced B cell proliferation in the presence of anti-CD19 bound to only 92 mIgM molecules per cell, or 0.03% of the total mIgM expressed per cell.

We measured the fraction of CD19 that must be ligated to augment B cell proliferation by incubating replicate samples of B cells and Fc_γRII–L cells in the presence of suboptimal anti-mIgM, IL-4, and incremental concentrations of anti-CD19 for 2.5 days, after which incorporation of [³H]thymidine was assayed. The enhancing effect of anti-CD19 increased incrementally from 6.7×10^{-12} M, which bound <0.2% of CD19, up to a saturating concentration of 6.7×10^{-8} M (Fig. 2A). To determine

Fig. 1. Enhancement of DNA synthesis in B lymphocytes by the coaggregation of mIgM and CD19. Replicate samples of 5×10^4 peripheral blood B cells (B) were cultured in 0.2 ml of RPMI-1640 with 10% fetal calf serum in flat-bottom culture plates with 2×10^4 mitomycin-treated fibroblastic L cells expressing human CDw32/Fc_γRII and recombinant IL-4 (200 units/ml, Genzyme). The B cells were stimulated with a range of concentrations of MAb DA4.4 anti-IgM (9) together with either control antibody MOPC-21 (○) or HD37 anti-CD19 (●). Cells were pulsed with [³H]thymidine for the last 16 hours of a 60-hour culture. Results are means \pm SD of triplicates. Cells cultured with either control antibody or anti-CD19 alone incorporated 2726 ± 256 and 2475 ± 255 counts per minute (cpm), respectively. Representative of four experiments.

From *Science*, Vol. 256 (3 April 1992): 105–107 DOI: 10.1126/science.1373518 © 1992 reprinted with permission from AAAS.

whether coaggregation of CD19 with mIgM on the B cells was necessary for augmented cellular proliferation, in replicate samples of cells, F(ab')₂ anti-CD19 was substituted for IgG anti-CD19. The F(ab')₂ form of anti-CD19, which could not interact with the Fc_γR2, suppressed rather than enhanced mIgM-induced B cell proliferation (Fig. 2B). Thus, CD19 augments signaling by mIgM only when the two membrane proteins are coaggregated, probably reflecting a requirement for their proximity.

Anti-CD19 enhanced mIgM-dependent stimulation both in the absence and presence of IL-4 and did not alter the dose response characteristics of the B cells to IL-4. Anti-CD19 also did not induce long-term proliferation of B cells stimulated with anti-IgM or change the time at which peak synthesis occurred (10).

To determine whether ligation of other members of the CD19 complex also enhances mIgM signaling, we incubated B cells with the Fc_γR2-L cells in the presence of saturating concentrations of anti-CD19, anti-TAPA-1, anti-CR2, or control MOPC antibody in combination with IL-4 and incremental concentrations of anti-IgM, and we assessed the cells for [³H]thymidine incorporation after 2.5 days. The anti-CD19 again decreased the concentration of anti-IgM required for B cell proliferation by approximately two orders of magnitude (Fig. 3).




Fig. 2. The requirement for coligation of CD19 with mIgM. B cells were cultured with mitomycin-treated Fc_γR2-L cells and IL-4 (200 units/ml), were stimulated with a range of concentrations of either (A) IgG or (B) F(ab')₂ HD37 anti-CD19 alone (●) or in combination with 10 ng/ml of IgG DA4.4 anti-IgM (○), and were assessed for their capacity to incorporate [³H]thymidine. Representative of three independent replications of the experiment.

Anti-TAPA-1, an antibody selected for its capacity to suppress the growth of a human B cell lymphoma (11), augmented proliferation when coligated with higher concentrations of anti-IgM, and anti-CR2 modestly

Table 1. Comparison of the effect of anti-CD19 with that of antibodies to other B cell surface proteins. B cells were cultured with mitomycin-treated Fc_γR2-L cells, 200 units/ml IL-4, and 5 μg/ml of each IgG1 monoclonal antibody either alone or together with 10 ng/ml of DA4.4 anti-IgM, and they were assessed for incorporation of [³H]thymidine. Anti-IgM at 1.0 μg/ml induced 19,835 ± 732 cpm. Replicate samples of B cells were incubated with the same antibody preparations at 0°C for 30 min, stained with fluorescein-labeled anti-mouse IgG, and analyzed for fluorescence by flow cytometry. The mean fluorescence channel (MFC) for each population is given.

MAb	CD	[³ H]Thymidine incorporation (10 ⁵ cpm)		MFC
		Alone	Anti-IgM added	
None		2.6±0.2	7.7±0.6	42
MHM24	11a	2.3±0.2	7.4±0.3	362
HD37	19	3.9±0.7	22.3±1.4	222
L27	20	2.1±0.8	4.4±0.1	886
4KB128	22	2.5±0.2	7.9±0.7	289
BU38	23	3.0±0.3	9.1±0.3	90
Yz-1	35	2.5±0.1	9.3±0.7	273
BU25p	CI II*	3.1±0.8	13.9±0.8	145

*MHC class II.




Fig. 3. Enhancement of DNA synthesis induced in B cells by the coaggregation of CD19, CR2, or TAPA-1 with mIgM. B cells were cultured with mitomycin-treated Fc_γR2-L cells and IL-4 (200 units/ml) and were stimulated with incremental concentrations of DA4.4 anti-IgM alone (○) or together with 1 μg/ml of HD37 anti-CD19 (●), BU32 anti-CR2 (■), or 5A6 anti-TAPA-1 (▲). Cells cultured without added antibody or stimulated with antibodies to CD19, CR2, or TAPA-1 alone incorporated 2450 ± 155, 2556 ± 180, 2205 ± 103, and 1776 ± 329 cpm, respectively. Representative of three experiments.

augmented the B cell response. The greater activity of anti-CD19 relative to that of anti-TAPA-1 or anti-CR2 suggests that, among the components of the complex, CD19 is the primary mediator of augmented signaling through the mIgM complex. The other components may be less effective because they are reversibly associated with CD19, as demonstrated for CR2, which also forms a complex with CD35 (12).

We assessed the role of CD19 in augmenting mIgM signaling when coligated with the antigen receptor by determining the capacity of saturating concentrations of isotypically matched MAbs specific for six other membrane proteins on B cells to augment cellular proliferation in the presence of limited anti-IgM, Fc_γR2-L cells, and IL-4. Among these antibodies, only anti-MHC class II significantly increased [³H]thymidine incorporation (Table 1). In an additional experiment, anti-MHC class II was examined for its effect on proliferation induced by a range of concentrations of anti-IgM and was found not to decrease the threshold for proliferation in response to anti-IgM (10).

These data show that CD19 participates in a fundamental problem of the immune system, the competing requirements for sensitivity and broad specificity. Coligation of CD19 with mIgM lowered the threshold for effective mIgM signaling so that ligation of only 0.03% of antigen receptors was sufficient to induce cellular proliferation. This function was unique to CD19 and was dependent on coligation with mIgM, linking CD19 with antigen-specific stimulation of the B cell.

This function of CD19 is consistent with its large, conserved cytoplasmic domain and the cocapping of CD19 with mIg (13). The enhancing effect of CD19 also parallels the function of the T cell accessory proteins, CD4 and CD8. Each is a member of the Ig superfamily, each is physically or functionally coupled to a protein tyrosine kinase, and each decreases the threshold for antigen receptor-mediated increases in free intracellular Ca²⁺ concentration and for lymphocyte proliferation when coligated with the antigen receptor; moreover, independent ligation of CD19, CD4, or CD8 suppresses signaling through the antigen receptors of B and T cells (1, 6, 14) (Fig. 2). This analogy predicts that there is a ligand for CD19 to permit complement-independent stimulation of the complex and that an antigen-presenting cell for B cells exists on which both antigen and the CD19 ligand reside.

REFERENCES AND NOTES

1. P. Anderson, M.-L. Blue, C. Morimoto, S.F. Schlossman, *J. Immunol.* **139**, 678 (1987); J. A. Ledbetter *et al.*, *Eur. J. Immunol.* **18**, 525 (1988).
2. L. M. Nadler *et al.*, *J. Immunol.* **131**, 244 (1983); B.

From [Science](#), Vol. 256 (3 April 1992): 105–107 DOI: 10.1126/science.1373518 © 1992 reprinted with permission from AAAS.

- Dörken, P. Möller, A. Pezzutto, R. Schwartz-Albiez, G. Moldenhauer, in *Leucocyte Typing IV: White Cell Differentiation Antigens*, W. Knapp *et al.*, Eds. (Oxford Univ. Press, Oxford, 1989), pp. 34–36.
3. A. K. Matsumoto *et al.*, *J. Exp. Med.* **173**, 55 (1991); L. Bradbury, G. Kansas, S. Levy, R. L. Evans, T. F. Tedder, *FASEB J.* **5**, A611 (1991).
 4. B. Heyman, E. J. Wiersman, T. Kinoshita, *J. Exp. Med.* **172**, 665 (1990); G. Thyphronitis *et al.*, *J. Immunol.* **147**, 224 (1991); T. Hebell, J. M. Ahearn, D. T. Fearon, *Science* **254**, 102 (1991).
 5. R. H. Carter, M. O. Spycher, Y. C. Ng, R. Hoffman, D. T. Fearon, *J. Immunol.* **141**, 457 (1988); H. Yamada, C. H. June, E. D. Anastassiou, F. D. Finkelman, G. C. Tsokos, *FASEB J.* **4**, A1882 (1990).
 6. R. H. Carter, D. A. Tuveson, D. J. Park, S. G. Rhee, D. T. Fearon, *J. Immunol.* **147**, 3663 (1991); A. Pezzutto *et al.*, *ibid.* **138**, 2793 (1987).
 7. J. Banchereau, P. de Paoli, A. Vallé, E. Garcia, F. Rousset, *Science* **251**, 70 (1991). The Fc_γRII-L cells were provided by J. Ravetch [J. V. Ravetch *et al.*, *Science* **234**, 718 (1986)].
 8. B cells were purified by centrifugation over Ficoll-Hypaque (Pharmacia LKB Biotechnology), adherence to plastic in RPMI-1640 medium with 5% calf serum for 60 min at 37°C, treatment with 5 mM leucine methyl ester in RPMI-1640 for 40 min at 37°C, two rounds of rosetting with aminoethylisothiuronium bromide-treated sheep erythrocytes, and then recentrifugation over Ficoll-Hypaque. The resulting population was found to be >80% CD20⁺ and <1% CD3⁺ by flow cytometric analysis.
 9. The IgG1 MAbs used were DA4.4 anti-IgM (American Type Culture Collection), MOPC-21 control antibody (Organon Teknika), HD37 anti-CD19 (Dako, Carpinteria, CA, and G. Moldenhauer), BU32 anti-CR2/CD21 (Binding Site), 5A6 anti-TAPA-1 (17), MHM24 anti-LFA-1/CD11a (Dako), L27 anti-CD20 (Becton-Dickinson), 4KB128 anti-CD22 (Dako), BU36 anti-CD23 (Binding Site), Yz-1 anti-CR1/CD35 [P. S. Changellian, R. M. Jack, L. A. Collins, D. T. Fearon, *J. Immunol.* **134**, 1851 (1985)], and BU25 anti-MHC class II (Binding Site).
 10. R. H. Carter and D. T. Fearon, unpublished observations.
 11. TAPA, target of anti-proliferative antibody [S. Takahashi, C. Doss, S. Levy, R. Levy, *J. Immunol.* **145**, 2207 (1990)].
 12. D. A. Tuveson, J. M. Ahearn, A. K. Matsumoto, D. T. Fearon, *J. Exp. Med.* **173**, 1083 (1991).
 13. I. Stamenkovic and B. Seed, *ibid.* **168**, 1205 (1988); L.-J. Zhou, D. C. Ord, A. L. Hughes, T. F. Tedder, *ibid.* **174**, 1424 (1991); J. M. Pesando, L. S. Bouchard, B. E. McMaster, *ibid.* **170**, 2159 (1989).
 14. I. Bank and L. Chess, *ibid.* **162**, 1294 (1985); C. E. Rudd, J. M. Trevisan, J. D. Dasgupta, L. L. Wong, S. F. Schlossman, *Proc. Natl. Acad. Sci. U.S.A.* **85**, 5190 (1988); A. Veillette, M. A. Bookman, E. M. Horak, J. B. Bolen, *Cell* **55**, 301 (1988).
 15. We thank G. Moldenhauer and S. Levy for providing antibody and J. Ravetch for providing Fc_γRII-L cells. Supported by NIH grants AI-22833 and AI-28191 and a Terri Gotthelf Lupus Research Foundation award (R.H.C.).

2 December 1991; accepted 19 February 1992