

Structural Basis for Fc γ RIIa Recognition of Human IgG and Formation of Inflammatory Signaling Complexes

This information is current as of May 25, 2019.

Paul A. Ramsland, William Farrugia, Tessa M. Bradford, Caroline Tan Sardjono, Sandra Esparon, Halina M. Trist, Maree S. Powell, Peck Szee Tan, Angela C. Cendron, Bruce D. Wines, Andrew M. Scott and P. Mark Hogarth

J Immunol 2011; 187:3208-3217; Prepublished online 19 August 2011;
doi: 10.4049/jimmunol.1101467
<http://www.jimmunol.org/content/187/6/3208>

Supplementary Material <http://www.jimmunol.org/content/suppl/2011/08/19/jimmunol.1101467.DC1>

References This article **cites 61 articles**, 20 of which you can access for free at:
<http://www.jimmunol.org/content/187/6/3208.full#ref-list-1>

Why *The JI*? [Submit online.](#)

- **Rapid Reviews! 30 days*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

**average*

Subscription Information about subscribing to *The Journal of Immunology* is online at:
<http://jimmunol.org/subscription>

Permissions Submit copyright permission requests at:
<http://www.aai.org/About/Publications/JI/copyright.html>

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://jimmunol.org/alerts>

Structural Basis for Fc γ RIIa Recognition of Human IgG and Formation of Inflammatory Signaling Complexes

Paul A. Ramsland,^{*,†,‡} William Farrugia,^{*} Tessa M. Bradford,^{*} Caroline Tan Sardjono,^{*} Sandra Esparon,^{*} Halina M. Trist,^{*} Maree S. Powell,^{*,‡,§} Peck Szee Tan,^{*} Angela C. Cendron,^{*} Bruce D. Wines,^{*,‡,§} Andrew M. Scott,[¶] and P. Mark Hogarth^{*,‡,§}

The interaction of Abs with their specific FcRs is of primary importance in host immune effector systems involved in infection and inflammation, and are the target for immune evasion by pathogens. Fc γ RIIa is a unique and the most widespread activating FcR in humans that through avid binding of immune complexes potently triggers inflammation. Polymorphisms of Fc γ RIIa (high responder/low responder [HR/LR]) are linked to susceptibility to infections, autoimmune diseases, and the efficacy of therapeutic Abs. In this article, we define the three-dimensional structure of the complex between the HR (arginine, R134) allele of Fc γ RIIa (Fc γ RIIa-HR) and the Fc region of a humanized IgG1 Ab, hu3S193. The structure suggests how the HR/LR polymorphism may influence Fc γ RIIa interactions with different IgG subclasses and glycoforms. In addition, mutagenesis defined the basis of the epitopes detected by FcR blocking mAbs specific for Fc γ RIIa (IV.3), Fc γ RIIb (X63-21), and a pan Fc γ RII Ab (8.7). The epitopes detected by these Abs are distinct, but all overlap with residues defined by crystallography to contact IgG. Finally, crystal structures of LR (histidine, H134) allele of Fc γ RIIa and Fc γ RIIa-HR reveal two distinct receptor dimers that may represent quaternary states on the cell surface. A model is presented whereby a dimer of Fc γ RIIa-HR binds Ag-Ab complexes in an arrangement that possibly occurs on the cell membrane as part of a larger signaling assembly. *The Journal of Immunology*, 2011, 187: 3208–3217.

The interaction between Ig complexes and FcRs induces potent and diverse immune responses. In normal immunity, these include inflammation, Ab-dependent killing of target cells, mast cell degranulation, phagocytosis, and regulation of Ag receptor activation of B cells. However, in pathological situations like autoimmunity, immune complex FcR-mediated activation of effector cells is a major pathway in the development of tissue injury and, indeed, the early events of disease pathogenesis (1, 2).

The Fc γ RII receptors (CD32 group of receptors) are key activating and inhibitory effectors of the IgG-mediated immune functions of leukocytes. Fc γ RIIa triggers both host protective and damaging proinflammatory activities, whereas Fc γ RIIb modulates signaling from the activating Fc γ Rs, including Fc γ RIIa, and the B-cell Ag receptor complex. The evidence of many models of autoimmune disease and the association of receptor gene polymorphisms with human disease concur that the balanced integration of signals from activating receptors and inhibitory Fc γ RIIb is necessary for a normal host response to infection and resistance to autoimmune pathologies. Abs that specifically target these receptors to manipulate this balance of proinflammatory and inhibitory signals are of particular interest as therapeutics (3).

Fc γ RIIa is unique to higher primates and is the most widespread FcR. This low-affinity FcR is different from all other activating FcRs because its signaling ITAM is contained in the ligand-binding chain. Other activating FcRs and related multisubunit immunoreceptors, including the Ag receptors and NK receptors, signal via their noncovalent association with dimeric accessory molecules such as the common FcR- γ -chain dimer that contain ITAMs (4, 5). Animal and human studies indicate that Fc γ RIIa plays a major role in the development of destructive inflammation. Mice transgenic for human Fc γ RIIa exhibit extreme sensitivity to pathogenic Abs and also develop a spontaneous autoimmune disease with features of human rheumatoid arthritis including joint destruction (6) and are sensitive to thrombocytopenia-inducing Abs (7). The transgenic Fc γ RIIa mice are particularly susceptible to collagen-induced arthritis, which can be largely suppressed by small chemical inhibitors designed to bind Fc γ RIIa (8). In addition, *in vitro* studies of human plasmacytoid dendritic cells from systemic lupus erythematosus (SLE) patients show the targeting of DNA to TLR9 is dependent on Ab and Fc γ RIIa, as opposed to other FcRs (9).

^{*}Centre for Immunology, Burnet Institute, Melbourne, Victoria 3004, Australia; [†]Department of Surgery, Austin Hospital, University of Melbourne, Heidelberg, Victoria 3084, Australia; [‡]Department of Immunology, Monash University, Melbourne, Victoria 3004, Australia; [§]Department of Pathology, University of Melbourne, Parkville, Victoria 3056, Australia; and [¶]Tumour Targeting Program, Ludwig Institute for Cancer Research, Austin Health, Heidelberg, Victoria 3084, Australia

Received for publication May 18, 2011. Accepted for publication July 6, 2011.

This work was supported by project Grant 315625 from the National Health and Medical Research Council of Australia and the Victorian state government Operational Infrastructure Support Scheme. P.A.R. was supported by a National Health and Medical Research Council R. Douglas Wright Career Development award (Grant 365209). This work, including use of the BioCARS sector, was supported by the Australian Synchrotron Research Program, which is supported by the Commonwealth of Australia under the Major National Research Facilities Programme. Use of BioCARS Sector 14 was also supported by the National Institutes of Health, National Center for Research Resources (Grant RR07707). Use of the Advanced Photon Source was supported by the U.S. Department of Energy, Basic Energy Sciences, Office of Energy Research (Contract W-31-109-Eng-38).

Address correspondence and reprint requests to Prof. P. Mark Hogarth, Centre for Immunology, Burnet Institute, GPO Box 2284, Melbourne, Victoria 3001, Australia. E-mail address: pmhogarth@burnet.edu.au

The online version of this article contains supplemental material.

Abbreviations used in this article: Fc γ RIIa-HR, high-responder (arginine, R134) allele of Fc γ RIIa; Fc γ RIIa-LR, low-responder (histidine, H134) allele of Fc γ RIIa; GlcNAc, *N*-acetylglucosamine; HR, high responder; LR, low responder; PDB, Protein Data Bank; PEG, polyethylene glycol; SLE, systemic lupus erythematosus.

Copyright © 2011 by The American Association of Immunologists, Inc. 0022-1767/11/\$16.00

Genetic polymorphisms of Fc γ RIIa have also been linked to susceptibility to several autoimmune diseases, notably SLE (10–13) and ulcerative colitis (14), as well as resistance to Gram-negative bacterial infection (15) and to the outcome of therapeutic Ab treatment in lymphoma (16). The most extensively studied polymorphism is the high-responder/low-responder (HR/LR) polymorphism, alleles of which code either arginine (HR) or histidine (LR) at position 134 [numbering is based on the experimentally determined N-terminal sequence (17); this polymorphic site is frequently referred to as position 131 in the amino acid sequence (18, 19)]. The functional differences between the HR (arginine, R134) allele of Fc γ RIIa (Fc γ RIIa-HR) and LR (histidine, H134) allele of Fc γ RIIa (Fc γ RIIa-LR) relate to different abilities to bind mouse IgG1 or human IgG2, respectively (19, 20). Indeed, the Fc γ RIIa-LR is the only receptor that binds human IgG2 (21), which interestingly is a major IgG class in autoimmunity (11) and in resistance to Gram-negative bacterial infection (15) and severe swine flu infection (22). A higher frequency of Fc γ RIIa-LR homozygous genotype was found to be associated with *Plasmodium falciparum* malaria resistance and consistently higher levels of anti-malarial IgG2 and IgG3 serum Ab (23).

Despite the extensive functional, genetic, and biochemical characterization of this receptor, the three-dimensional structural basis for the interaction of Fc γ RIIa with IgG and functional differences of the HR/LR polymorphism was unknown. Soluble Fc γ RIIa-LR glycoprotein can form a crystallographic dimer as described for orthorhombic $P2_12_12$ crystals (24). Such a dimer of cellular receptors would place the two ITAM-containing cytoplasmic tails in close proximity, similar to other ITAM-activating molecules, in an arrangement presumably suitable for downstream intracellular signaling (25). It is generally accepted that, in activating leukocytes, the FcRs cluster or aggregate, a process that is driven by Ag–Ab complexes (26–28), yet the structural arrangement of receptors within these signaling assemblies is poorly understood.

In this article, the resolution of the crystal structure of Fc γ RIIa-LR is extended to 1.5 Å from orthorhombic $P2_12_12$ crystals (24). In addition, we have determined the 2.3-Å resolution structure of glycosylated Fc γ RIIa-HR from orthorhombic $C222_1$ crystals and show it forms a crystallographic dimer distinct from that of Fc γ RIIa-LR. We also report, to our knowledge, the first three-dimensional structure for the complex of Fc γ RIIa-HR and the Fc region of human IgG1 at a resolution of 3.8 Å. Interestingly, this Fc γ RIIa–Fc complex could itself form a dimer equivalent to the crystallographic dimer of Fc γ RIIa-HR (unlike Fc γ RIIa-LR) without interfering with ligand binding. Hence we predict signaling by Fc γ RIIa occurs through dimeric receptors with bound immune complexes (Ag–IgG), possibly as part of larger aggregates, which are important for intracellular signaling, cell activation, and ultimately inflammation.

Materials and Methods

Production and purification of recombinant human Fc γ RIIa and the IgG1 Fc

Fc γ RIIa-LR (H134) was prepared as previously described; a point mutation replacing S88 with F88 introduced during the production of soluble Fc γ RIIa-LR did not alter its ability to bind IgG or immune complexes (17). Fc γ RIIa-HR (R134) cDNA was produced by splice overlap extension PCR and expressed using a baculovirus system in SF21 insect cells using Life Technologies SF900 media (Invitrogen Australia Pty, Rowville, VIC, Australia) at 27°C for 72 h. The receptor was purified from supernatant by anion exchange over Q-Sepharose (GE Healthcare, Rydalmere, NSW, Australia), followed by affinity chromatography on heat-aggregated Ig (Novartis, Basel, Switzerland) coupled Sepharose, as previously described for Fc γ RIIa-LR (17). For crystallization, Fc γ RIIa-HR was prepared in 75 mM NaCl, 5 mM Tris-HCl pH 7.4 at between 5 and 10 mg/ml.

The Fc portion was released by plasmin proteolysis from a humanized IgG1, hu3S193, which was expressed in an NSO myeloma cell line (29). Purified recombinant hu3S193 at 5 mg/ml was dialyzed against 0.15 M NaCl, 50 mM Tris pH 8.0, and was digested with 1 U/ml plasmin (Roche, Basel, Switzerland) for 48 h at 21°C. The Fc was purified to homogeneity by separation from Fab by protein A-Sepharose, followed by removal of residual intact IgG by protein L-Sepharose (Pierce Protein Research Products, Thermo Fisher Scientific, Scoresby, VIC, Australia). Purified Fc was dialyzed against 75 mM NaCl, 5 mM Tris buffer pH 7.4 before crystallization.

Crystallization of Fc γ RIIa and complexes with the Fc of IgG1

Crystals of Fc γ RIIa-LR were produced in 5- μ l hanging drops with the protein solution at 8 mg/ml in 0.2 M ammonium acetate, 0.1 M sodium citrate buffer pH 5.6, and 30% (w/v) polyethylene glycol (PEG) 4000. Crystals were nucleated at 37°C (typically 5–7 d for small crystals to appear), and were grown and maintained at 18°C. Fc γ RIIa-HR crystals were prepared and maintained at 18°C in 2 μ l sitting droplets, with the protein at 4 mg/ml in 30% (w/v) PEG 4000 and 0.2 M ammonium sulfate.

Cocrystallization of IgG1–Fc was successful with both polymorphic variants of Fc γ RIIa, but only with wild-type Fc γ RIIa-HR were the crystals suitable for structure determination. However, both systems produced large, thin plates with growth habits of single crystals and crystalline clusters in 2 μ l sitting and hanging droplets in 2–4 wk at 18°C. Typically, 1 μ l protein solution (containing 5–8 mg/ml total protein with Fc γ RIIa and Fc mixed at a 1:1 molar ratio) was mixed with 1 μ l of the reservoir solution that contained 0.2 M trisodium citrate or tripotassium citrate and 20% PEG 3350. A range of additives was tested with the best crystals of the complex grown from an initial protein concentration of 5.5 mg/ml in 5 mM Tris pH 7.4, 0.15 M NaCl, 2 mM MOPS, 0.2 M tripotassium citrate, and 20% PEG 3350. Supplies and reagents for crystallization were purchased from Hampton Research (Aliso Viejo, CA).

Data collection and structure determinations

For Fc γ RIIa-LR, a large (0.6 \times 0.2 \times 0.2 mm) orthorhombic $P2_12_12$ crystal was briefly transferred into artificial mother liquor containing 20% (v/v) glycerol before flash-cooling to 100 K in a nitrogen cryostat (Cryojet; Oxford Instruments, Abingdon, U.K.). Diffraction data to 1.5-Å resolution were obtained at 100 K using synchrotron radiation on beamline 14-BMC (BioCARS) at the Advanced Photon Source and processed with the HKL package, version 1.97 (30). Initial crystallographic refinement involved rigid body fitting of domains 1 and 2 (residues 4–88 and 89–173, respectively) of LR-Fc γ RIIa using the coordinates of Protein Data Bank (PDB) entry 1FCG (24). Subsequent cycles of interactive model building and solvent fitting into σ_A -weighted $2F_o - F_c$ and $F_o - F_c$ electron density maps were performed in TURBO-FRODO version 5.5 (BioGraphics, Marseille, France). Each cycle of model building was followed by simulated annealing, energy minimization, and individual temperature (B) factor refinement protocols against maximum likelihood target functions in the CNS version 1.0 program suite (31). Bulk solvent correction, anisotropic overall B factor refinements, and cross-validation by assigning 10% of intensities for calculation of an R_{free} value were applied throughout the structure determination. Relevant values and statistics for evaluation of the X-ray diffraction data and the final atomic model are presented in Table I.

Crystals of Fc γ RIIa-HR were extremely sensitive to glycerol, but crystals were relatively stable in artificial mother liquor containing 20% (v/v) PEG 400. Using this cryoprotectant, we used an orthorhombic $C222_1$ crystal (0.2 \times 0.1 \times 0.1 mm) to collect X-ray diffraction data at 100 K on a MicroMax007/R-Axis IV²⁺ rotating anode generator system (Rigaku Americas, The Woodlands, TX), operated at 40 kV and 20 mA. Processing of X-ray diffraction data and crystallographic refinement to a resolution of 2.3 Å proceeded essentially as described earlier (Table I), but in this case, the coordinates of the high-resolution structure of Fc γ RIIa-LR were used as a starting template for molecular replacement. Two residues were replaced (F88 with S88 and H134 with R134) to correspond with the sequence of the wild-type and HR polymorphism of Fc γ RIIa, respectively.

Crystals of the Fc γ RIIa–IgG1–Fc complex proved extremely sensitive to cryopreservation, with evident large increases in mosaic spread and uneven shrinkage of the unit cell such that different cryopreserved crystals were generally nonisomorphous. Moreover, the volumes of the crystals that were essentially thin plates (<10 μ m in the smallest dimension) were not suitable for room temperature data collection. Our best crystals were flash-cooled in artificial mother liquor in the presence of 20% (v/v) glycerol and diffracted X-rays anisotropically to between 3.4 and 4.5 Å resolutions. Consequently, the structure determination extended to a resolution of 3.8 Å and was solved by molecular replacement using the coordinates of the unliganded form of Fc γ RIIa-HR (described in this

Table I. Summary of data collection and crystallographic refinements

	Fc γ RIIa-LR	Fc γ RIIa-HR	Fc γ RIIa-HR-Fc
Data collection			
Space group	$P2_12_12$	$C222_1$	$C222_1$
Cell dimensions <i>a</i> , <i>b</i> , <i>c</i> (Å)	79.34, 100.41, 27.94	50.04, 77.94, 88.10	153.38, 255.55, 58.44
Resolution (Å)	20–1.5 (1.55–1.5) ^a	30–2.3 (2.38–2.3)	30–3.8 (3.94–3.8)
R_{sym}	0.052 (0.32)	0.10 (0.27)	0.14 (0.35)
Mean $I/\sigma(I)$ ^a	21.8 (3.8)	13.2 (4.9)	10.8 (3.78)
Completeness (%)	94.5 (85.5)	97.2 (96.0)	98.9 (99.7)
Redundancy	6.1 (4.0)	4.6 (4.3)	5.0 (4.8)
Refinement			
Resolution (Å)	19.8–1.5	20.4–2.3	29.7–3.8
No. of reflections	34,590	7,554	11,110
$R_{\text{work}}/R_{\text{free}}$	0.20/0.23	0.21/0.28	0.25/0.42
No. of atoms			
Protein	1,354	1,350	4,745
Ligand	14	0	376
Water	273	85	24
<i>B</i> -factors			
Protein	22.7	20.7	34.5
Water	36.5	23.1	26.6
RMSD bond lengths (Å)	0.005	0.006	0.010
RMSD bond angles (°)	1.3	1.4	1.8

^aValues in parentheses are for the highest resolution shell. RMSD, root-mean-square deviation.

article) and the Fc from PDB entry 1E4K (32). Crystals contained one Fc γ RIIa-HR-IgG1-Fc complex in the asymmetric unit. Crystallographic refinements were performed against a crystallographic residual target as described earlier, but only group and overall B factors were refined (Table I). The high R_{free} and below average quality of the model geometry is due to the anisotropy and low resolution of the diffraction data, but in previous studies with glycosylated Fc molecules free and in complexes, similar high R_{free} values and diffraction properties have been reported, even when data extend to higher resolutions (32). However, electron density maps were of sufficient quality for tracing the main chains for the Fc γ RIIa-HR monomer and the two H chains of the Fc of hu3S193 IgG1, and placement of 29 saccharide units at four N-linked glycosylation sites. Atomic coordinates and structure factors have been deposited in the PDB (PDB ID codes 3RY4, 3RY5, and 3RY6; <http://www.pdb.org>).

mAb and epitope mapping

The anti-Fc γ RIIa mAbs 8.7 (murine IgG1) (33) and IV.3 have been previously described (34). The anti-Fc γ RIIb mAb X63-21/7.2 was made by immunizing transgenic mice expressing the human Fc γ RIIa with mouse IIA1.6 lymphoma cells transfected with, and expressing, human Fc γ RIIb2. Mice were immunized on three occasions with 10^6 cells, boosted i.v. with 10^6 cells, and splenocytes fused with NS1 cells. Hybridoma cell culture supernatants were harvested from cloned X63-21/7.2 cells and IgG purified by affinity chromatography as described. Fab and F(ab')₂ fragments were generated by pepsin (Boehringer Mannheim, Mannheim, Germany) digestion at an enzyme/substrate ratio of 1:10 to 1:50 at 37°C for 1 h in 0.1 M sodium citrate pH 3.5 as described previously (33). Ab fragments were biotinylated with EZ-link biotin (Pierce Protein Research Products, Thermo Scientific). Fc γ RIIa-HR (R134) mutant constructs or with equivalent sequence elements exchanged between Fc γ RIIa and the IgE receptor, Fc ϵ R1, were produced as described previously (5, 35) and expressed by transient transfection of COS-7 or CHOP cells using Fugene-6 (Boehringer Mannheim) or Lipofectamine 2000 (Invitrogen Australia Pty). Transfected cells were reacted with unbiotinylated or biotinylated mAb fragments and further reacted with FITC-conjugated anti-mouse Ig or R-phycoerythrin-conjugated streptavidin as appropriate and analyzed by flow cytometry.

Results

Crystal structures of the Fc γ RIIa-HR and Fc γ RIIa-LR

The two allelic variants of Fc γ RIIa were produced in insect cells and purified to homogeneity. Surface plasmon resonance established equilibrium dissociation constants (K_D) for solid-phase human IgG1 of 0.5 ± 0.02 mM and 0.6 ± 0.03 mM for the HR and LR versions of Fc γ RIIa, respectively. This is consistent with

affinities measured in previous studies of Fc γ RIIa (17) and for Fc γ RIIb (36), and is also consistent with the observed binding of IgG to whole cells (35). Thus, the relatively simple glycosylation pattern of the insect cells did not appear to significantly influence IgG binding activity of each of these forms. Higher resolution information was obtained for Fc γ RIIa-LR in this study (Fig. 1A, 1B), and the glycosylated HR form (Fig. 1C, 1D) was analyzed for the first time to our knowledge and in complex with the Fc of human IgG1 (Figs. 2, 3). The Fc γ RIIa-HR protein crystallized in the orthorhombic $C222_1$ space group (Table I), which is distinct from the orthorhombic $P2_12_12$ crystals of the LR form previously described (17, 24). Analysis of the crystal lattice to 2.3-Å resolution identified a dimer of Fc γ RIIa-HR that is markedly distinct from the LR dimer (Fig. 1) (24). The Fc γ RIIa-HR dimer interface consists of a total buried surface area of 1564 Å² and involves regions of both extracellular Ig-like domains (D1 and D2) of the receptor monomers. Interactions in the Fc γ RIIa-HR dimer occur principally between the A'B strands and loop, as well as the E




FIGURE 1. Surface representations of the two distinct crystallographic dimers of Fc γ RIIa. *A*, Side view of the Fc γ RIIa-LR dimer with monomer 1 in cyan, monomer 2 in magenta, and the GlcNAc residue attached to N145 shown in green. *B*, End-on view of the Fc γ RIIa-LR dimer, looking down onto the surfaces involved in IgG binding. *C*, Side view of the Fc γ RIIa-HR dimer. *D*, End-on view of the Fc γ RIIa-HR dimer.




FIGURE 2. Overview of the Fc γ RIIa-HR complex with Fc of human IgG1. *A*, Surface representation with Fc γ RIIa in white (N-linked oligosaccharides at positions N64 and N145 are in green) and the Fc in magenta (chain A) and cyan (chain B) for the two H chains. Oligosaccharides attached to N297 of the Fc are in orange. *B*, Surface view of the Fc γ RIIa-Fc complex rotated by 90° from its orientation in *A*. *C*, α -Carbon polypeptide trace of the Fc γ RIIa-Fc complex with the N-linked oligosaccharide chains displayed as space-filling (CPK) representations. *D*, Close-up of the Fc γ RIIa interaction with Fc. The extracellular domains 1 and 2 (D1 and D2) and the F-G loop of Fc γ RIIa are indicated, as are the constant domains, CH2 and CH3, of the H chains.

strand of D2 and the A'B and E strands of D1; notably, residues T26, R33, Q54, P55, S56, R58, E102, G103, T105, P142, and Q143 contribute to the receptor pairing. Importantly, the Fc γ RIIa-HR dimer interface is distinct from the Fc γ RIIa-LR interface that involves residues only of D2 of each monomer and does not overlap. As previously described, the Fc γ RIIa-LR interface involves residues from the C' strands interacting with the F and G strands of neighboring monomers (24).

The second major distinction between the HR and LR crystallographic forms is the relationship between the IgG binding surfaces and the dimer interfaces. Mutagenesis studies of Fc γ RIIa (24, 37) combined with homology modeling predict that the IgG binding surface involves residues from the A strands, the BC, C'E, and FG loops, which are not within the LR or HR dimer interfaces. Thus, for both dimers, the predicted IgG binding surfaces are accessible to solvent if the receptors are oriented in a hypothetical cell membrane (Fig. 1). However, in the LR dimer, the IgG binding surfaces of each monomer are "closely" apposed so that access of the IgG ligand for binding one receptor monomer is prevented by the neighboring monomer, whereas in the HR dimer, the relative position of the binding surfaces is more "open" and are predicted to allow access for binding of one or two IgG ligands.

Although both the LR and HR forms of Fc γ RIIa were glycosylated, we observed electron density corresponding to only a single *N*-acetylglucosamine (GlcNAc) moiety attached to N145 in the Fc γ RIIa-LR form of the receptor at a resolution of 1.5 Å




FIGURE 3. Sites of interaction between Fc γ RIIa and human IgG1 revealed by the crystal structure of the Fc γ RIIa-HR complex with Fc. *A*, Binding site on FcR for its Fc ligand with residues that interact with Fc shaded on the solvent-accessible surface. The N-linked glycans at positions 64 and 145 are also included, but these do not participate directly in the interaction with Fc. *B*, Close-up of the FcR residues involved in the interaction with Fc. Note the site of interaction of the IgG fucose of N297 (S129 and K128) is shown in orange. *C*, Solvent-accessible view of the Fc ligand with the residues in contact with FcR shaded. Note the polymorphic R134 residue of Fc γ RIIa-HR binds in a shallow depression on the Fc formed by D265, V266, S267, and N297. *D*, Close-up of the Fc residues in contact with FcR. Numbering for the Fc residues is according to the Eu Index (62). The coloring scheme is the same as Fig. 2 with contacts shaded according to the Fc ligand.

(Fig. 1). The carbohydrates were disordered at N64 in both allelic forms used in these studies. However, we have observed carbohydrates at both N-linked sites in a heavily glycosylated sample of Fc γ RIIa-LR, but these crystals diffracted to a resolution of only 2.8 Å. In this glycoform, a GlcNAc[α 1 \rightarrow 6Fuc] β 1 \rightarrow 4GlcNAc trisaccharide was clearly visible at the N145 site, whereas a single GlcNAc was attached to N64 (P.A.R., W.F., and P.M.H., unpublished observations). We have attributed the decreased X-ray diffraction quality of crystals of the second Fc γ RIIa-LR glycoform to the oligosaccharide at position 145, which comes into contact with a symmetry-related molecule. Notably, in the two distinct dimers of Fc γ RIIa, the glycosylation sites are not involved in the interface region (Fig. 1). Thus, the carbohydrates may act to limit lateral interactions with other receptors in crystals and in their native context in membranes.

Overview of the Fc γ RIIa-HR complex with Fc of human IgG1

The 3.8-Å resolution crystal structure (Figs. 2, 3) was determined for Fc γ RIIa-HR bound to Fc of human IgG1. Fc γ RIIa bound its Fc ligand in an asymmetric fashion between the two CH2 domains and the lower hinge regions (Fig. 2), which is similar to Fc γ RIII (CD16) (32, 38), and consistent with mutagenesis of Fc and functional studies of IgG (39). The protruding loop connecting the F and G β -strands in D2 of Fc γ RIIa penetrates, with Y160 in the lead, between the lower hinge regions and the first β -strands of the two CH2 domains (Fig. 2C, 2D). As a result, the hinge regions were pushed from a central location and now lie almost perpendicular to the pseudo-2-fold axis of the Fc and are directed away from the Fc γ RIIa binding site.

Rather than a contiguous binding site, the binding surfaces of Fc γ RIIa that recognize Fc are composed of three separate regions

(Fig. 3). First, a large, protruding ridge is formed by the tip of the F-G loop (Y160), the neighboring B-C loop (P117, V119, and K120), and the C' β-strand (F132, R134, and L135; Fig. 3A, 3B). This ridge participates in the majority of binding interactions with the Fc ligand by fitting between the two lower hinge regions (Leu-Leu-Gly-Gly; positions 234–237) and binding to a large area (S239, D265, S267, V266, and N297) on one side of the Fc (chain B; Fig. 3C, 3D). Second, a small, yet important, binding site is formed by FcγRIIa residues W90 and W113 that trap P329 from Fc chain A (Fig. 3B). Such an arrangement has been observed in complexes of FcεRI and FcγRIII (32, 38, 40), and is almost certainly a feature of all conventional FcγRs. The Pro residue binding in this alcove on FcγRIIa is probably mimicked by a peptide that was selected by phage-display and blocks IgG binding (41). The third FcγRIIa binding region involves a small patch (S129 and K128), which contacts the Fc-associated carbohydrate of chain B (N297-glycan) and may influence recognition of various IgG glycoforms (see later).

Influence of carbohydrate on the FcγRIIa interaction with human IgG1

A considerable amount of carbohydrate was observed in the electron density maps calculated for the FcγRIIa-Fc complex (Fig. 2). Unlike the unliganded FcγRIIa-HR that had a notable absence of ordered carbohydrate (Fig. 1), the same receptor in complex with Fc displayed an ordered branched oligosaccharide chain at N145 and a clearly defined GlcNAc residue at N64 (Fig. 2). Both FcγRIIa N-linked carbohydrates are distant from the binding interaction with the Fc ligand.

Carbohydrates attached to N297 of the two H chains reside at the interface of the CH2 domains and are consequently ordered in most crystal structures of Fc and Fc-ligand complexes (32, 42, 43). In the hu3S193 Fc complex with FcγRIIa, we observed biantennary and fucosylated oligosaccharide on both H chains (Fig. 2). The oligosaccharide attached to H chain A was missing only the terminal disaccharide Neu5Acα2→6Gal of the Manα1→6 branch. In H chain B, a complete branched N-linked glycan, including the two terminal sialic acids, was observed. A bisecting GlcNAc residue was not observed for either Fc-associated glycan, which is not surprising because this moiety occurs in only 5–10% of normal human IgG.

Because of the asymmetric nature of the FcγRIIa-Fc interaction, only the carbohydrate associated with one H chain (B) is directly involved in the interaction through the proximity of the Fuc to S129 and K128 of the receptor (Figs. 2, 3). Interestingly, the F132 and R134 side chains of FcγRIIa-HR bind in a shallow depression on Fc (chain B), formed by D265, V266, S267, and N297, which is adjacent to the Fuc moiety. The binding of R134 near the Fuc moiety of the IgG-Fc may relate to the previously reported modest increase in affinity of FcγRIIa-HR for Fuc-deficient IgG1 (44). In addition, the structure provides a glimpse of the interactions with IgG involving the HR/LR (R134/H134) polymorphic site, which has been linked to disease susceptibility (45), and in H134/H134 individuals higher response rates to a therapeutic Ab, rituximab (16).

Structural comparison of FcγR interactions with different IgG subclasses

Amino acids involved in the FcγRIIa-Fc interaction were assessed for their role in determining the specificity of different FcRs and the variation in IgG subclass binding by FcγRIIa (Table II). The related FcγRIII and FcγRI that both bind IgG displayed low sequence identities (5/11 and 6/11, respectively) to the FcγRIIa binding-site residues, which may explain the unique specificity of

Table II. Comparison of residues of the FcγRIIa-IgG1-Fc interaction with those of other human FcR and IgG subclasses

Position ^a	FcγRIIa ^b	FcγRIII	FcγRI	FcεRI
Human FcRs				
90	W	W	W	W
113	W	W	W	W
117	P	A	I	D
119	V	H	V	V
120	K	K	N	K
128	K	K	K	F
129	S	G	A	A
132	F	Y	F	Y
134	H/R (R)	H	H	Y
135	L (S)	H	W	F
160	Y	S	K	Q
Eu Index ^c				
	IgG1	IgG2	IgG3	IgG4
Human IgG subclasses				
234	L	V	L	F
235	L	A	L	L
236	G	–	G	G
237	G	G	G	G
239	S	S	S	S
265	D	D	D	D
266	V	V	V	V
267	S	S	S	S
297	N	N	N	N
327	A	G	A	G
328	L	L	L	L
329	P	P	P	P

^aPosition in amino acid sequence corresponding to the numbering scheme for crystal structures described in this article and reported previously (17, 24).

^bResidues in parentheses occur in FcγRIIb. The same numbering scheme has been used for FcγRIIb.

^cNumbering for IgG is the Eu Index (62).

–, deletion in the sequence.

FcγRIIa for various IgG subclasses. In addition, the related IgG inhibitory receptor, FcγRIIb, has R134 similar to the HR polymorphism, but differs by having S135 replacing L135 of FcγRIIa (Table II). It was not surprising that the IgE receptor (FcεRI) shared only 4/11 binding site residues used by FcγRIIa.

The HR and LR alleles of FcγRIIa were initially named for their ability to bind mouse IgG1. However, the FcγRIIa-LR reacts efficiently with human IgG2, whereas the FcγRIIa-HR does not (21), and the relative reactivity of FcγRIIa toward human IgG subclasses is reported to be: IgG3 > IgG1/IgG2 > > IgG4 for LR-FcγRIIa (17) and IgG3/IgG1 > > IgG2 > IgG4 for FcγRIIa-HR (21). The near-equivalent binding of human IgG3 and human IgG1 is reflected in the identity of the Fc residues involved in the interaction with FcγRIIa-HR (Table II). It is likely that differences in affinity of FcγRIIa for the IgG subclasses are also contributed by different sequences and lengths of the hinge region. In IgG2, 8/12 (including a deletion at position 236 with respect to IgG1) contact residues were identical, whereas in IgG4, 10/12 residues were identical to IgG1. Interestingly, most of the differences in human IgG1 versus IgG2 are located near the R134/H134 polymorphic binding site, suggesting that cooperative conformational effects are responsible for the different binding capacities of the HR and LR receptor alleles toward the human IgG2 subclass. The LL to VA (234–235) substitutions and the deletion of G236 in the “lower hinge” region of IgG2 is where the F-G loop and Y160 residue of the receptor binds, which is immediately adjacent to the cavity where R134 is located. In contrast, the A to G (327) substitution in IgG2 is close to residue P329, which on the other side of the Fc is sandwiched between W90 and W113 of FcγRIIa-HR. Thus, the noted differences in IgG2 in the lower hinge may ac-

count for specificity of this subclass for the Fc γ RIIa-LR (also see later). It is noteworthy that like human IgG, the mouse IgG classes also vary in hinge sequence and length; in particular, mouse IgG1 originally used to distinguish between the HR and LR phenotypes contains a truncated lower hinge region, which is two residues shorter than in human IgG1 (L234 and L235 are absent), whereas 6/10 remaining Fc γ RIIa contact residues in mouse IgG1 are identical (S239, D265, S267, N297, A327, and P329). Thus, the sequence and length of the lower hinge appears to be important in dictating the strength of binding to the polymorphic variants of Fc γ RIIa.

The interaction of Fc γ RIIa and IgG is distinct from that of Fc γ RIII (CD16a) with IgG. The Fc γ RIII-IgG complex has been defined and predictions made for the interactions of human IgG1 with Fc γ RIIa-HR and Fc γ RI (32, 46). In Fc γ RIII, the binding site for Fc residue L235 of chain A involves residues T113, A114, and V155, and on this basis, the equivalent residues in Fc γ RIIa (K116, P117, and I158) were predicted to bind IgG (46). However, in the crystal structure of Fc γ RIIa-HR-Fc, whereas L235 on Fc binds close to the predicted binding site, only P117 of the receptor contacts the Fc ligand (Fig. 3B). Furthermore, L235 on the other H chain (B) in the Fc γ RIII-IgG complex is bound between Fc γ RIII residues H116 and H132 that correspond to Fc γ RIIa residues V119 and L135, which were predicted to bring L235 closer into hydrophobic contact with the receptor (46). In contrast, wedging of the Fc γ RIIa F-G loop into the lower hinge region where Y160 interacts with L234 (of H chain B) forces L235 of the Fc away (>12 Å) from the predicted V119/L135 hydrophobic binding pocket. Instead, V119 of Fc γ RIIa interacts with Fc residue G237 closer to the binding pocket for R134 of the HR allele, which as correctly predicted (46) does not have sufficient space (i.e., this residue adopts a bent or cramped conformation) in its binding site when compared with H134 of Fc γ RIIa-LR.

Structural mapping of epitopes of Fc γ RII blocking Abs

To define how three distinct anti-Fc γ RII Abs block IgG binding, we mapped their epitopes using a mutagenesis approach and correlated the findings to the crystal structures of Fc γ RIIa. The mAbs include IV.3 that specifically binds to Fc γ RIIa, X63-21/7.2 specifically recognizing Fc γ RIIb, and 8.7, which recognizes both Fc γ RIIa and Fc γ RIIb. All Abs inhibit immune complex IgG binding to the respective receptors.

mAbs 8.7 and IV.3 block IgG binding and have been shown to bind to the second extracellular domain through the use of chimeric

receptors containing sequence elements derived from Fc γ RIIa and Fc ϵ RI (35). This was evident from their binding to chimera $\epsilon\gamma$ containing only the second domain of Fc γ RIIa (Supplemental Fig. 1B). Further analyses of these Fc γ RIIa/Fc ϵ RI receptor chimeras indicate that 8.7 recognizes a discontinuous epitope (Supplemental Figs. 1, 2). Ab 8.7 bound to only those chimeric receptors containing residues 112–119 together with 157–162 (chimeras 3 and 4) and failed to bind to any receptor lacking either one of these segments (chimeras 1 and 2; Supplemental Figs. 1, 2). Most importantly, 8.7 bound to chimera 4 wherein the only segments derived from Fc γ RIIa are residues 112–119 and 157–162. These two segments also encompass two IgG contact regions defined in the complex of Fc γ RIIa-IgG1-Fc (Fig. 3B).

Point mutations were then introduced into the Fc γ RIIa to further define the 8.7 epitope. First, residues L162 and F163 were mutated to introduce the equivalent N-linked glycosylation site of Fc γ RIIIa (47), N162 and V163, into Fc γ RIIa. Normal Fc γ RIIa bound aggregated IgG, as well as IV.3 and 8.7. However, the L162N mutant showed only weak binding of 8.7, whereas the binding of IV.3 was unaffected (Supplemental Fig. 3), indicating a role for this residue in the 8.7 epitope. Not surprisingly, the binding of aggregated IgG was diminished. The second component of the 8.7 epitope involves residues 112–119 and includes W113, which is pivotally involved in the interaction with IgG (Fig. 3B). The mutation of W90 and W113 completely abolished 8.7 binding, whereas the mutation of W90 alone had no effect, indicating that W113 is essential for the 8.7 epitope (Table III). Furthermore, both W113 and L162 are present in Fc γ RIIb to which 8.7 also binds.

The IV.3 mAb specifically binds Fc γ RIIa, but not Fc γ RIIb, and its epitope was localized to residues 132–137. Replacement of this sequence in Fc γ RIIa by the equivalent six amino acids from Fc ϵ RI (chimera 6) destroys IV.3 binding, whereas all other chimeras containing this Fc γ RIIa sequence bound IV.3 (chimeras 1, 2, 3, 5; Table III, Supplemental Figs. 1, 2). This epitope contains the HR/LR polymorphism of Fc γ RIIa, but the fact that IV.3 binds to both Fc γ RIIa-HR and Fc γ RIIa-LR indicates either arginine or histidine at position 134 is compatible with IV.3 binding. However, in Fc γ RIIa and Fc γ RIIb, this sequence differs at residue 135 and it is also flanked by two additional sequence differences at residues 130 and 138. Mutation of Q130 or T138 in Fc γ RIIa to K130 or N138 of Fc γ RIIb had no detectable effect on IV.3 binding. However, replacement of L135 in both the Fc γ RIIa-HR and Fc γ RIIa-LR with S135 of Fc γ RIIb completely destroyed the IV.3

Table III. Summary of epitope mapping data using Fc γ RIIa mutants

Receptor Construct	Residue Number ^a						Binding Activity			
	90	113	130	134–138	162	163	8.7	IV.3	X63-21	HAGG
Fc γ RIIa-LR WT	W	W	Q	HLDPT	L	F	++	++	–	++
Fc γ RIIa-HR WT	.	.	.	R	++	++	–	ND
Fc γ RIIb WT	.	.	K	RS . . N	.	.	++	–	++	++
Fc γ RIIa-LR W90A	A	++	++	ND	+
Fc γ RIIa-LR W90A,W113A	A	A	–	++	ND	±
Fc γ RIIa-LR Q130K	.	.	K	++	++	ND	++
Fc γ RIIa-LR L135SS	++	–	–	++
Fc γ RIIa-HR L135S	.	.	.	RS	++	–	++	++
Fc γ RIIa-HR T138N N	.	.	++	++	ND	++
Fc γ RIIa-HR L135S, T138NS . . N	.	.	++	–	++	++
Fc γ RIIa-HR L162N, F163V	N	V	±	++	ND	+

^a“.” indicates residues that are unaltered from the wild-type sequence of Fc γ RIIa-LR, Fc γ RIIa-HR, or Fc γ RIIb. Residue W113 is essential to the 8.7 epitope. Residues L135 and R134-S135 are essential to the IV.3 and X63-21/7.2 epitopes in Fc γ RIIa and Fc γ RIIb, respectively.

++, maximal binding equivalent to WT Fc γ RIIa; +, intermediate binding; ±, weak detectable binding; –, undetectable binding over background, as exemplified in Fig. 4 and Supplemental Fig. 3; ND, no data.




FIGURE 4. FACS analyses of mAb binding to mutants of FcγRII. Mock-transfected CHOP cells (blue line), FcγRIIa-LR WT (green line), or FcγRIIa-LR L135S mutant receptor (black line) were incubated with (A) biotinylated 8.7 F(ab')₂ or (B) biotinylated IV.3 Fab. C, FcγRIIb transfectants (red line) and mutant receptors FcγRIIa-HR L135S (orange line) and FcγRIIa-HR L135S T138N (magenta line) were reacted with the FcγRIIb-selective biotinylated F(ab')₂ of mAb X63-21/7.2. D, Ligand binding activity of the transfected cells was tested with heat-aggregated gamma globulin (HAGG) and allophycocyanin-conjugated anti-human IgG. C and D, Transfections were spiked with 1/40 EGFP-N1 (Clontech) and FL1⁺ cells gated to select only transfected cells for analysis.

epitope and had no effect on 8.7 (Fig. 4, Table III), indicating L135 is an essential part of the IV.3 epitope and determines its selectivity for FcγRIIa.

The epitope detected by FcγRIIb-specific blocking mAb X63-21/7.2 was mapped to an area equivalent to that of the IV.3 epitope in FcγRIIa. In a reciprocal pattern to IV.3 binding, X63-21/7.2 fails to bind FcγRIIa or the FcγRIIa-LR L135S, but does stain FcγRIIa-HR L135S, T138N, and most importantly, FcγRIIa-HR L135S (Fig. 4), indicating both R134 and S135 are essential to the X63-21/7.2 epitope and S135 for its selectivity for FcγRIIb.

When the epitopes defined by the chimeric receptor and mutagenesis experiments were mapped onto the solvent-accessible surfaces of FcγRIIa, it was apparent that the epitopes recognized by 8.7 and IV.3 were likely to be nonoverlapping (Fig. 5). Importantly, both blocking Abs also bound to regions that were clearly used for interactions with IgG1-Fc as defined by the crystal structure of the FcγRIIa-HR-IgG1-Fc. Thus, these mAbs are clearly competitive inhibitors of binding of immune complexes to the FcγRII class of receptors.

Model of the activation complex of FcγRIIa

Immune complexes (Ag-IgG) drive the aggregation of FcγRIIa on the cell surface to initiate intracellular receptor phosphorylation and downstream signaling (25, 27, 48). However, the arrangement of Ag-IgG-FcγRIIa complexes within these aggregates has not been characterized. Furthermore, we previously demonstrated that a proportion of FcγRIIa is constitutively present as a dimer on the cell surface (25). Thus, the dimeric forms of FcγRIIa that we have observed in crystals of glycosylated receptor (Fig. 1) may represent the receptor presentation on the surface of cells. We attempted to overlay the coordinates for the FcγRIIa-Fc complex onto the two crystallographic FcγRIIa dimers. The dimer represented in crystals of FcγRIIa-LR was not capable of binding a single Fc without clashing with the neighboring receptor mono-




FIGURE 5. Structural location of epitopes recognized by FcγRIIa blocking mAb. A, FcγRIIa-HR with IgG1 binding residues mapped to the surface colored according to Fig. 3. B, Epitope regions defined to bind to 8.7. C, Epitope region defined to bind to IV.3. The views of FcγRIIa are shown in the same orientation.

mer. In contrast, the FcγRIIa-HR crystallographic dimer was capable of accommodating either one or two Fc ligands without any major steric clashes (Fig. 6). In this model, the hinge regions of the two trapped Fc ligands are nearly abutting, but are bent away from the proposed location of the FcγRIIa molecules and the membrane. As a result, the Fab arms, tethered by flexible hinge regions, are positioned a considerable distance away from the cell membrane and are readily available for binding Ag. This arrangement suggests that the elements in the hinges and CH2 domains of opposing receptor-bound Fc can interact (e.g., lower hinge residue E233 from one Fc can form a potential salt-bridge with K326 on the second Fc). This is entirely consistent with a comprehensive mutagenesis study where mutations in the lower hinge region including E233 reduced FcR binding to IgG1 (39). It is important to note that rather than the free IgG bound to FcγRIIa as in crystals and the model (Fig. 6), *in vivo* Ag-IgG binds to the receptors. However, the structures reported in this article may represent snapshots of the possible arrangements of receptors within the membrane and in signaling assemblies.

Discussion

The Fc region of IgG provides humoral immunity with a cell-based effector system through the binding of IgG immune complexes to their specific activating receptors: FcγRI (CD64), FcγRIII (CD16), and FcγRIIa (CD32) in humans. These interactions are critical for normal immunity, but in autoimmunity are capable, especially through FcγRIIa, of inducing powerful inflammatory responses with adverse consequences in diseases like rheumatoid arthritis and SLE (1).

In addition, the pivotal role of the Ab Fc in humoral immunity has directed the evolution of bacterial and viral virulence




FIGURE 6. Structural considerations for clustering and activation of Fc γ RIIa by immune complexes. *A*, Molecular model for the activation complex of Fc γ RIIa with two bound IgG ligands. The model was generated by rigid body superposition of the coordinates for the unliganded Fc γ RIIa-HR dimer with those of its complex with the Fc region of human IgG1, and the Fabs were derived from hu3S193 Fab (PDB code 1S3K). Two orthogonal views are shown of IgG bound to dimeric receptor. Sufficient space is available to accommodate the Fab arms that emerge as they are tethered to Fc by long, flexible hinge regions. *B*, Schematic view of FcR clustering based on the available crystallographic information. An inactive or resting Fc γ RIIa dimer (state I) dissociates for steric reasons with the addition of Ag-IgG (immune complexes) to produce an Ag-IgG-Fc γ RIIa complex (state II). Rearrangements of Ag-IgG-Fc γ RIIa complexes can form a dimeric activation complex (state III) composed of either two Ag-IgG bound to the FcR dimer (shown in *A*) or alternatively a single Ag-IgG interacting with receptor associated with an unliganded monomer. These component assemblies may cluster to form larger aggregates, similar to microdomains, and act as potent inducers of inflammation.

factors in their development of immune evasion mechanisms. Crystallographic studies of many immune system (host) and pathogen proteins in complex with Ab Fc regions have provided a wealth of information on the mechanisms of effector functions and immune evasion (32, 38, 40, 42, 49–52). These studies suggest that recognition strategies have converged toward the binding of proteins at one of two hot spots on the Fc portion of Ab. First, the conventional FcRs interact with the CH2 domains and adjacent lower hinge region, including Fc γ RIIa (this study), Fc γ RIII (32, 38), and Fc ϵ RI (40) as a complex with Ce2-Ce4 of IgE-Fc (53); this is also likely to be the case for Fc γ RI (46, 54). The second important Fc hot spot is located at the junction of the CH2 and CH3 domains, and is recognized by many molecules such as the MHC-related neonatal FcR (55). Numerous proteins produced by significant human pathogens also bind the CH2-CH3 domain junction such as protein A from *Staphylococcus aureus* (42); the equivalent site in IgA is the target for host and pathogen proteins (50, 52, 56). The CH2-CH3 domain junction has also been identified as a target for bacterial Ig-binding proteins across species (57). Furthermore, the CH2-CH3 domain junction is the primary site of rheumatoid factor recognition directed against IgG (49). In IgM, the CH3-CH4 domain junction is analogous to CH2-CH3 of IgG and appears to be a recognition site for both the Fca/ μ receptor (IgA and IgM receptor) (58) and some strains of *P. falciparum* erythrocyte membrane protein 1 (59).

Clinically important polymorphisms occur in Fc γ RIIa (R134/H134; HR/LR) and Fc γ RIII (V158/F158), which influence IgG binding and are known to be associated with susceptibility to Gram-negative infection (15), autoimmune diseases (10, 13, 45), and malaria (23). Furthermore, a structural understanding of the impact of the receptor polymorphisms on recognition of different IgG glycoforms or subclasses is of particular importance for the development of effective Ab-based therapies. For example, the homozygous H134/H134 (Fc γ RIIa) and V158/V158 (Fc γ RIII) genotypes were independently associated with higher response rates and freedom from disease progression in patients being treated for follicular lymphoma with a CD20-specific mAb, rituximab (16).

In this study, we observed the R134 of the Fc γ RIIa-HR binds in a shallow depression in the Fc close to the Fuc from the adjacent N297-linked carbohydrate on Fc. Removal of Fuc from human IgG1 has been shown to slightly improve the binding of the R134 (Fc γ RIIa-HR and Fc γ RIIb), but not the H134 (Fc γ RIIa-LR) form. The effect was even more pronounced in Fc γ RIII where binding to Fuc-deficient IgG1 was improved by up to 50-fold (F158 > V158) compared with native IgG1 (44). In the Fc γ RIII complex with Fc, the fucose does not participate in the interaction (32, 38). Furthermore, the carbohydrate on N162 of Fc γ RIII is needed for the high-affinity binding to Fuc-deficient IgG1, and this oligosaccharide (not present on Fc γ RIIa) is predicted to interact with the carbohydrate moieties of the Fc (60). Thus, different mechanisms may mediate the variation in binding of IgG glycoforms observed for Fc γ RII and Fc γ RIII. However, further studies are required to establish the link between IgG fucosylation and molecular interactions involving Fc γ RIIa polymorphisms.

The IgG binding regions on Fc γ RIIa are localized to two nonoverlapping sites on the receptor (Fig. 5A). The same regions on Fc γ RII that bind IgG were determined to be the predominant epitopes recognized by mAbs 8.7, IV.3, and X63-21/7.2, which establishes that these blocking Abs are direct inhibitors of immune complexes binding by the Fc γ RII class of receptors. The two Fc γ RIIa binding Abs, 8.7 (Fig. 5B) and IV.3 (Fig. 5C), bind to the nonoverlapping regions that correspond with the two distinct regions used by the receptor to bind IgG. The selectivity of mAb IV.3 for Fc γ RIIa (L135) and X63-21/7.2 for Fc γ RIIb (S135) is dependent on the difference at residue 135 between these two receptors. Although our studies have established the structural locations for the epitopes of these Fc γ RII blocking Abs, there remain significant challenges in developing intact IgG as potential therapeutics for immune complex-mediated disease. Importantly, we have determined the crystal structure of Fc γ RIIa in complex with IgG, which provides details of the target site for binding immune complexes by this human inflammatory receptor.

The Fc in the complex with Fc γ RIIa-HR was prepared from the hu3S193 IgG1 that has potent FcR-dependent, Ab-dependent cellular cytotoxicity against tumor cells bearing its target Le^y Ag. We have previously determined the crystal structure of the hu3S193 Fab in complex with the Le^y tetrasaccharide and shown that this compact epitope binds end-on into the Ag-binding cavity (29). Because the Le^y epitopes are displayed at very high densities on the surface of tumor cells, avid crosslinking of Le^y by hu3S193 IgG (61) would enhance clustering of FcRs (including Fc γ RIIa) on effector cells, thereby eliciting Ab-dependent cellular cytotoxicity. Similar mechanisms are certainly involved in FcR-mediated killing observed with other therapeutic Abs. In addition to the forces of Ag-driven clustering, it is possible that interactions between the FcRs themselves are involved in organization and activation of these cell surface receptors.

Fc γ RIIa, like all activating FcRs and indeed other multisubunit immunoreceptors including the Ag receptors, relies on ITAM-

based signaling pathways to initiate signaling cascades. The ITAM signaling pathway typically relies on the noncovalent association of ligand-binding chains with covalently dimerized polypeptides such as the common FcR- γ -chain, the CD3 complex of TCR complex, or Ig α / β associated with the B cell Ag receptor complex. However, in this respect, Fc γ RIIa is unique, because it does not associate with, nor require, separate ITAM-containing dimers for activation of cells. Receptor dimerization as anticipated in this study (Figs. 1, 6) would juxtapose the ITAM-containing cytoplasmic tails. We postulate that there is a dynamic interplay between quaternary states of Fc γ RIIa, which may be involved in immune complex activation leading to inflammation (Fig. 6). The structural information on Fc γ RIIa suggests that a minimum of three quaternary states may account for the available biochemical data on the interplay between Ag-IgG complexes and the cell surface-associated receptor. In such a model, Fc γ RIIa exists as a resting or inactive dimer (state I, Fig. 6) that requires dissociation before, or upon, binding an Ag-IgG complex (state II, Fig. 6). Lateral packing of receptors (as in crystals of the Fc γ RIIa-HR) with bound Ag-IgG complexes will bring two ITAMs together to form signaling competent complex (state III, Fig. 6), a building block for a larger cluster, which efficiently mediates cell activation. Depending on the presentation of Ag-Ab complexes, a variation on state III of one, not two, Ag-IgG complexes and a receptor dimer may also occur. In any event, for both cases, the liganded dimers contain cytoplasmic tails arranged analogously to the dimeric FcR- γ subunit of other immunoreceptors, and so could mediate intracellular signaling and inflammation.

In summary, we provide in this article a structural rationale for the interplay between glycosylated Fc γ RIIa monomers, their dimers, and binding of immune complexes on the cell surface that lead to ordered receptor clustering, signaling cascades, and inflammation. Collectively, the structural data suggest how the HR/LR polymorphism of Fc γ RIIa influences binding to different IgG subclasses and possibly glycoforms such as fucose-deficient IgG. Finally, the same sites that are involved in binding IgG were found to be targeted by blocking Abs specific for Fc γ RII.

Disclosures

The authors have no financial conflicts of interest.

References

- Hogarth, P. M. 2002. Fc receptors are major mediators of antibody based inflammation in autoimmunity. *Curr. Opin. Immunol.* 14: 798–802.
- Takai, T. 2005. Fc receptors and their role in immune regulation and autoimmunity. *J. Clin. Immunol.* 25: 1–18.
- Masuda, A., M. Yoshida, H. Shiomi, Y. Morita, H. Kutsumi, H. Inokuchi, S. Mizuno, A. Nakamura, T. Takai, R. S. Blumberg, and T. Azuma. 2009. Role of Fc receptors as a therapeutic target. *Inflamm. Allergy Drug Targets* 8: 80–86.
- Kinet, J. P. 1992. The gamma-zeta dimers of Fc receptors as connectors to signal transduction. *Curr. Opin. Immunol.* 4: 43–48.
- Wines, B. D., H. M. Trist, P. A. Ramsland, and P. M. Hogarth. 2006. A common site of the Fc receptor gamma subunit interacts with the unrelated immunoreceptors Fc α RI and Fc ϵ RI. *J. Biol. Chem.* 281: 17108–17113.
- Tan Sardjono, C., P. L. Mottram, and P. M. Hogarth. 2003. The role of Fc γ RIIa as an inflammatory mediator in rheumatoid arthritis and systemic lupus erythematosus. *Immunol. Cell Biol.* 81: 374–381.
- McKenzie, S. E., S. M. Taylor, P. Malladi, H. Yuhan, D. L. Cassel, P. Chien, E. Schwartz, A. D. Schreiber, S. Surrey, and M. P. Reilly. 1999. The role of the human Fc receptor Fc γ RIIa in the immune clearance of platelets: a transgenic mouse model. *J. Immunol.* 162: 4311–4318.
- Pietersz, G. A., P. L. Mottram, N. C. van de Velde, C. T. Sardjono, S. Esparon, P. A. Ramsland, G. Moloney, J. B. Baell, T. D. McCarthy, B. R. Matthews, et al. 2009. Inhibition of destructive autoimmune arthritis in Fc γ RIIa transgenic mice by small chemical entities. *Immunol. Cell Biol.* 87: 3–12.
- Means, T. K., E. Latz, F. Hayashi, M. R. Murali, D. T. Golenbock, and A. D. Luster. 2005. Human lupus autoantibody-DNA complexes activate DCs through cooperation of CD32 and TLR9. *J. Clin. Invest.* 115: 407–417.
- Salmon, J. E., S. Millard, L. A. Schachter, F. C. Arnett, E. M. Ginzler, M. F. Gourley, R. Ramsey-Goldman, M. G. Peterson, and R. P. Kimberly. 1996.

- Fc gamma RIIA alleles are heritable risk factors for lupus nephritis in African Americans. *J. Clin. Invest.* 97: 1348–1354.
- Zuniga, R., G. S. Markowitz, T. Arkachaisri, E. A. Imperatore, V. D. D'Agati, and J. E. Salmon. 2003. Identification of IgG subclasses and C-reactive protein in lupus nephritis: the relationship between the composition of immune deposits and Fc γ receptor type IIA alleles. *Arthritis Rheum.* 48: 460–470.
- Loos, B. G., F. G. J. Leppers-Van de Straat, J. G. J. Van de Winkel, and U. Van der Velden. 2003. Fc γ receptor polymorphisms in relation to periodontitis. *J. Clin. Periodontol.* 30: 595–602.
- Dijstelbloem, H. M., M. Bijl, R. Fijnheer, R. H. Scheepers, W. W. Oost, M. D. Jansen, W. J. Sluiter, P. C. Limburg, R. H. Derksen, J. G. van de Winkel, and C. G. Kallenberg. 2000. Fc γ receptor polymorphisms in systemic lupus erythematosus: association with disease and in vivo clearance of immune complexes. *Arthritis Rheum.* 43: 2793–2800.
- Asano, K., T. Matsushita, J. Umeno, N. Hosono, A. Takahashi, T. Kawaguchi, T. Matsumoto, T. Matsui, Y. Kakuta, Y. Kinouchi, et al. 2009. A genome-wide association study identifies three new susceptibility loci for ulcerative colitis in the Japanese population. *Nat. Genet.* 41: 1325–1329.
- Sanders, L. A., R. G. Feldman, M. M. Voorhorst-Ogink, M. de Haas, G. T. Rijkers, P. J. Capel, B. J. Zegers, and J. G. van de Winkel. 1995. Human immunoglobulin G (IgG) Fc receptor IIA (CD32) polymorphism and IgG2-mediated bacterial phagocytosis by neutrophils. *Infect. Immun.* 63: 73–81.
- Weng, W. K., and R. Levy. 2003. Two immunoglobulin G fragment C receptor polymorphisms independently predict response to rituximab in patients with follicular lymphoma. *J. Clin. Oncol.* 21: 3940–3947.
- Powell, M. S., P. A. Barton, D. Emmanouilidis, B. D. Wines, G. M. Neumann, G. A. Peitersz, K. F. Maxwell, T. P. Garrett, and P. M. Hogarth. 1999. Biochemical analysis and crystallisation of Fc gamma RIIa, the low affinity receptor for IgG. *Immunol. Lett.* 68: 17–23.
- Warmerdam, P. A., J. G. van de Winkel, E. J. Gosselin, and P. J. Capel. 1990. Molecular basis for a polymorphism of human Fc gamma receptor II (CD32). *J. Exp. Med.* 172: 19–25.
- Tate, B. J., E. Witort, I. F. McKenzie, and P. M. Hogarth. 1992. Expression of the high responder/non-responder human Fc gamma RII. Analysis by PCR and transfection into FcR-COS cells. *Immunol. Cell Biol.* 70: 79–87.
- Tax, W. J., H. W. Willems, P. P. Reekers, P. J. Capel, and R. A. Koene. 1983. Polymorphism in mitogenic effect of IgG1 monoclonal antibodies against T3 antigen on human T cells. *Nature* 304: 445–447.
- Parren, P. W., P. A. Warmerdam, L. C. Boeije, J. Arts, N. A. Westerdaal, A. Vlugs, P. J. Capel, L. A. Aarden, and J. G. van de Winkel. 1992. On the interaction of IgG subclasses with the low affinity Fc gamma RIIa (CD32) on human monocytes, neutrophils, and platelets. Analysis of a functional polymorphism to human IgG2. *J. Clin. Invest.* 90: 1537–1546.
- Gordon, C. L., P. D. Johnson, M. Permezel, N. E. Holmes, G. Gutteridge, C. F. McDonald, D. P. Eisen, A. J. Stewardson, J. Edington, P. G. Charles, et al. 2010. Association between severe pandemic 2009 influenza A (H1N1) virus infection and immunoglobulin G(2) subclass deficiency. *Clin. Infect. Dis.* 50: 672–678.
- Nasr, A., N. C. Iriemenam, H. A. Gihla, H. A. Balogun, R. F. Anders, M. Troye-Blomberg, G. ElGhazali, and K. Berzins. 2009. Fc γ RIIa (CD32) polymorphism and anti-malarial IgG subclass pattern among Fulani and sympatric ethnic groups living in eastern Sudan. *Malar. J.* 8: 43.
- Maxwell, K. F., M. S. Powell, M. D. Hulett, P. A. Barton, I. F. McKenzie, T. P. Garrett, and P. M. Hogarth. 1999. Crystal structure of the human leukocyte Fc receptor, Fc gammaRIIa. *Nat. Struct. Biol.* 6: 437–442.
- Powell, M. S., N. C. Barnes, T. M. Bradford, I. F. Musgrave, B. D. Wines, J. C. Cambier, and P. M. Hogarth. 2006. Alteration of the Fc gamma RIIa dimer interface affects receptor signaling but not ligand binding. *J. Immunol.* 176: 7489–7494.
- Faeder, J. R., W. S. Hlavacek, I. Reischl, M. L. Blinov, H. Metzger, A. Redondo, C. Wofsy, and B. Goldstein. 2003. Investigation of early events in Fc epsilon RI-mediated signaling using a detailed mathematical model. *J. Immunol.* 170: 3769–3781.
- Metzger, H. 1992. Transmembrane signaling: the joy of aggregation. *J. Immunol.* 149: 1477–1487.
- Metzger, H. 2002. Molecular versatility of antibodies. *Immunol. Rev.* 185: 186–205.
- Ramsland, P. A., W. Farrugia, T. M. Bradford, P. Mark Hogarth, and A. M. Scott. 2004. Structural convergence of antibody binding of carbohydrate determinants in Lewis Y tumor antigens. *J. Mol. Biol.* 340: 809–818.
- Otwinowski, Z., and W. Minor. 1997. Processing of X-ray diffraction data collected in oscillation mode. In *Macromolecular Crystallography. Part A. Methods in Enzymology*, Vol. 276. C. W. Carter, Jr. and R. M. Sweet, eds. Academic Press, New York, p. 307–326.
- Brünger, A. T., P. D. Adams, G. M. Clore, W. L. DeLano, P. Gros, R. W. Grosse-Kunstleve, J. S. Jiang, J. Kuszewski, M. Nilges, N. S. Pannu, et al. 1998. Crystallography & NMR system: a new software suite for macromolecular structure determination. *Acta Crystallogr. D Biol. Crystallogr.* 54: 905–921.
- Sondermann, P., R. Huber, V. Oosthuizen, and U. Jacob. 2000. The 3.2-Å crystal structure of the human IgG1 Fc fragment-Fc gammaRIIIc complex. *Nature* 406: 267–273.
- Ierino, F. L., M. D. Hulett, I. F. McKenzie, and P. M. Hogarth. 1993. Mapping epitopes of human Fc gamma RII (CD32) with monoclonal antibodies and recombinant receptors. *J. Immunol.* 150: 1794–1803.
- Looney, R. J., G. N. Abraham, and C. L. Anderson. 1986. Human monocytes and U937 cells bear two distinct Fc receptors for IgG. *J. Immunol.* 136: 1641–1647.

35. Hulett, M. D., I. F. McKenzie, and P. M. Hogarth. 1993. Chimeric Fc receptors identify immunoglobulin-binding regions in human Fc gamma RII and Fc epsilon RI. *Eur. J. Immunol.* 23: 640–645.
36. Sondermann, P., and U. Jacob. 1999. Human Fc gamma receptor IIb expressed in *Escherichia coli* reveals IgG binding capability. *Biol. Chem.* 380: 717–721.
37. Hulett, M. D., E. Witort, R. I. Brinkworth, I. F. McKenzie, and P. M. Hogarth. 1995. Multiple regions of human Fc gamma RII (CD32) contribute to the binding of IgG. *J. Biol. Chem.* 270: 21188–21194.
38. Radaev, S., S. Motyka, W. H. Fridman, C. Sautes-Fridman, and P. D. Sun. 2001. The structure of a human type III Fc gamma receptor in complex with Fc. *J. Biol. Chem.* 276: 16469–16477.
39. Shields, R. L., A. K. Namenuk, K. Hong, Y. G. Meng, J. Rae, J. Briggs, D. Xie, J. Lai, A. Stadlen, B. Li, et al. 2001. High resolution mapping of the binding site on human IgG1 for Fc gamma RI, Fc gamma RII, Fc gamma RIII, and Fc gamma Rn and design of IgG1 variants with improved binding to the Fc gamma R. *J. Biol. Chem.* 276: 6591–6604.
40. Garman, S. C., B. A. Wurzburg, S. S. Tarchevskaya, J. P. Kinet, and T. S. Jardetzky. 2000. Structure of the Fc fragment of human IgE bound to its high-affinity receptor Fc epsilon RI alpha. *Nature* 406: 259–266.
41. Cendron, A. C., B. D. Wines, R. T. Brownlee, P. A. Ramsland, G. A. Pietersz, and P. M. Hogarth. 2008. An Fc gamma RIIa-binding peptide that mimics the interaction between Fc gamma RIIa and IgG. *Mol. Immunol.* 45: 307–319.
42. Deisenhofer, J. 1981. Crystallographic refinement and atomic models of a human Fc fragment and its complex with fragment B of protein A from *Staphylococcus aureus* at 2.9- and 2.8-A resolution. *Biochemistry* 20: 2361–2370.
43. Krapp, S., Y. Mimura, R. Jefferis, R. Huber, and P. Sondermann. 2003. Structural analysis of human IgG-Fc glycoforms reveals a correlation between glycosylation and structural integrity. *J. Mol. Biol.* 325: 979–989.
44. Shields, R. L., J. Lai, R. Keck, L. Y. O'Connell, K. Hong, Y. G. Meng, S. H. A. Weikert, and L. G. Presta. 2002. Lack of fucose on human IgG1 N-linked oligosaccharide improves binding to human Fc gamma RIII and antibody-dependent cellular toxicity. *J. Biol. Chem.* 277: 26733–26740.
45. Fossati, G., R. C. Bucknall, and S. W. Edwards. 2001. Fc gamma receptors in autoimmune diseases. *Eur. J. Clin. Invest.* 31: 821–831.
46. Sondermann, P., J. Kaiser, and U. Jacob. 2001. Molecular basis for immune complex recognition: a comparison of Fc-receptor structures. *J. Mol. Biol.* 309: 737–749.
47. Drescher, B., T. Witte, and R. E. Schmidt. 2003. Glycosylation of Fc gamma RIII in N163 as mechanism of regulating receptor affinity. *Immunology* 110: 335–340.
48. Bournazos, S., S. P. Hart, L. H. Chamberlain, M. J. Glennie, and I. Dransfield. 2009. Association of Fc gamma RIIa (CD32a) with lipid rafts regulates ligand binding activity. *J. Immunol.* 182: 8026–8036.
49. Corper, A. L., M. K. Sohi, V. R. Bonagura, M. Steinitz, R. Jefferis, A. Feinstein, D. Beale, M. J. Taussig, and B. J. Sutton. 1997. Structure of human IgM rheumatoid factor Fab bound to its autoantigen IgG Fc reveals a novel topology of antibody-antigen interaction. *Nat. Struct. Biol.* 4: 374–381.
50. Herr, A. B., E. R. Ballister, and P. J. Bjorkman. 2003. Insights into IgA-mediated immune responses from the crystal structures of human Fc alpha RI and its complex with IgA1-Fc. *Nature* 423: 614–620.
51. Sprague, E. R., C. Wang, D. Baker, and P. J. Bjorkman. 2006. Crystal structure of the HSV-1 Fc receptor bound to Fc reveals a mechanism for antibody bipolar bridging. *PLoS Biol.* 4: e148.
52. Ramsland, P. A., N. Willoughby, H. M. Trist, W. Farrugia, P. M. Hogarth, J. D. Fraser, and B. D. Wines. 2007. Structural basis for evasion of IgA immunity by *Staphylococcus aureus* revealed in the complex of SSL7 with Fc of human IgA1. *Proc. Natl. Acad. Sci. USA* 104: 15051–15056.
53. Holdom, M. D., A. M. Davies, J. E. Nettleship, S. C. Bagby, B. Dhaliwal, E. Girardi, J. Hunt, H. J. Gould, A. J. Bevil, J. M. McDonnell, et al. 2011. Conformational changes in IgE contribute to its uniquely slow dissociation rate from receptor Fc epsilon RI. *Nat. Struct. Mol. Biol.* 18: 571–576.
54. Lund, J., G. Winter, P. T. Jones, J. D. Pound, T. Tanaka, M. R. Walker, P. J. Artymiuk, Y. Arata, D. R. Burton, R. Jefferis, and J. M. Woof. 1991. Human Fc gamma RI and Fc gamma RII interact with distinct but overlapping sites on human IgG. *J. Immunol.* 147: 2657–2662.
55. Burmeister, W. P., L. N. Gastinel, N. E. Simister, M. L. Blum, and P. J. Bjorkman. 1994. Crystal structure at 2.2 A resolution of the MHC-related neonatal Fc receptor. *Nature* 372: 336–343.
56. Pleass, R. J., T. Areschoug, G. Lindahl, and J. M. Woof. 2001. Streptococcal IgA-binding proteins bind in the Alpha 2-Calpha 3 interdomain region and inhibit binding of IgA to human CD89. *J. Biol. Chem.* 276: 8197–8204.
57. Lewis, M. J., M. Meehan, P. Owen, and J. M. Woof. 2008. A common theme in interaction of bacterial immunoglobulin-binding proteins with immunoglobulins illustrated in the equine system. *J. Biol. Chem.* 283: 17615–17623.
58. Ghumra, A., J. Shi, R. S. McIntosh, I. B. Rasmussen, R. Braathen, F. E. Johansen, I. Sandlie, P. K. Mongini, T. Areschoug, G. Lindahl, et al. 2009. Structural requirements for the interaction of human IgM and IgA with the human Fc alpha/mu receptor. *Eur. J. Immunol.* 39: 1147–1156.
59. Ghumra, A., J. P. Semblat, R. S. McIntosh, A. Raza, I. B. Rasmussen, R. Braathen, F. E. Johansen, I. Sandlie, P. K. Mongini, J. A. Rowe, and R. J. Pleass. 2008. Identification of residues in the Cmu4 domain of polymeric IgM essential for interaction with *Plasmodium falciparum* erythrocyte membrane protein 1 (PfEMP1). *J. Immunol.* 181: 1988–2000.
60. Ferrara, C., F. Stuart, P. Sondermann, P. Brunker, and P. Umaña. 2006. The carbohydrate at Fc gamma RIIIa Asn-162. An element required for high affinity binding to non-fucosylated IgG glycoforms. *J. Biol. Chem.* 281: 5032–5036.
61. Farrugia, W., A. M. Scott, and P. A. Ramsland. 2009. A possible role for metallic ions in the carbohydrate cluster recognition displayed by a Lewis Y specific antibody. *PLoS ONE* 4: e7777.
62. Kabat, E. A., T. T. Wu, H. M. Perry, K. S. Gottesman, and C. Foeller. 1991. *Sequences of Proteins of Immunological Interest*. Public Health Service, U. S. Department of Health and Human Services. National Institutes of Health, Bethesda, MD.