



Mabtech IRIS™
SPOT ANALYSIS REINVENTED

Read how >

MABTECH

Capture | Detect | Discover



In This Issue

J Immunol 2013; 190:2475-2476; ;

doi: 10.4049/jimmunol.1390009

<http://www.jimmunol.org/content/190/6/2475>

This information is current as
of October 23, 2018.

Supplementary Material <http://www.jimmunol.org/content/suppl/2013/03/01/190.6.2475.DC1>

Why *The JI*? Submit online.

- **Rapid Reviews! 30 days*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

**average*

Subscription Information about subscribing to *The Journal of Immunology* is online at:
<http://jimmunol.org/subscription>

Permissions Submit copyright permission requests at:
<http://www.aai.org/About/Publications/JI/copyright.html>


Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://jimmunol.org/alerts>

The Journal of Immunology is published twice each month by
The American Association of Immunologists, Inc.,
1451 Rockville Pike, Suite 650, Rockville, MD 20852
Copyright © 2013 by The American Association of
Immunologists, Inc. All rights reserved.
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



Autoimmunity through Negative Regulation

A single point mutation, E613R, in the cytoplasmic membrane-proximal wedge domain of CD45 causes a lupus-like autoimmune disease in susceptible mice. With this point mutation, the CD45 cytoplasmic domain no longer dimerizes and inhibits the phosphatase activity that regulates BCR signaling. To determine the mechanism by which a dysfunctional CD45 contributes to disease, Zikherman et al. (p. 2527) examined BCR signaling in mice with different CD45 expression levels. Interestingly, both B cells that overexpressed CD45 and E613R B cells displayed hyper-responsive B cell signaling. However, B cells with supraphysiologic levels of CD45 became anergic and only mice with the E613R mutation became frankly autoimmune. This difference was due to the E613R mutation changing the substrate specificity of CD45 for Lyn. Thus, autoimmunity in E613R mice was caused by blocking the action of the Lyn-specific negative regulatory pathway. By demonstrating that autoimmunity in these mice results from relaxing the negative regulation of BCR signaling rather than an overdrive of positive regulation, the authors have provided an important mechanistic insight into autoimmune disease.



LAT Links *Listeria* and T Lymphocytes

Following an initial pathogenic infection, naive CD8⁺ T cells undergo a three-phase response: expansion, contraction, and memory formation. The linker for activation of T cells (LAT), which links TCR activation to downstream signaling, is critical for thymocyte development and T cell activation. To examine the role of LAT in this three-phase response following *Listeria* infection, Ou-Yang et al. (p. 2938) used an OVA-specific LAT conditional knockout mouse (LATKO) to delete LAT in Ag-specific CD8⁺ T cells at different stages of the immune response. T cell expansion was shown to be dependent upon LAT, since CD8⁺ LATKO T cells adoptively transferred into wild-type (WT) mice prior to infection with *Listeria* failed to proliferate compared with WT T cells. CD8⁺ LATKO T cells did not demonstrate an increase in programmed cell death, suggesting that the contraction phase is LAT independent. Deletion of LAT during the effector-to-memory transition led to an increase in the formation of long-lived memory precursor cells (MPECs) and central memory T cells (T_{CM}) compared with WT CD8⁺ T cells. However, Ag stimulation through the TCR demonstrated a profound defect in effector cytokine production and calcium mobilization by LATKO memory CD8⁺ T cells, and these memory cells failed to proliferate in response to a recall


challenge with *Listeria*. The authors conclude that although LAT is dispensable for the contraction phase and memory maintenance of CD8⁺ T cell responses to *Listeria*, it is essential for the recall memory response against these pathogens.

Persistence Pays Off

Memory cell persistence is important to long-lasting host protection from infection. Compared to their CD8⁺ T cell counterparts, memory CD4⁺ T cell populations persist for a shorter time. To better understand the development of CD4⁺ T cell memory, Nelson et al. (p. 2828) examined the stability of pathogen-specific CD4⁺ T cells during a sustained *Salmonella* infection. Using Nramp1-resistant (Nramp1^R) mice, which maintain a persistent intracellular infection after oral administration of *Salmonella*, the authors found that the number of CD4⁺ T cells specific for *Salmonella* peptide:MHC class II (MHCII) ligands remained stable for a year. The stability of this CD4⁺ T cell population was maintained by proliferation in secondary lymphoid organs containing bacteria and continuous presentation of the peptide:MHCII ligand. Phenotypically, these cells were Th1, upregulated PD-1, and became exhausted when exposed to systemic expression of *Salmonella* peptide:MHCII ligand. Taken together, these data indicate that pathogen-specific CD4⁺ T cells can be maintained by Ag-driven proliferation of a small population of cells in discrete lymphoid sites.

Baby Steps to Anergy


Complete activation of naive CD4⁺ T cell responses to pathogens requires engagement of the αβ-TCR/CD3 complex and CD28. This leads to calcium release and phosphorylation of Erk in a MAPK cascade critical for activation of AP-1. Human neonates have a reduced and delayed CD4⁺ T cell immune response to pathogens; however, the exact mechanism for this distinction is unclear. Palin et al. (p. 2682) compared calcium signaling in naive CD4⁺ T cells from cord blood (CB) and adult peripheral blood (APB) and demonstrated an increase in signaling in CB T cells following CD3 cross-linking, reflecting retention of a thymocyte-like phenotype. Additionally, the percent of phosphorylated Erk⁺ cells was higher in CB compared with APB CD4⁺ T cells. Quantitative PCR showed elevated levels of microRNA (miR)-181a, which regulates calcium flux and Erk phosphorylation, in naive CD4⁺ T cells from CB relative to APB, and overexpression of miR-181a in APB T cells significantly increased mean calcium flux. Transfection of an AP-1-dependent luciferase reporter gene demonstrated that CB T cells have significantly lower levels of AP-1 transcription and higher levels of Cbl-b induction following CD3 and CD28 stimulation compared with APB T cells, suggesting a tendency for CB



T cells to become anergic following activation. Thus, calcium signaling and AP-1 transcription are decoupled during the activation of CB CD4⁺ T cells, providing a possible mechanism for the observed impairment in neonatal T cell immunity.

InterFeriNg with Encephalitis

Herpes simplex encephalitis (HSE), marked by enlargement of the lateral ventricles during HSV-1 infection, is a rapidly progressive disease with mortality rates of 30% in treated patients. Data suggest that IFN- γ is protective against the incidence of encephalitis; however, the events leading to the neuropathology and morbidity of HSE remain elusive. Conrady et al. (p. 2807) showed that type I IFN receptor α -chain-deficient (CD118^{-/-}) mice had a significantly increased HSV-1 viral burden in the cerebral cortex and lateral ventricular enlargement relative to wild-type (WT) mice. These CD118^{-/-} mice had 100% mortality, whereas WT mice showed a delayed viral accumulation and 20% mortality. RT-PCR on CNS sections showed that HSV-1 entered the CNS through a neural route and selectively progressed to the ependymal cells. The authors' in vivo encephalitic model limited expression of HSV-1 Ag to the ependymal cells in the CNS of CD118^{-/-} mice and demonstrated a loss of ciliated ependymal cells leading to reduced clearance of cerebrospinal fluid and a resulting expansion of the lateral ventricles. A therapeutic peptide, anti-secretory factor 16 (AF-16), used to reduce intracranial pressure without decreasing viral titer, significantly lowered mortality in HSV-infected WT mice but not CD118^{-/-} mice. Organotypic brain slice cultures from CD118^{-/-} mice infected with HSV-1 showed higher viral titers than WT brain slices, supporting a role for IFN-responsive cells. Depletion of microglia in these cultures further increased viral load, supporting a role for microglia in innate resistance to HSV-1. In conclusion, a functional type I IFN pathway in resident microglial cells is required for resistance to HSV-1-associated lateral ventricle enlargement.



Tuberculosis as Transcript Regulator

The pathogenic *Mycobacterium tuberculosis* enhances its survival by altering the gene expression and IFN- γ response of host macrophages. Salamon et al. (p. 2747) examined how this pathogen affected the mRNA transcripts of STAT1 and IRF-1, IFN- γ responsive transcription factors, in *M. tuberculosis*-infected cells. The authors examined steps involved in the postinitiation processes of mRNA biogenesis to determine if these were responsible for changes in STAT1 and IRF-1 expression levels. These mRNA postinitiation steps can include 3' end cleavage, 5' end capping, elongation, polyadenylation, and splicing of message. Transcriptome analysis of infected cells treated with and without IFN- γ revealed that *M. tuberculosis* infection negatively regulated postinitiation mRNA biogenesis. Patient samples were then examined to determine the clinical relevance of this effect. The authors found that postinitiation mRNA biogenesis steps are suppressed in both latent and active tuberculosis disease, although to a differing extent. Thus, *M. tuberculosis* suppresses macrophage responses through its

action on the postinitiation mRNA biogenesis events of key regulators of IFN- γ signaling.

Taking the STING out of *Listeria*

MPY5 (also known as STING) is thought to be a critical participant in the host response to *Listeria monocytogenes*. Although MPY5 has been shown to be an essential component of host defense against DNA virus infections, its in vivo role in defense against nonviral pathogens is unknown. Jin et al. (p. 2835) observed an increase in bacterial burden in the liver but not the spleen of *Listeria*-infected MPY5-deficient (MPY5^{-/-}) mice compared with wild-type (WT) mice. No differences were observed in serum IFN- γ , IL-1 β , and TNF levels or in the number of B, T, or NK cells in the livers of MPY5^{-/-} mice relative to WT mice. MPY5 is predominantly expressed on CD45⁺ liver cells, of which ~50% are Ly6C^{hi} monocytes newly immigrated from the bone marrow during *Listeria* infection. Compared with WT mice, MPY5^{-/-} mice displayed a significant reduction in the number of Ly6C^{hi} monocytes in the liver and the bloodstream during *Listeria* infection. Diminished production of the CCR2 ligands, MCP-1 and MCP-3, which control emigration of Ly6C^{hi} monocytes, led to their impaired emigration from the bone marrow to the liver in MPY5^{-/-} mice. Adoptive transfer of WT Ly6C^{hi} monocytes into *Listeria*-infected MPY5^{-/-} mice increased the number of Ly6C^{hi} monocytes recruited to the liver and significantly decreased the bacterial burden. Taken together, these data demonstrate a role for MPY5 in the recruitment of Ly6C^{hi} monocytes to the liver, which is essential for bacterial clearance during *Listeria* infection.

More Ways To Kill Brain Cells

Multiple sclerosis (MS) is an inflammatory disorder characterized by the infiltration of immune cells into the CNS and the destruction of oligodendrocytes and myelin membranes. Although myelin-specific CD4⁺ T cells expressing CD56 are cytotoxic toward oligodendrocytes, a previous study showed that an anti-CD56 blocking Ab did not inhibit their cytotoxicity, suggesting the involvement of additional cytotoxic effector molecules. To identify these molecules, Zaguia et al. (p. 2510) analyzed myelin basic protein (MBP)-specific CD4⁺ T cell lines and PHA-activated CD4⁺ T cells for NK-associated markers. The majority of CD56⁺CD4⁺ T cells (85%) expressed NKG2C and had elevated levels of cytotoxic molecules, including FasL, granzyme B, and perforin. Treatment of oligodendrocytes with proinflammatory cytokines increased the expression of HLA-E, the cognate ligand of NKG2C, and blocking this interaction inhibited CD4⁺ T cell-mediated killing of oligodendrocytes. To evaluate the physiological relevance, the authors examined the expression of NKG2C on PBMCs from MS patients and healthy donors by flow cytometry. Increased levels of NKG2C were seen on CD4⁺ T cells from MS patients relative to healthy controls, and patient T cells also expressed higher levels of FasL, granzyme B, and perforin. Additionally, immune cells expressing NKG2C and oligodendrocytes expressing HLA-E were observed in situ in postmortem CNS tissue. Taken together, these data demonstrate a novel mechanism by which CD4⁺ T cells contribute to tissue injury in MS.

