

Inflammasome Reporter Cells

All you have to do is ASC

InvivoGen



IFN- β Expression Is Directly Activated in Human Neutrophils Transfected with Plasmid DNA and Is Further Increased via TLR-4-Mediated Signaling

This information is current as of July 16, 2018.

Nicola Tamassia, Flavia Bazzoni, Vincent Le Moigne, Federica Calzetti, Caterina Masala, Giulia Grisendi, Uta Bussmeyer, Sara Scutera, Marzia De Gironcoli, Claudio Costantini, Tiziana Musso and Marco A. Cassatella

J Immunol published online 22 June 2012
<http://www.jimmunol.org/content/early/2012/06/22/jimmunol.1102985>

Supplementary Material <http://www.jimmunol.org/content/suppl/2012/06/22/jimmunol.1102985.DC1>

Why *The JI*? [Submit online.](#)

- **Rapid Reviews! 30 days*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

**average*

Subscription Information about subscribing to *The Journal of Immunology* is online at: <http://jimmunol.org/subscription>

Permissions Submit copyright permission requests at: <http://www.aai.org/About/Publications/JI/copyright.html>

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at: <http://jimmunol.org/alerts>

The Journal of Immunology is published twice each month by The American Association of Immunologists, Inc., 1451 Rockville Pike, Suite 650, Rockville, MD 20852
Copyright © 2012 by The American Association of Immunologists, Inc. All rights reserved.
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



IFN- β Expression Is Directly Activated in Human Neutrophils Transfected with Plasmid DNA and Is Further Increased via TLR-4–Mediated Signaling

Nicola Tamassia,* Flavia Bazzoni,* Vincent Le Moigne,* Federica Calzetti,* Caterina Masala,* Giulia Grisendi,* Uta Bussmeyer,* Sara Scutera,[†] Marzia De Gironcoli,[‡] Claudio Costantini,* Tiziana Musso,[†] and Marco A. Cassatella*

Upon LPS binding, TLR4 activates a MyD88-dependent pathway leading to the transcriptional activation of proinflammatory genes, as well as a MyD88-independent/TRIF-dependent pathway, responsible for the transcriptional induction of IFN- β . Previous findings delineated that human neutrophils are unable to induce the transcription of IFN- β in response to TLR4 stimulation. Because neutrophils do not express protein kinase C ϵ , a molecule recently reported as essential for initiating the MyD88-independent/TRIF-dependent pathway, we optimized an electroporation method to transfect PKC ϵ into neutrophils with very high efficiency. By doing so, a significant IFN- β mRNA expression was induced, in the absence of LPS stimulation, not only in PKC ϵ -overexpressing neutrophils but also in cells transfected with a series of empty DNA plasmids; however, LPS further upregulated the IFN- β transcript levels in plasmid-transfected neutrophils, regardless of PKC ϵ overexpression. Phosphoimmunoblotting studies, as well as chromatin immunoprecipitation assays targeting the IFN- β promoter, revealed that IFN- β mRNA induction occurred through the cooperative action of IRF3, activated by transfected DNA, and NF- κ B, activated by LPS. Additional immunoblotting and coimmunoprecipitation studies revealed that neutrophils constitutively express various cytosolic DNA sensors, including IFN-inducible protein 16, leucine-rich repeat (in Flightless I) interacting protein-1, and DDX41, as well as that IFN-inducible protein 16 is the intracellular receptor recognizing transfected DNA. Consistently, infection of neutrophils with intracellular pathogens, such as *Bartonella henselae*, *Listeria monocytogenes*, *Legionella pneumophila*, or adenovirus type 5, promoted a marked induction of IFN- β mRNA expression. Taken together, these data raise questions about the role of PKC ϵ in driving the MyD88-independent/TRIF-dependent response and indicate that human neutrophils are able to recognize and respond to microbial cytosolic DNA. *The Journal of Immunology*, 2012, 189: 000–000.

The innate immune system, to which polymorphonuclear neutrophils belong, is highly specialized in its capacity to recognize foreign pathogens as a result of the expression of evolutionarily conserved families of receptor proteins: the pattern recognition receptors (1). The latter include, among others, the TLRs (2), which are constitutively expressed and functional in human neutrophils, with the exception of TLR3 (3–5). For in-

stance, TLR4, which is the specific receptor for endotoxin (6), potently triggers TNF- α , CXCL8, CCL3, CCL4, CCL19, CCL20, IL-1ra, and IL-12p40 mRNA expression and production in neutrophils stimulated with LPS (7). However, unlike human monocytes, endotoxin-activated neutrophils do not express type I IFN (IFN- β and/or IFN- α) or IFN-dependent genes, such as CXCL10 or CXCL9 (8). This was recently attributed to the inability of neutrophils to mobilize the so-called “MyD88-independent/TRIF-dependent” pathway (8). In fact, it has emerged from studies mainly performed in macrophages or dendritic cells of gene-targeted mice that LPS triggers two classes of genes via TLR4 (1). One class is defined as MyD88 dependent, because it is not induced in MyD88^{-/-} mice and mostly includes proinflammatory mediators, such as TNF- α , IL-1, IL-12p40, and CXCL8 (1). In this pathway, MyD88 and TIR domain-containing adapter protein/MyD88 adapter-like protein mediate a rapid and early activation of the transcription factor NF- κ B, which is essential for transcriptional induction of the above-mentioned proinflammatory genes (1). The other class of genes is defined as MyD88-independent, because it relies on a more delayed activation of both NF- κ B and IRF3 transcription factors in MyD88^{-/-} mice that ultimately leads to the expression of IFN- β and, subsequently, to IFN- β -dependent STAT1 activation (1). It includes, for instance, a number of antiviral genes; Th1-activating chemokines, such as monokine induced by IFN- γ /CXCL9, IFN-inducible protein-10/CXCL10, and IFN-inducible T cell α chemoattractant/CXCL11; and anticancer molecules, such as TRAIL (9). Along the MyD88-independent pathway, TRAM (TIR domain-containing adapter-

*Section of General Pathology, Department of Pathology and Diagnostics, University of Verona, 37134 Verona, Italy; [†]Department of Public Health and Microbiology, University of Torino, 10126 Turin, Italy; and [‡]Servizio Trasfusionale, Azienda Ospedaliera Universitaria Integrata di Verona, 37134 Verona, Italy

Received for publication October 17, 2011. Accepted for publication May 21, 2012.

This work was supported by grants from Ministero dell'Istruzione, dell'Università e della Ricerca, Associazione Italiana per la Ricerca sul Cancro, and Fondazione Cassa di Risparmio (to M.A.C.). N.T. holds a fellowship from Fondazione Italiana per la Ricerca sul Cancro.

Address correspondence and reprint requests to Dr. Marco A. Cassatella, Department of Pathology and Diagnostics, Section of General Pathology, University of Verona, Strada Le Grazie 4, 37134 Verona, Italy. E-mail address: marco.cassatella@univr.it

The online version of this article contains supplemental material.

Abbreviations used in this article: ChIP, chromatin immunoprecipitation; DAI, DNA-dependent activator of IRF; IFI16, IFN-inducible protein 16; LRRFIP1, leucine-rich repeat (in Flightless I) interacting protein-1; MNE, mean normalized expression; MOI, multiplicity of infection; PDTC, pyrrolidine dithiocarbamate; pEGFP, enhanced GFP plasmid; PI, propidium iodide; PKC, protein kinase C; poly(dA:dT), poly(deoxyadenylic-deoxythymidyllic) acid; poly(I:C), polyinosinic-polycytidylic acid; PRL, prolactin; qPCR, quantitative PCR; RIG-I, retinoic acid-inducible gene; STING, stimulator of IFN genes; TBK1, TRAF family-associated NF- κ B-binding kinase-binding kinase-1; TRAM, TIR domain-containing adapter-inducing IFN- β -related adapter molecule; T4SS, type IV secretion system.

Copyright © 2012 by The American Association of Immunologists, Inc. 0022-1767/12/\$16.00

inducing IFN- β -related adapter molecule) and TRIF adapter proteins both transduce the activation of redundant protein kinases (i.e., TRAF family-associated NF- κ B-binding kinase-binding kinase-1 [TBK1] and I κ B kinase- ϵ), which phosphorylate IRF3 on Ser/Thr residues (10). As a result, IRF3 dimerizes, translocates to the nucleus, associates with other coactivators, and ultimately contributes to activate IFN- β gene transcription (11). Interestingly, the MyD88-independent/TRIF-dependent cascade seems to be initiated by a rapid activation of protein kinase C (PKC ϵ) in response to LPS, which, according to recent findings, would lead to phosphorylation of TRAM on serine residues (12). Such a post-translational modification would cause TRAM disappearance from the membrane concomitantly with activation of a downstream signaling cascade that would finally lead to activation of the TBK1-IRF3 axis and, ultimately, the transcriptional induction of IFN- β (12).

In this study, given the absence of PKC ϵ in human neutrophils (13–16), we verified whether the lack of PKC ϵ expression is the defect that prevents activation of the LPS-triggered MyD88-independent/TRIF-dependent pathway. For this purpose, we developed a transfection procedure to overexpress PKC ϵ in human neutrophils and ultimately test whether we could rescue the MyD88-independent/TRIF-dependent pathway. Although our experiments disproved a role for PKC ϵ in driving the MyD88-independent/TRIF-dependent response, they helped us to subsequently demonstrate that, similarly to all cell types tested to date, human neutrophils constitutively express various cytosolic DNA sensors. Consequently, we found that neutrophils can promptly express IFN- β mRNA upon recognition of transfected DNA or upon infection with intracellular pathogens, such as *Bartonella henselae*, *Listeria monocytogenes*, *Legionella pneumophila*, or adenovirus type 5.

Materials and Methods

Abs

PKC ϵ (sc-214), p-PKC ϵ (Ser⁷²⁹; sc-12355), PKC β II (sc-210), IRF3 (sc-9082), NF- κ B p65 (sc-372), NF- κ B p50 (sc-7178), I κ B- α (sc-371), IFN-inducible protein 16 (IFI16; sc-8023), DDX41 (sc-166225), and leucine-rich repeat (in Flightless I) interacting protein-1 (LRRFIP1; sc-135917) Abs were from Santa Cruz Biotechnology (Santa Cruz, CA). TBK1 (no. 3013), p-IRF3 (Ser³⁹⁶; no. 4947), p-NF- κ B p65 (Ser⁵³⁶; no. 3031), and p-p44/42 MAPK (no. 9106) Abs were from Cell Signaling (Beverly, MA); anti- β -tubulin (T5293) and anti-actin (A5060) Abs were from Sigma (St. Louis, MO). TBK1 mAbs (IMG-139A) and rabbit polyclonal stimulator of IFN genes (STING) Abs (IMG-6422A) were from IMGENEX (San Diego, CA); rabbit polyclonal ISG15 Abs were kindly provided by Dr. Arthur L. Haas (Louisiana State University Health Sciences Center, New Orleans, LA).

Cell purification and culture

Highly purified granulocytes (neutrophils > 96.5%, eosinophils < 3%, $n = 30$) and Percoll-purified monocytes ($n = 15$) were isolated and prepared under endotoxin-free conditions from buffy coats of healthy donors, as previously described (17). Ficoll-Paque-isolated neutrophils were further enriched by positively removing any eventual contaminating cells, to reach > 99.7% purity (high purity neutrophils), at least once for each type of experiment presented in this article (18). Immediately after purification, neutrophils were either subjected to the transfection procedure (see below) or suspended in RPMI 1640 medium supplemented with 10% low-endotoxin FBS (<0.5 EU/ml; BioWhittaker, Verviers, Belgium), treated or not with stimuli (including 100 ng/ml ultrapure *Escherichia coli* LPS [0111:B4; Alexis], 10 μ M R848 [InvivoGen], 50 μ g/ml polyinosinic-polycytidylic acid [poly(I:C); InvivoGen], 10 ng/ml PMA [Sigma], or 100 U/ml IFN- γ [R&D Systems, Minneapolis, MN]), and then plated either in 6/24-well tissue culture plates (Nunc, Roskilde, Denmark) or in polystyrene flasks (Orange, Trasadingen, Switzerland) to be cultured at 37°C, 5% CO₂ atmosphere. After the desired incubation period, cells were collected and spun at 300 \times g for 5 min, and cell pellets were either extracted for total RNA or lysed for protein analysis, as described below.

HEK293T cells (German Collection of Microorganism and Cell Cultures, Braunschweig, Germany) were cultured in 24-well plates using DMEM medium (Lonza) supplemented with 10% FBS and transfected with 2 μ l Lipofectamine 2000 (Invitrogen) complexed with 0.8 μ g poly(deoxyadenylic-deoxythymidylic) acid [poly(dA:dT); Sigma] or enhanced GFP plasmid (pEGFP; Clontech). All reagents used were of the highest available grade and were dissolved in pyrogen-free water for clinical use.

Transfection of neutrophils

Neutrophils were transfected with different types of plasmids, including pmaxGFP (Lonza), pEGFP, and pEGFP-PKC ϵ (kindly provided by Professor P. Parker, Cancer Research UK, London Research Institute, London, U.K.) previously purified using the EndoFree Plasmid Maxi Kit (QIAGEN), as well as with *E. coli* DNA (#D4889; RNA and contaminant free, according to the Sigma datasheet, as well as to our own checking), poly(dA:dT), and poly(I:C). Transfection was performed using the human monocyte nucleofector kit and the Amaxa nucleofector II device (both from Lonza), according to the reagent instructions, yet with minor modifications. After isolation, 5 \times 10⁶ neutrophils were resuspended in 100 μ l complete nucleofector solution containing 2.5–15 μ g DNA (or equal volumes of PBS for mock transfection) and then transferred to a nucleoporation cuvette. Electroporation was performed using the Y001 program in the nucleofector II device. Cells were then recovered and left for 5 min in 2 ml human monocyte nucleofector medium supplemented with 2 mM glutamine and 10% FBS. Thereafter, neutrophils were washed once with PBS to remove broken cells and subjected to flow cytometry analysis using a FACScan flow cytometer (Becton-Dickinson, Franklin Lakes, NJ) to determine GFP expression. In selected experiments, transfected neutrophils were incubated for 30 min with different inhibitors, including BAY 117082, pyrrolidine dithiocarbamate (PDTTC), sc-514 (Calbiochem, San Diego, CA), or 10 μ M MG132 (Sigma) prior to stimulation with LPS.

Quantitative PCR

RNA isolation and reverse transcription were accomplished as previously described (19). Quantitative PCR (qPCR) was performed using SYBR Premix Ex Taq (Takara) and gene-specific primers (Invitrogen, Carlsbad, CA) available in the public database RTPrimerDB (<http://medgen.ugent.be/rtprimerdb/index.php>) under the following entry codes: IFN- β (3542), TNF- α (3551), CXCL8/IL-8 (3553), CXCL10/IFN-inducible protein-10 (3537), G1P2/ISG15 (3547), IFIT1 (3540), and GAPDH (3539). Data were calculated with Q-Gene software (<http://www.gene-quantification.de/download.html>) and are expressed as mean normalized expression (MNE) units after GAPDH normalization.

Apoptosis assessment

Apoptosis of transfected neutrophils was determined, as previously described (5), by the propidium iodide (PI) staining procedure, according to the “quick method” described by Riccardi and Nicoletti (20). Data were analyzed by FlowJo software (TreeStar, Ashland, OR).

Immunoprecipitations and immunoblots

Whole-cell extracts were prepared for immunoblots using the RNeasy mini kit (QIAGEN), as described (21), or according to the chromatin immunoprecipitation (ChIP) assay procedure (22) for IFI16, LRRFIP1, and DDX41 immunoprecipitations. Cytoplasmic extracts for PKC ϵ and TBK1 immunoprecipitations were prepared using the nitrogen cavitation procedure and processed exactly as described previously (23). Blotted proteins were detected and quantified using the Odyssey infrared imaging system (LI-COR Biosciences, Lincoln, NE) (24).

ChIP assays

ChIP experiments were performed as described elsewhere (22), with minor modifications. For immunoprecipitation of transcription factors, nuclear extracts from 10⁷ sonicated neutrophils were incubated with 7 μ g anti-IRF3 or 5 μ g anti-NF- κ B p65/p50 Abs, whereas for cytoplasmic DNA sensor/plasmid DNA coimmunoprecipitations, whole-cell extracts were prepared from 5 \times 10⁶ pEGFP-transfected neutrophils were immunoprecipitated using 5 μ g α DDX41, α IFI16, and α LRRFIP1 Abs. In all ChIP experiments, protein recruitment to the prolactin (PRL) promoter (a gene that is completely silent in our cell types) was used as negative control. The coimmunoprecipitated material was then subjected to qPCR analysis using the following specific primers (purchased from Invitrogen): IFN- β promoter forward: 5'-TCCCAGGAAGTCAATGAAGG-3', IFN- β promoter reverse: 5'-GTGTCGCAATGGAGTGTGT-3'; CXCL8 promoter forward: 5'-CTTAGTGGGGTTGAAAGTGAC-3', CXCL8 promoter reverse: 5'-AAGAAATAGTCACTCACCCAAG-3'; pEGFP-backbone

forward: 5'-ACGGCATCAAGGTGAACCTTC-3', pEGFP-backbone reverse: 5'-GCTTCTCGTTGGGGTCTTTT-3'; pEGFP-GFP forward: 5'-TGCC-ATAGCCTCAGGTTACTC-3', pEGFP-GFP reverse: 5'-GACGCTCAGTG-GAACGAAAAC-3'; and PRL promoter forward: 5'-AGGGAACGAA-TGCCTGATT-3', PRL promoter reverse: 5'-GCAGGAAACACTTCA-CCA-3'. pEGFP-backbone and pEGFP-GFP primers of pEGFP (accession number: U55763, <http://www.ncbi.nlm.nih.gov/genbank/>) amplify the 3854–3999-bp and 1091–1259-bp regions, respectively.

Infection of neutrophils

A total of 2×10^6 high-purity neutrophils was infected with 10 multiplicity of infection (MOI) *L. monocytogenes*, *L. pneumophila*, or *B. henselae* and then cultured in 24-well tissue culture plates for up to 6 h prior to RNA extraction and qPCR analysis. At 1 h postinfection, 50 μ g/ml gentamicin was added to limit the growth of extracellular bacteria. In other experiments, neutrophils were infected with 1000 MOI adenovirus type 5 (E1 deleted, E3 defective, encoding for GFP; a kind gift of Prof. B. M. Foxwell, Imperial College, London, U.K.). Bacterial strains were all grown to mid-logarithmic phase as follows: wild-type *L. monocytogenes* (ATCC 19115) were grown in brain–heart infusion medium (Difco, Detroit, MI); *L. pneumophila* (serogroup 1 NCTC 12821, kindly provided by Prof. M.C. Zotti, University of Torino) were grown in N-(2-acetamido)-2-aminoethanesulfonic acid buffered yeast extract broth supplemented with 0.4 mg/ml L-cysteine and 0.135 mg/ml ferric nitrate (Sigma); and *B. henselae* Houston-1 (ATCC 49882) were grown in Schneider's medium supplemented with 10% FCS and 2 mM glutamine at 37°C with 5% CO₂ (25). Prior to infection, bacteria were pelleted and washed twice with PBS.

Statistical analysis

Data are expressed as mean \pm SE. Statistical evaluation was performed using the Student *t* test or one-way ANOVA followed by the Tukey post hoc test. The *p* values < 0.05 were considered statistically significant.

Results

Overexpression of PKC ϵ in human neutrophils by electroporation

Immunoblot and immunoprecipitation analyses of either cytoplasmic cavities (26) (Supplemental Fig. 1A, *top panel*) or whole-cell lysates (data not shown) confirmed (13–16) that neutrophils do not express PKC ϵ , unlike autologous monocytes. Moreover, PKC ϵ protein expression in neutrophils and monocytes was not inducible or upregulated, respectively, by 24 h-treatment with either LPS or IFN- γ (Supplemental Fig. 1B). Therefore, to clarify whether the absence of PKC ϵ might be responsible for the inability of human neutrophils to mobilize the MyD88-independent/TRIF-dependent pathway, we attempted to overexpress PKC ϵ in these cells. However, a series of preliminary experiments was necessary to develop a reliable transfection protocol, because neutrophils are extremely difficult to manipulate for exogenous gene expression as a result of their inability to proliferate and to survive in culture for long periods. Ultimately, we achieved success by modifying a recently reported nucleofection protocol (27), with which the investigators were able to transfect both GFP and p47^{phox}-GFP proteins into neutrophils. By our procedure, we transfected a pmaxGFP plasmid into neutrophils with a greater efficiency than the reported 5% (27) (see the fluorescence microscopy image in Supplemental Fig. 2A, *left panel*). Accordingly, flow cytometry analysis revealed that the percentage of GFP⁺ neutrophils correlated with the amounts of transfected pmaxGFP plasmid (Supplemental Fig. 2B). For example, >50% of GFP⁺ neutrophils were measured by nucleofecting 15 μ g pmaxGFP plasmid (Supplemental Fig. 2B). Importantly, such remarkable levels of GFP⁺ neutrophils were obtained as early as 4 h postnucleofection, and they remained stable for up to 20 h of culture (Supplemental Fig. 2C). However, because neutrophils became mostly apoptotic after 20 h (nearly 80% were PI⁺; Supplemental Fig. 2D), whereas they remained alive for up to 7–8 h posttransfection (75–80% of PI⁻ cells) (Supplemental Fig. 2D), all subsequent experiments were performed using neutrophils that

were first transfected for 4 h and then stimulated with LPS for no more than 3 h.

The validity of our experimental procedure was confirmed by transfecting neutrophils with a plasmid encoding a GFP-tagged full-length PKC ϵ (pEGFP-PKC ϵ), which produced de novo expression of remarkable quantities of exogenous PKC ϵ -GFP (Fig. 1A, 1B). Indeed, pEGFP-PKC ϵ -transfected neutrophils displayed specific PKC ϵ -immunoreactive signals, unlike the cells transfected with the corresponding empty plasmid (pEGFP) (Fig. 1A). Again, expression levels of exogenous PKC ϵ correlated with the amounts of transfected plasmid (Fig. 1B); for instance, the levels of exogenous PKC ϵ -GFP expressed in 2×10^6 neutrophils transfected with 15 μ g PKC ϵ -GFP plasmid were comparable to the levels of endogenous PKC ϵ present in 1×10^6 monocytes




FIGURE 1. Transgenic expression of PKC ϵ -GFP in human neutrophils. **(A)** Neutrophils were electroporated with 15 μ g pEGFP-PKC ϵ , 15 μ g pmaxGFP, or PBS (i.e., mock electroporated), as described in *Materials and Methods*. After 4 h, neutrophil and monocyte whole-cell extracts were prepared and, together with extracts from HEK293T cells, were transfected with pEGFP-PKC ϵ or not and electrophoresed (50 μ g neutrophils, 30 μ g monocytes, 10 μ g for HEK293T) for immunoblotting studies using Abs specific for PKC ϵ and PKC β II. Panels depict a representative experiment of three independent ones. **(B)** Neutrophils were electroporated with 5 or 15 μ g of pEGFP-PKC ϵ or with 15 μ g of pEGFP. After 4 h, whole-cell extracts were prepared and then electrophoresed for PKC ϵ and actin immunoblotting. Panels depict a representative experiment (*n* = 2). **(C)** Neutrophils were electroporated with 15 μ g of pEGFP-PKC ϵ for 4 h, treated with 5 ng/ml PMA, and then lysed after an additional 20 min. Whole-cell extracts were prepared, electrophoresed, and immunoblotted using Abs specific for p-PKC ϵ , total PKC ϵ , p-ERK(p42/p44), and actin. **(D)** Whole-cell extracts prepared from monocytes stimulated for 20 min with 5 ng/ml PMA were processed as in (C) and used as positive control. For (C) and (D), data show one representative experiment of two.

(Fig. 1A, 1B). Importantly, PKC ϵ -GFP overexpressed in neutrophils underwent serine phosphorylation in response to PMA stimulation, as revealed by the use of specific anti-p-PKC ϵ Abs (Fig. 1C). Similar data could be extrapolated by using anti-PKC ϵ Abs that detect a doublet (28), with the more slowly migrating band likely corresponding to phosphorylated PKC ϵ , as clearly evidenced in monocytes (Fig. 1D). Importantly, transfected neutrophils displayed an unaltered capacity to respond to external agonists, as proven by the findings that PMA triggered ERK phosphorylation in electroporated neutrophils at levels very similar to those observed in nontransfected cells (Fig. 1C). Taken together, our data demonstrate that it is possible to successfully transfect PKC ϵ -GFP in neutrophils and that exogenous PKC ϵ -GFP can be rapidly phosphorylated by PMA.

LPS acquires the capacity to upregulate IFN- β mRNA expression in plasmid DNA-transfected neutrophils

To assess whether transfected PKC ϵ could restore the MyD88-independent/TRIF-dependent pathway, neutrophils were incubated 4 h post-PKC ϵ -GFP plasmid or pEGFP plasmid transfection and then stimulated with 100 ng/ml ultrapure LPS for an additional 90 min (Fig. 2A, upper panel) to investigate the state of IFN- β and CXCL10 mRNA expression by qPCR. In line with our previous study (8), no IFN- β or CXCL10 mRNA induction occurred in nontransfected neutrophils stimulated with LPS, as opposed to the induction of two classical MyD88-dependent genes: *TNF- α* and *CXCL8* (Fig. 2A, lower panels). In contrast, both IFN- β and CXCL10 (but not *TNF- α* or *CXCL8*) gene expression was significantly increased by LPS in PKC ϵ -GFP-overexpressing neutrophils (Fig. 2A, lower panels), as if the MyD88-independent/TRIF-dependent pathway was rescued by the introduction of an exogenous PKC ϵ . However, this presumed rescue also occurred in neutrophils transfected with a pEGFP plasmid, with increased amounts of IFN- β and CXCL10 transcripts in response to LPS at levels comparable to those measured in PKC ϵ -GFP-overexpressing neutrophils (Fig. 2A, center panels). Unexpectedly, neutrophils transfected with either PKC ϵ -GFP or pEGFP plasmids expressed elevated amounts of IFN- β , CXCL10, *TNF- α* , and *CXCL8* transcripts, even in the absence of LPS stimulation (Fig. 2A), indicating that the simple introduction of DNA into neutrophils strongly promotes their gene activation. Accordingly, data depicted in Fig. 2B clearly demonstrate that, upon transfection with pEGFP plasmid (or PKC ϵ -GFP plasmid or *E. coli* DNA, data not shown), a time-dependent increase in IFN- β and CXCL10 mRNA is already detectable in neutrophils after 45 min, which increases thereafter up to 7 h. Also, neutrophil electroporation with poly(dA:dT), a chemically synthesized DNA commonly used to study the immunological response to B-form DNA (29), directly triggered IFN- β and CXCL10 mRNA expression (Fig. 2C, lower panels). The latter experiments not only demonstrate that the ability of neutrophils to recognize foreign DNA is not restricted to plasmid DNA, but they also prove that the stimulating agent is genuine DNA and not a contaminating product derived from the plasmid-isolation procedure (e.g., bacterial RNA). Moreover, additional stimulation with LPS of poly(dA:dT)-electroporated neutrophils again resulted in an upregulation in the levels of IFN- β and CXCL10 transcripts relative to nontransfected neutrophils (Fig. 2C, lower panels). Such an LPS-mediated effect was also observed in neutrophils transfected with *E. coli* DNA (Fig. 2C, lower panels) but, interestingly, not with poly(I:C) (Fig. 2C, lower panels), a synthetic analog of viral dsRNA that we previously showed to very strongly upregulate IFN- β mRNA expression when electroporated into human neutrophils (5). Altogether, our data demonstrate that the mere DNA/plasmid DNA transfection




FIGURE 2. Modulation of IFN- β mRNA expression in plasmid-transfected neutrophils incubated or not with LPS. Neutrophils were electroporated with 15 μ g pEGFP-PKC ϵ , 15 μ g pEGFP (empty vector) (A, B), 15 μ g of poly(dA:dT), 15 μ g *E. coli* DNA, 2 μ g of poly(I:C), or PBS (mock electroporated) (C) and then incubated for 4 h before the addition of 100 ng/ml LPS (A, C) or incubated for the times indicated (B). Total RNA was extracted after an additional 90 min (A) or 180 min (C) post-LPS stimulation or as indicated (B) and then analyzed for IFN- β , CXCL10, *TNF- α* , *CXCL8*, and GAPDH mRNA expression by qPCR. The schemes illustrating the experimental protocols used are shown in the uppermost panels of (A) and (C). Gene expression is depicted as MNE \pm SE ($n = 6$) after GAPDH normalization of triplicate reactions for each sample. For (B) and (C), results shown are from one representative experiment of at least three performed. Asterisks indicate a significant increase exerted by LPS. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

tion of neutrophils markedly activates the expression of IFN- β and CXCL10 mRNA. Our data also demonstrate that LPS further upregulates the expression of IFN- β and CXCL10 mRNA in DNA/plasmid DNA-transfected neutrophils, yet in a manner that appears independent from the overexpression of exogenous PKC ϵ -GFP.

Identification of IRF3 as the critical factor for the transcription of IFN- β mRNA in plasmid DNA-transfected neutrophils

Subsequent experiments were aimed at elucidating the molecular bases responsible for the induction of IFN- β mRNA expression in neutrophils transfected with plasmid DNA, as well as for its further upregulation by LPS. For this purpose, we analyzed the activation status of IRF3, which represents a crucial intermediate for the transcriptional induction of IFN- β (11). Using the protocol depicted in the scheme of Fig. 3A (*upper panel*), we found that the mere pEGFP transfection of neutrophils triggered a direct ser396 phosphorylation of IRF3 (Fig. 3A, *center panel*) that was relatively weak after 2 h but became very strong after 5 h (for a quantitative densitometric analysis see Fig. 3A, *lower panel*, $n = 3$). Notably, similar results were obtained with the PKC ϵ -GFP plasmid (data not shown), whereas a 1-h stimulation of DNA-transfected neutrophils with LPS did not augment the levels of ser396-phosphorylated IRF3 (Fig. 3A). Consistent with these findings were the results from IRF3 ChIP assays, which revealed a strong recruitment of IRF3 to the IFN- β promoter in plasmid-transfected neutrophils (Fig. 3B, *lower panel*) and that such recruitment is not increased further by subsequent LPS stimulation (Fig. 3B, *lower panel*). The specificity of the IRF3 ChIP assay was proved by the fact that the PRL promoter was not amplified under the same experimental conditions (Fig. 3B, *lower panel*), as well as by the fact that matched control Abs did not coprecipitate any IFN- β or PRL promoter (data not shown). Altogether, data demonstrate that, although the transfection of neutrophils with plasmid DNA directly activates the phosphorylation of IRF3 and its binding to the IFN- β promoter, LPS remains unable to modify the activation status of IRF3, even if it upregulates the accumulation of IFN- β transcripts. It follows that the upregulatory effect of LPS at the level of IFN- β gene expression in plasmid-transfected neutrophils does not occur via the MyD88-independent/TRIF-dependent cascade.

Identification of the transcription factors promoting the expression of IFN- β mRNA in plasmid DNA-transfected neutrophils upon LPS stimulation: role of NF- κ B

Because NF- κ B is a transcription factor cooperating with IRF3 in promoting IFN- β transcription (30), we subsequently analyzed its activation status in plasmid-transfected neutrophils treated or not with LPS for 45 min (Fig. 4A, 4B). By doing so, we observed considerable levels of phosphorylated NF- κ B p65 (Fig. 4A) and diminished amounts of I κ B α (Fig. 4B) in neutrophils transfected with DNA plasmids for 4 h and 45 min. Moreover, the addition of LPS to 4-h plasmid-transfected neutrophils resulted in a further increase in the levels of NF- κ B p65 phosphorylation (Fig. 4A) and I κ B α degradation (Fig. 4B) after 45 min (for a quantitative densitometric analysis see Fig. 4A, 4B, *lower panels*, $n = 3$). Importantly, ChIP assays performed according to the scheme depicted in Fig. 4C, using Abs toward NF- κ B p65 (Fig. 4D) and NF- κ B p50 (Fig. 4E), highlighted a crucial role for NF- κ B in driving the effect of LPS on IFN- β mRNA expression in plasmid-transfected neutrophils. In fact, both p50 and p65 NF- κ B subunits were strongly recruited to the IFN- β promoter upon LPS stimulation of plasmid-transfected neutrophils (Fig. 4D, 4E, *left panels*). In contrast, no recruitment of either NF- κ B p50 or NF- κ B p65 to the IFN- β




FIGURE 3. Activation of IRF3 in plasmid-transfected neutrophils. **(A)** Neutrophils were electroporated with 15 μ g pEGFP and cultured for up to 5 h, the last hour in the presence or the absence of 100 ng/ml LPS. Neutrophils were lysed at the 2- and 5-h time points (*uppermost panel*) and whole-cell extracts were electrophoresed and immunoblotted using anti-p-IRF3 (Ser³⁹⁶), anti-IRF3, and anti-actin Abs. The *center panel* shows a representative immunoblot ($n = 3$), and the *lower panel* displays the densitometric quantification of p-IRF3 levels (normalized by total IRF3; $n = 3$) by LI-COR Odyssey software, expressed as densitometric units (D.U.). **(B)** Neutrophils were electroporated with 15 μ g pEGFP and cultured for 5 h, the last hour in the presence or absence of 100 ng/ml LPS. Cells were then processed for ChIP analysis, as described in *Materials and Methods*. Enrichment of IFN- β and PRL promoters in coprecipitated DNA was analyzed by qPCR using promoter-specific primers. Data from qPCR are expressed as percentages over input DNA (mean \pm SE; $n = 3$). * $p < 0.05$, ** $p < 0.01$ versus nontransfected cells.

promoter was observed in nontransfected neutrophils treated with LPS (Fig. 4D, 4E, *left panels*), consistent with the inability of LPS to induce IFN- β gene expression under the latter conditions (8). However, under the latter conditions, both NF- κ B p65 and NF- κ B p50 bound strongly to the CXCL8 promoter (Fig. 4D, 4E, *right panels*). Notably, in plasmid-transfected neutrophils, the recruitment of NF- κ B p65 and NF- κ B p50 to the CXCL8 promoter, although negligible after 60 min of LPS stimulation (Fig. 4D, 4E, *right panels*), was ≥ 3 -fold higher after 30 min (data not shown), in line with the enhanced CXCL8 mRNA expression observed in parallel (Fig. 2A).

The role of activated NF- κ B as a crucial mediator of the additional upregulation of IFN- β mRNA expression by LPS in




FIGURE 4. LPS-mediated NF- κ B activation and binding to the IFN- β promoter in plasmid-transfected neutrophils. (A, B) Neutrophils were electroporated with 15 μ g pEGFP and cultured for 4 h and 45 min, with the last 45 min in the presence or absence of 100 ng/ml LPS (as depicted in the top panels). Whole-cell extracts were prepared, electrophoresed, and immunoblotted using anti-p-NF- κ B p65 (Ser⁵³⁶), anti-total NF- κ B p65 (A), anti-I κ B α , and anti-actin Abs (B). Shown is a representative immunoblot split into two parts ($n = 3$) (center panels). The graphs (lower panels) illustrate the densitometric quantification of p-NF- κ B p65 levels (normalized by the total NF- κ B p65) (A) and I κ B α levels (normalized by the total actin) (B), expressed as mean \pm SE ($n = 3$). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Neutrophils were treated as in (C), and processed for ChIP analysis using anti-NF- κ B p65 (D) or anti-NF- κ B p50 (E) Abs. Enrichment in IFN- β , CXCL8, and PRL promoter in coprecipitated DNA was analyzed by qPCR using promoter-specific primers. Data from qPCR are expressed as percentages over input DNA (mean \pm SE; $n = 3$). ** $p < 0.01$, *** $p < 0.001$ versus control or nontransfected cells.

plasmid-transfected neutrophils was additionally supported by the effect of four chemical NF- κ B blockers: BAY 117082, an inhibitor of I κ B α phosphorylation and degradation (31); PDTC, an inhibitor of NF- κ B activation (32); sc-514, a selective I κ B kinase-2

inhibitor (33) (Fig. 5); and MG132, a proteasomal inhibitor (34) (data not shown). In fact, although they had no effect on the induction of IFN- β mRNA directly triggered by the simple plasmid transfection, all of them significantly reduced the upregulatory effect of LPS on IFN- β mRNA expression (Fig. 5). Altogether, our data suggest that the increased expression of IFN- β mRNA observed in response to LPS in plasmid-transfected neutrophils is likely the consequence of the LPS-triggered recruitment of p65 and p50 NF- κ B subunits to the IFN- β promoter, which, for unknown reasons, does not take place in nontransfected neutrophils.

Intracellular recognition of plasmid DNA by human neutrophils is dependent on IFI16

Finally, we aimed at defining how human neutrophils recognize transfected plasmid DNA. Initially, we investigated the role of TLR9, because this molecule is constitutively expressed and responsive in human neutrophils (4, 35). TLR9 recognizes unmethylated CG dinucleotides (CpG) present in the extracellular space within the endosomal compartment (36); consequently, drugs able to block endosome acidification (37), such as chloroquine and bafylomycin A1, inhibit its ability to activate NF- κ B and to induce cytokines (37, 38). Experiments in which neutrophils were treated with the two inhibitors for 30 min prior to plasmid DNA electroporation (for 4 h) revealed that both chloroquine and bafylomycin A1 do not prevent the induction of IFN- β or TNF- α gene expression (Supplemental Fig. 3A, left and center panels). In contrast, both inhibitors markedly suppressed the induction of TNF- α mRNA expression in R848-stimulated neutrophils (Supplemental Fig. 3A, right panel), thereby proving that they were functioning appropriately. Then we tested the role of RNA polymerase III, which functions as an indirect intracellular DNA sensor (39, 40). In fact, RNA polymerase III recognizes AT-rich DNA and consequently synthesizes uncapped 5' triphosphate-bearing RNA, which, in turn, serves as an agonist for retinoic acid-inducible gene (RIG-I) (39, 40), which is constitutively expressed by human neutrophils (Fig. 6A) (5). Once again, a specific RNA polymerase III inhibitor-based approach, using ML-60218 (39, 40), did not suppress the induction of IFN- β mRNA exerted by plasmid DNA or poly(dA:dT) electroporation (Supplemental Fig. 3B). In contrast, ML-60218 dose dependently reduced the




FIGURE 5. Effect of NF- κ B inhibitors on the upregulation of IFN- β mRNA expression triggered by LPS in plasmid-transfected neutrophils. Neutrophils were electroporated with 15 μ g of pEGFP; after 3.5 h of culture, they were incubated with 300 μ M PDTC, 100 μ M sc-514, or 5 μ M BAY 117082. After 30 min, neutrophils were stimulated with 100 ng/ml LPS for an additional 90 min (upper panel). Total RNA was then extracted and analyzed for IFN- β and GAPDH mRNA expression by qPCR. IFN- β mRNA expression is depicted as MNE \pm SE ($n = 3$) after GAPDH normalization of triplicate reactions for each sample. ** $p < 0.01$, *** $p < 0.001$.




FIGURE 6. Neutrophil recognition of intracellular plasmid DNA is dependent on IFI16. (A) Whole neutrophil and monocyte extracts (60 μg) were electrophoresed and immunoblotted using Abs specific for IFI16, LRRFIP1, DDX41, RIG-I, STING, and tubulin. (B) Whole neutrophil lysates (500 μg), prepared using the ChIP assay buffer, were subjected to immunoprecipitation with IFI16, LRRFIP1, DDX41, or anti-mouse IgG1 (control) Abs. Representative immunoblot experiments are shown (n = 3) for (A) and (B). (C) Neutrophils were transfected with pEGFP, cultured for 3 and 5 h, and then disrupted by ChIP assay buffer. Whole-cell lysates were then subjected to immunoprecipitation with IFI16, LRRFIP1, DDX41, or anti-mouse IgG1 control Abs. Finally, coimmunoprecipitated DNA was purified and analyzed by qPCR using two sets of primers specific for pEGFP (pEGFP-backbone and pEGFP-GFP). Data from qPCR are expressed as percentages over input DNA. Results are from one experiment representative of four.

induction of IFN-β mRNA in HEK293T cells transfected with poly(dA:dT) (Supplemental Fig. 3C), as expected (39, 40). Thus, we concluded that neither TLR9 nor RNA polymerase III functions as the intracellular receptor(s) recognizing plasmid DNA in electroporated neutrophils. In contrast, we confirmed that neutrophils express IFI16 (41), one of the newly identified intracellular receptors for foreign DNA (42). Accordingly, immunoblotting experiments revealed that whole neutrophil and monocyte lysates contain IFI16, as a cluster of proteins of 85–95 kDa (Fig. 6A), similar to what was originally described in the nuclear extracts of IFN-γ-treated HL-60 cells (41). Under the same conditions, we also detected, in both neutrophils and monocytes, two more recently discovered DNA sensors: LRRFIP1, which promotes IFN-β gene expression through β-catenin engagement and not via IRF3 activation (43), and DDX41, a member of the DEXDC family of helicases (44) (Fig. 6A). Furthermore, we found that neu-

trophils also express STING (Fig. 6A), a transmembrane protein located in the endoplasmic reticulum that is crucial for DNA-mediated signaling (45–48).

In an attempt to identify the specific DNA sensor(s) binding transfected DNA in neutrophils, we used the strategy that allowed Zhang et al. (44) to identify the intracellular binding protein recognizing *L. monocytogenes* DNA. Specifically, we immunoprecipitated IFI16, LRRFIP1, and DDX41 after DNA transfection (Fig. 6B) and then analyzed, by qPCR, to which of these receptors plasmid DNA was stably bound (Fig. 6C). By doing so, we found that IFI16 functions as the intracellular DNA sensor that, in transfected neutrophils, is predominantly involved in recognizing plasmid DNA (Fig. 6C).

Induction of IFN-β mRNA expression by infection of neutrophils with intracellular pathogens

In a final series of experiments, we tested a number of intracellular pathogens, including *B. henselae*, *L. monocytogenes*, *L. pneumophila*, and adenovirus type 5, which, according to the literature (49) are known to activate IFN-β gene expression via cytosolic sensing of their DNA. As shown in Fig. 7A, all intracellular bacteria under examination triggered a remarkable IFN-β and CXCL10 mRNA expression in neutrophils infected for 6 h but not 3 h. In addition, all bacteria upregulated the expression of TNFα (Fig. 7A) and CXCL8 mRNA (data not shown), indicating that




FIGURE 7. Induction of IFN-β mRNA expression in neutrophils infected by various intracellular pathogens. (A) Neutrophils were infected for 3 and 6 h with *B. henselae*, *L. monocytogenes*, or *L. pneumophila* at MOI 10. Total RNA was then extracted and analyzed for IFN-β, CXCL10, TNF-α, and GAPDH mRNA expression by qPCR. (B) Neutrophils were infected with adenovirus type 5 at MOI 1000. Total RNA was extracted after 1.5 and 5 h of incubation and analyzed for IFN-β, CXCL10, GFP, and GAPDH mRNA expression by qPCR. Gene expression in (A) and (B) is depicted as MNE after GAPDH normalization of triplicate reactions for each sample (MNE ± SE). Depicted results are from one experiment representative of at least three independent ones.

they are able to activate a wide spectrum of neutrophil genes upon their recognition. A time-dependent induction of IFN- β and CXCL10 mRNA expression was also observed in neutrophils infected with adenovirus type 5 (Fig. 7B). The latter phenomenon was genuine, as testified by the amplification of transcripts encoding GFP, which are virus specific (Fig. 7B).

Discussion

A characteristic feature that differentiates human neutrophils from monocytes in terms of LPS responsiveness is that neutrophils are unable to mobilize the MyD88-independent/TRIF-dependent cascade upon TLR4 engagement (8). In this study, in light of the identification of PKC ϵ as a crucial molecule for initiating the MyD88-independent/TRIF-dependent pathway (12), and having confirmed that PKC ϵ is undetectable in human neutrophils (13–16), we attempted to overexpress PKC ϵ with the aim to restore the MyD88-independent/TRIF-dependent pathway. By optimizing a previously described electroporation method (27) to obtain >50% GFP⁺ cells, we were able to successfully transfect PKC ϵ in neutrophils. However, despite the very high amounts of exogenous PKC ϵ protein expressed by transfected neutrophils, the results of our subsequent experiments did not permit us to conclude that overexpressed PKC ϵ is instrumental in rescuing the MyD88-independent/TRIF-dependent pathway. In fact, although LPS acquired the capacity to upregulate the expression of IFN- β and CXCL10 mRNA (as well as ISG15 and IFIT-1 mRNA [N. Tamassia and M.A. Cassatella, unpublished observations]) in PKC ϵ -transfected neutrophils, a similar phenomenon was observed in neutrophils transfected with *E. coli* DNA, poly(dA:dT), or a variety of empty plasmids but not in mock-transfected cells. Moreover, the fact that LPS did not modify the state of IRF3 activation/phosphorylation in either plasmid- or PKC ϵ -transfected neutrophils further demonstrates that the MyD88-independent/TRIF-dependent cascade was not activated in DNA-nucleofected neutrophils. Assuming that overexpressed PKC ϵ was fully functional in neutrophils (in this regard, we show that PKC ϵ underwrote serine phosphorylation upon cell treatment with PMA), an additional implication of our results is that PKC ϵ may not be as critical for the activation of the MyD88-independent/TRIF-dependent cascade as previously proposed (12). Consistent with such a notion, the role of PKC ϵ in driving the MyD88-independent/TRIF-dependent response was also questioned recently by Parker and colleagues (50), who demonstrated that PKC ϵ associates with MyD88 and that PKC ϵ phosphorylation is important for NF- κ B activation via TLR2, which, unlike TLR4, does not use the TRIF or TRAM adaptors (51).

Whatever the case may be, we also show in this study that the mere transfection of neutrophils with plasmid DNA (regardless of the presence of a PKC ϵ -coding region within the vector) directly activates IRF3, promotes its recruitment to the IFN- β promoter, and, in turn, strongly induces the expression of IFN- β and other type I IFN-dependent genes (such as CXCL10, ISG15, and IFIT-1). Together with the findings described above, our data imply that, in plasmid-transfected neutrophils, LPS becomes able to further upregulate the expression of IFN- β mRNA without directly targeting IRF3. ChIP assays confirmed this hypothesis, as well as established the presumed molecular mechanisms underlying the induction of IFN- β mRNA, either by plasmid transfection or by LPS in plasmid-transfected neutrophils. Accordingly, we found that plasmid DNA transfection promotes a time-dependent recruitment of IRF3 to the IFN- β promoter; the addition of LPS to plasmid-transfected neutrophils does not increase such IRF3 recruitment; although cytosolic DNA markedly activates NF- κ B in the cytoplasm, it is unable to promote NF- κ B

recruitment at the IFN- β promoter; LPS promotes the binding of NF- κ B to the IFN- β promoter in plasmid-transfected neutrophils but not in nontransfected cells; and NF- κ B inhibitors significantly suppress the upregulatory effect of LPS on IFN- β mRNA expression in transfected neutrophils. Based on these findings, we propose that the capacity of LPS to increase the expression of IFN- β mRNA in plasmid-transfected neutrophils relies on its ability to activate NF- κ B, which, under conditions in which IRF3 is already present on the IFN- β promoter (i.e., after plasmid transfection), becomes readily recruited (see the scheme depicted in Supplemental Fig. 4). In fact, given the close proximity of the IRF3 binding site to the NF- κ B locus within the IFN- β promoter, it is possible that binding of NF- κ B occurs only when IRF3 is already bound. In this regard, it was demonstrated that binding of IRF3 to the IFN- β promoter produces a nucleosome shift that functions as a prerequisite for the binding of NF- κ B and AP1 to the IFN- β promoter (52) (i.e., the nucleosome remodeling exerted by IRF3 might unmask the NF- κ B binding sites present in the IFN- β promoter, in this way allowing NF- κ B binding and consequent increased IFN- β transcription). Altogether, our data are in accordance with the view that, even if not essential unlike IRF3 (11), NF- κ B plays an important role in modulating the degree of IFN- β promoter activation (11, 53). This is also in line with other experimental systems in which the induction of IFN- β results from a cooperative action of two different signals (54, 55). Similarly, the present data are consistent with previous findings demonstrating that human neutrophils often require stimulation by two concurrent, but different, stimulatory pathways to optimally express a given gene [e.g., IFN- γ plus LPS for CXCL10 (56) and IL-12 (57) expression or IL-10 plus LPS for IL-1ra (22) expression]. Curiously, LPS was unable to further upregulate the levels of IFN- β and CXCL10 transcripts in poly(I:C)-transfected neutrophils. Although we did not specifically explore, at the molecular levels, the reasons for this, some speculations can be made. First, because we showed that poly(I:C) transfection itself represents one of the most potent stimulatory conditions for gene expression in neutrophils (5), it is plausible that IFN- β and CXCL10 mRNA levels are already induced to the maximum and, therefore, cannot be increased further. Second, and in agreement with the former effects, poly(I:C) transfection activates NF- κ B so powerfully (5) that LPS cannot further activate it. However, the fact that LPS maintains the ability to promote lactoferrin release (58) in poly(I:C)-transfected neutrophils (N. Tamassia and M.A. Cassatella, unpublished observations) indicates that, under the latter experimental conditions, TLR4 is expressed and functional.

This study also demonstrates that human neutrophils can promptly respond to transfected DNA, whether of bacterial or synthetic origin. In this context, our observations complement a series of previous studies describing the capacity of neutrophils to respond to exogenous DNA. Neutrophils, in fact, can readily respond to CpG oligodeoxynucleotides in terms of CXCL8 production and inhibition of apoptosis because they express their cognate receptor, TLR9 (4). Other findings demonstrated that extracellular bacterial DNA can trigger, through an unidentified receptor, another pathway functioning in a TLR9- and CpG-independent, but MyD88-dependent, manner (59). Our new observations additionally suggest that human neutrophils possess intracellular sensor system(s) that allow(s) the recognition of foreign and potentially dangerous DNA and, consequently, the induction of a distinct and potent immune response (60). The intracellular sensors that recognize exogenous DNA include TLR9 (36), RIG-I through RNA polymerase III (39, 40), absent in melanoma 2 (61–64), LRRFIP1 (43), DNA-dependent activator of IRFs (DAI) (65), IFI16 (42), and DDX41 (44). In this study, we provide evidence

that human neutrophils constitutively express, in addition to RIG-I (5), DDX41, LRRFIP1, and, confirming previous findings (41), IFI16. We also show that neutrophils constitutively express STING, a transmembrane protein essential for the signaling necessary for the production of IFN- β , via IRF3 activation, mediated by various cytosolic DNA sensors (45, 66). In this context, we did not investigate the role of DAI, the first identified cytosolic DNA sensor (65), because wild-type and DAI-deficient cells produce comparable amounts of type I IFN in response to cytosolic DNA (67) or poly(dA:dT) (68). We also did not focus on absent in melanoma 2, a member of the pyrin and HIN200 domain-containing protein family, because, based on the literature, this receptor is not essential for type I IFN induction by transfected DNA, whereas it is crucial for inflammasome activation and IL-1 β secretion (61–64). Moreover, we excluded the involvement of both TLR9 and RNA polymerase III by using specific inhibitors. However, by performing coimmunoprecipitation studies of DDX41, LRRFIP1, and IFI16, and, presumably bound plasmid DNA, we were able to identify IFI16 as the critical intracellular sensor for plasmid DNA in human neutrophils. The latter findings are consistent with the biological features of IFI16, which, differently from TLR9, DAI, and RNA polymerase III, does not recognize specific DNA structure (42), and, which, through the activation of NF- κ B and IRF3, is known to induce a strong production of proinflammatory cytokines and type I IFN in response to transfection with DNA motifs (42). In any case, the presence of RIG-I, DDX41, and LRRFIP1 in neutrophils provides a solid molecular basis to explain our additional findings on the ability of *L. monocytogenes*, *L. pneumophila*, and adenovirus type 5 to induce the expression of IFN- β and other ISG mRNA in infected neutrophils, because those pathogens require, for such a function, DDX41 plus LRRFIP1, RIG-I, and DDX41 plus RIG, respectively. In contrast, we show that neutrophils also express IFN- β mRNA following infection with *B. henselae*, a Gram-negative facultative intracellular microorganism that can invade many cells (69, 70). *Bartonella* uses several virulence factors for its interaction with host cells, including the VirB/D4 type IV secretion system (T4SS), which was recently shown to mediate plasmid DNA and protein transfer into eukaryotic host cells (71). Because VirB/VirD4 T4SS is related to the dot/Icm T4SS used by *L. pneumophila*, which is essential to trigger IFN- β expression (72), likely via the RNA polymerase III/RIG cascade (40), it is tempting to speculate that *B. henselae* induces IFN- β mRNA through mechanisms similar to those used by *L. pneumophila*.

Regardless, further studies are necessary to meticulously decipher which neutrophil sensor(s) is/are responsible for the recognition of foreign DNA from different sources and in which cell compartment they locate to do so. Nonetheless, the findings of this study shed new light on our understanding of the mechanisms by which human neutrophils recognize and respond to intracellular pathogens and, in turn, activate innate immune responses.

Acknowledgments

We thank Dr. Arthur L. Haas, Prof. P. Parker, and Prof. B.M. Foxwell for providing the anti-ISG15 Ab, the pEGFP-PKCe plasmid, and adenovirus type 5, respectively.

Disclosures


The authors have no financial conflicts of interest.

References

1. Kawai, T., and S. Akira. 2010. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat. Immunol.* 11: 373–384.


2. O'Neill, L. A. 2004. TLRs: Professor Mechnikov, sit on your hat. *Trends Immunol.* 25: 687–693.
3. Muzio, M., D. Bosisio, N. Polentarutti, G. D'amico, A. Stoppacciaro, R. Mancinelli, C. van't Veer, G. Penton-Rol, L. P. Ruco, P. Allavena, and A. Mantovani. 2000. Differential expression and regulation of toll-like receptors (TLR) in human leukocytes: selective expression of TLR3 in dendritic cells. *J. Immunol.* 164: 5998–6004.
4. Hayashi, F., T. K. Means, and A. D. Luster. 2003. Toll-like receptors stimulate human neutrophil function. *Blood* 102: 2660–2669.
5. Tamassia, N., V. Le Moigne, M. Rossato, M. Donini, S. McCartney, F. Calzetti, M. Colonna, F. Bazzoni, and M. A. Cassatella. 2008. Activation of an immunoregulatory and antiviral gene expression program in poly(I:C)-transfected human neutrophils. *J. Immunol.* 181: 6563–6573.
6. Poltorak, A., X. He, I. Smirnova, M. Y. Liu, C. Van Huffel, X. Du, D. Birdwell, E. Alejos, M. Silva, C. Galanos, et al. 1998. Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. *Science* 282: 2085–2088.
7. Mantovani, A., M. A. Cassatella, C. Costantini, and S. Jaillon. 2011. Neutrophils in the activation and regulation of innate and adaptive immunity. *Nat. Rev. Immunol.* 11: 519–531.
8. Tamassia, N., V. Le Moigne, F. Calzetti, M. Donini, S. Gasperini, T. Ear, A. Cloutier, F. O. Martinez, M. Fabbri, M. Locati, et al. 2007. The MyD88-independent pathway is not mobilized in human neutrophils stimulated via TLR4. *J. Immunol.* 178: 7344–7356.
9. Hirota, T., M. Yamamoto, Y. Kumagai, S. Uematsu, I. Kawase, O. Takeuchi, and S. Akira. 2005. Regulation of lipopolysaccharide-inducible genes by MyD88 and Toll/IL-1 domain containing adaptor inducing IFN- β . *Biochem. Biophys. Res. Commun.* 328: 383–392.
10. Fitzgerald, K. A., S. M. McWhirter, K. L. Faia, D. C. Rowe, E. Latz, D. T. Golenbock, A. J. Coyle, S. M. Liao, and T. Maniatis. 2003. IKKepsilon and TBK1 are essential components of the IRF3 signaling pathway. *Nat. Immunol.* 4: 491–496.
11. Hiscott, J., P. Pitha, P. Genin, H. Nguyen, C. Heylbroeck, Y. Mamane, M. Algarte, and R. Lin. 1999. Triggering the interferon response: the role of IRF-3 transcription factor. *J. Interferon Cytokine Res.* 19: 1–13.
12. McGettrick, A. F., E. K. Brint, E. M. Palsson-McDermott, D. C. Rowe, D. T. Golenbock, N. J. Gay, K. A. Fitzgerald, and L. A. O'Neill. 2006. Trif-related adapter molecule is phosphorylated by PKCepsilon during Toll-like receptor 4 signaling. *Proc. Natl. Acad. Sci. USA* 103: 9196–9201.
13. Dang, P. M., J. Hakim, and A. Périani. 1994. Immunochemical identification and translocation of protein kinase C zeta in human neutrophils. *FEBS Lett.* 349: 338–342.
14. Kent, J. D., S. Sergeant, D. J. Burns, and L. C. McPhail. 1996. Identification and regulation of protein kinase C-delta in human neutrophils. *J. Immunol.* 157: 4641–4647.
15. Khwaja, A., and L. Tatton. 1999. Caspase-mediated proteolysis and activation of protein kinase Cdelta plays a central role in neutrophil apoptosis. *Blood* 94: 291–301.
16. O'Flaherty, J. T., B. A. Chadwell, M. W. Kearns, S. Sergeant, and L. W. Daniel. 2001. Protein kinases C translocation responses to low concentrations of arachidonic acid. *J. Biol. Chem.* 276: 24743–24750.
17. Tecchio, C., V. Huber, P. Scapini, F. Calzetti, D. Margotto, G. Todeschini, L. Pilla, G. Martinelli, G. Pizzolo, L. Rivoltini, and M. A. Cassatella. 2004. IFNalpha-stimulated neutrophils and monocytes release a soluble form of TNF-related apoptosis-inducing ligand (TRAIL/Apo-2 ligand) displaying apoptotic activity on leukemic cells. *Blood* 103: 3837–3844.
18. Pelletier, M., L. Maggi, A. Micheletti, E. Lazzari, N. Tamassia, C. Costantini, L. Cosmi, C. Lunardi, F. Annunziato, S. Romagnani, and M. A. Cassatella. 2010. Evidence for a cross-talk between human neutrophils and Th17 cells. *Blood* 115: 335–343.
19. Cassatella, M. A., V. Huber, F. Calzetti, D. Margotto, N. Tamassia, G. Peri, A. Mantovani, L. Rivoltini, and C. Tecchio. 2006. Interferon-activated neutrophils store a TNF-related apoptosis-inducing ligand (TRAIL/Apo-2 ligand) intracellular pool that is readily mobilizable following exposure to proinflammatory mediators. *J. Leukoc. Biol.* 79: 123–132.
20. Riccardi, C., and I. Nicoletti. 2006. Analysis of apoptosis by propidium iodide staining and flow cytometry. *Nat. Protoc.* 1: 1458–1461.
21. Morse, S. M., G. Shaw, and S. F. Lerner. 2006. Concurrent mRNA and protein extraction from the same experimental sample using a commercially available column-based RNA preparation kit. *Biotechniques* 40: 54–, 56, 58.
22. Tamassia, N., M. Castellucci, M. Rossato, S. Gasperini, D. Bosisio, M. Giacomelli, R. Badolato, M. A. Cassatella, and F. Bazzoni. 2010. Uncovering an IL-10-dependent NF-kappaB recruitment to the IL-1ra promoter that is impaired in STAT3 functionally defective patients. *FASEB J.* 24: 1365–1375.
23. McDonald, P. P., A. Bald, and M. A. Cassatella. 1997. Activation of the NF-kappaB pathway by inflammatory stimuli in human neutrophils. *Blood* 89: 3421–3433.
24. Gatto, L., C. Berlato, V. Poli, S. Tininini, I. Kinjyo, A. Yoshimura, M. A. Cassatella, and F. Bazzoni. 2004. Analysis of SOCS-3 promoter responses to interferon gamma. *J. Biol. Chem.* 279: 13746–13754.
25. Riess, T., F. Dietrich, K. V. Schmidt, P. O. Kaiser, H. Schwarz, A. Schäfer, and V. A. Kempf. 2008. Analysis of a novel insect cell culture medium-based growth medium for *Bartonella* species. *Appl. Environ. Microbiol.* 74: 5224–5227.
26. McDonald, P. P., C. Bovolenta, and M. A. Cassatella. 1998. Activation of distinct transcription factors in neutrophils by bacterial LPS, interferon-gamma, and GM-CSF and the necessity to overcome the action of endogenous proteases. *Biochemistry* 37: 13165–13173.

27. Johnson, J. L., B. A. Ellis, D. B. Munafo, A. A. Brzezinska, and S. D. Catz. 2006. Gene transfer and expression in human neutrophils. The phox homology domain of p47phox translocates to the plasma membrane but not to the membrane of mature phagosomes. *BMC Immunol.* 7: 28.
28. Tourkina, E., P. Gooz, J. C. Oates, A. Ludwicka-Bradley, R. M. Silver, and S. Hoffman. 2004. Curcumin-induced apoptosis in scleroderma lung fibroblasts: role of protein kinase cepsilon. *Am. J. Respir. Cell Mol. Biol.* 31: 28–35.
29. Ishii, K. J., C. Coban, H. Kato, K. Takahashi, Y. Torii, F. Takeshita, H. Ludwig, G. Sutter, K. Suzuki, H. Hemmi, et al. 2006. A Toll-like receptor-independent antiviral response induced by double-stranded B-form DNA. *Nat. Immunol.* 7: 40–48.
30. Thanos, D., and T. Maniatis. 1995. Virus induction of human IFN beta gene expression requires the assembly of an enhancosome. *Cell* 83: 1091–1100.
31. Pierce, J. W., R. Schoenleber, G. Jesmok, J. Best, S. A. Moore, T. Collins, and M. E. Gerritsen. 1997. Novel inhibitors of cytokine-induced I κ B α phosphorylation and endothelial cell adhesion molecule expression show anti-inflammatory effects in vivo. *J. Biol. Chem.* 272: 21096–21103.
32. Schreck, R., B. Meier, D. N. Männel, W. Dröge, and P. A. Baeuerle. 1992. Dithiocarbamates as potent inhibitors of nuclear factor kappa B activation in intact cells. *J. Exp. Med.* 175: 1181–1194.
33. Kishore, N., C. Sommers, S. Mathialagan, J. Guzova, M. Yao, S. Hauser, K. Huynh, S. Bonar, C. Mielke, L. Albee, et al. 2003. A selective IKK-2 inhibitor blocks NF-kappa B-dependent gene expression in interleukin-1 beta-stimulated synovial fibroblasts. *J. Biol. Chem.* 278: 32861–32871.
34. Grisham, M. B., V. J. Palombella, P. J. Elliott, E. M. Conner, S. Brand, H. L. Wong, C. Pien, L. M. Mazzola, A. Destree, L. Parent, and J. Adams. 1999. Inhibition of NF-kappa B activation in vitro and in vivo: role of 26S proteasome. *Methods Enzymol.* 300: 345–363.
35. József, L., T. Khreiss, and J. G. Filep. 2004. CpG motifs in bacterial DNA delay apoptosis of neutrophil granulocytes. *FASEB J.* 18: 1776–1778.
36. Hemmi, H., O. Takeuchi, T. Kawai, T. Kaisho, S. Sato, H. Sanjo, M. Matsumoto, K. Hoshino, H. Wagner, K. Takeda, and S. Akira. 2000. A Toll-like receptor recognizes bacterial DNA. *Nature* 408: 740–745.
37. Yi, A. K., R. Tuetken, T. Redford, M. Waldschmidt, J. Kirsch, and A. M. Krieg. 1998. CpG motifs in bacterial DNA activate leukocytes through the pH-dependent generation of reactive oxygen species. *J. Immunol.* 160: 4755–4761.
38. Rutz, M., J. Metzger, T. Gellert, P. Lippa, G. B. Lipford, H. Wagner, and S. Bauer. 2004. Toll-like receptor 9 binds single-stranded CpG-DNA in a sequence- and pH-dependent manner. *Eur. J. Immunol.* 34: 2541–2550.
39. Ablasser, A., F. Bauernfeind, G. Hartmann, E. Latz, K. A. Fitzgerald, and V. Hornung. 2009. RIG-I-dependent sensing of poly(dA:dT) through the induction of an RNA polymerase III-transcribed RNA intermediate. *Nat. Immunol.* 10: 1065–1072.
40. Chiu, Y. H., J. B. Macmillan, and Z. J. Chen. 2009. RNA polymerase III detects cytosolic DNA and induces type I interferons through the RIG-I pathway. *Cell* 138: 576–591.
41. Dawson, M. J., and J. A. Trapani. 1995. IFI 16 gene encodes a nuclear protein whose expression is induced by interferons in human myeloid leukaemia cell lines. *J. Cell. Biochem.* 57: 39–51.
42. Unterholzer, L., S. E. Keating, M. Baran, K. A. Horan, S. B. Jensen, S. Sharma, C. M. Sirois, T. Jin, E. Latz, T. S. Xiao, et al. 2010. IFI16 is an innate immune sensor for intracellular DNA. *Nat. Immunol.* 11: 997–1004.
43. Yang, P., H. An, X. Liu, M. Wen, Y. Zheng, Y. Rui, and X. Cao. 2010. The cytosolic nucleic acid sensor LRRFIP1 mediates the production of type I interferon via a beta-catenin-dependent pathway. *Nat. Immunol.* 11: 487–494.
44. Zhang, Z., B. Yuan, H. Bao, N. Lu, T. Kim, and Y. J. Liu. 2011. The helicase DDX41 senses intracellular DNA mediated by the adaptor STING in dendritic cells. *Nat. Immunol.* 12: 959–965.
45. Ishikawa, H., and G. N. Barber. 2008. STING is an endoplasmic reticulum adaptor that facilitates innate immune signalling. *Nature* 455: 674–678.
46. Zhong, B., Y. Yang, S. Li, Y. Wang, Y. Li, F. Diao, C. Lei, X. He, L. Zhang, P. Tien, and H. B. Shu. 2008. The adaptor protein MITA links virus-sensing receptors to IRF3 transcription factor activation. *Immunity* 29: 538–550.
47. Jin, L., P. M. Waterman, K. R. Jonscher, C. M. Short, N. A. Reisdorph, and J. C. Cambier. 2008. MPYS, a novel membrane tetraspanner, is associated with major histocompatibility complex class II and mediates transduction of apoptotic signals. *Mol. Cell. Biol.* 28: 5014–5026.
48. Sun, W., Y. Li, L. Chen, H. Chen, F. You, X. Zhou, Y. Zhou, Z. Zhai, D. Chen, and Z. Jiang. 2009. ERIS, an endoplasmic reticulum IFN stimulator, activates innate immune signaling through dimerization. *Proc. Natl. Acad. Sci. USA* 106: 8653–8658.
49. Keating, S. E., M. Baran, and A. G. Bowie. 2011. Cytosolic DNA sensors regulating type I interferon induction. *Trends Immunol.* 32: 574–581.
50. Faisal, A., A. Saurin, B. Gregory, B. Foxwell, and P. J. Parker. 2008. The scaffold MyD88 acts to couple protein kinase Cepsilon to Toll-like receptors. *J. Biol. Chem.* 283: 18591–18600.
51. Fitzgerald, K. A., D. C. Rowe, B. J. Barnes, D. R. Caffrey, A. Visintin, E. Latz, B. Monks, P. M. Pitha, and D. T. Golenbock. 2003. LPS-TLR4 signaling to IRF-3/7 and NF-kappaB involves the toll adapters TRAM and TRIF. *J. Exp. Med.* 198: 1043–1055.
52. Lomvardas, S., and D. Thanos. 2002. Modifying gene expression programs by altering core promoter chromatin architecture. *Cell* 110: 261–271.
53. Yang, H., G. Ma, C. H. Lin, M. Orr, and M. G. Wathlet. 2004. Mechanism for transcriptional synergy between interferon regulatory factor (IRF)-3 and IRF-7 in activation of the interferon-beta gene promoter. *Eur. J. Biochem.* 271: 3693–3703.
54. Smith, J. A., M. J. Turner, M. L. DeLay, E. I. Klenk, D. P. Sowders, and R. A. Colbert. 2008. Endoplasmic reticulum stress and the unfolded protein response are linked to synergistic IFN-beta induction via X-box binding protein 1. *Eur. J. Immunol.* 38: 1194–1203.
55. Negishi, H., Y. Fujita, H. Yanai, S. Sakaguchi, X. Ouyang, M. Shinohara, H. Takayanagi, Y. Ohba, T. Taniguchi, and K. Honda. 2006. Evidence for licensing of IFN-gamma-induced IFN regulatory factor 1 transcription factor by MyD88 in Toll-like receptor-dependent gene induction program. *Proc. Natl. Acad. Sci. USA* 103: 15136–15141.
56. Tamassia, N., F. Calzetti, T. Ear, A. Cloutier, S. Gasperini, F. Bazzoni, P. P. McDonald, and M. A. Cassatella. 2007. Molecular mechanisms underlying the synergistic induction of CXCL10 by LPS and IFN-gamma in human neutrophils. *Eur. J. Immunol.* 37: 2627–2634.
57. Cassatella, M. A., L. Meda, S. Gasperini, A. D'Andrea, X. Ma, and G. Trinchieri. 1995. Interleukin-12 production by human polymorphonuclear leukocytes. *Eur. J. Immunol.* 25: 1–5.
58. Koivuranta-Vaara, P., D. Banda, and I. M. Goldstein. 1987. Bacterial-lipopolysaccharide-induced release of lactoferrin from human polymorphonuclear leukocytes: role of monocyte-derived tumor necrosis factor alpha. *Infect. Immun.* 55: 2956–2961.
59. Alvarez, M. E., J. I. Fuxman Bass, J. R. Geffner, P. X. Calotti, M. Costas, O. A. Coso, R. Gamberale, M. E. Vermeulen, G. Salamone, D. Martinez, et al. 2006. Neutrophil signaling pathways activated by bacterial DNA stimulation. *J. Immunol.* 177: 4037–4046.
60. Hornung, V., and E. Latz. 2010. Intracellular DNA recognition. *Nat. Rev. Immunol.* 10: 123–130.
61. Hornung, V., A. Ablasser, M. Charrel-Dennis, F. Bauernfeind, G. Horvath, D. R. Caffrey, E. Latz, and K. A. Fitzgerald. 2009. AIM2 recognizes cytosolic dsDNA and forms a caspase-1-activating inflammasome with ASC. *Nature* 458: 514–518.
62. Bürckstümmer, T., C. Baumann, S. Blüml, E. Dixit, G. Dürnberger, H. Jahn, M. Planyavsky, M. Bilban, J. Colinge, K. L. Bennett, and G. Superti-Furga. 2009. An orthogonal proteomic-genomic screen identifies AIM2 as a cytoplasmic DNA sensor for the inflammasome. *Nat. Immunol.* 10: 266–272.
63. Fernandez-Alnemri, T., J. W. Yu, P. Datta, J. Wu, and E. S. Alnemri. 2009. AIM2 activates the inflammasome and cell death in response to cytoplasmic DNA. *Nature* 458: 509–513.
64. Roberts, T. L., A. Idris, J. A. Dunn, G. M. Kelly, C. M. Burton, S. Hodgson, L. L. Hardy, V. Garceau, M. J. Sweet, I. L. Ross, et al. 2009. HIN-200 proteins regulate caspase activation in response to foreign cytoplasmic DNA. *Science* 323: 1057–1060.
65. Takaoka, A., Z. Wang, M. K. Choi, H. Yanai, H. Negishi, T. Ban, Y. Lu, M. Miyagishi, T. Kodama, K. Honda, et al. 2007. DAI (DLM-1/ZBP1) is a cytosolic DNA sensor and an activator of innate immune response. *Nature* 448: 501–505.
66. Ishikawa, H., Z. Ma, and G. N. Barber. 2009. STING regulates intracellular DNA-mediated, type I interferon-dependent innate immunity. *Nature* 461: 788–792.
67. Ishii, K. J., T. Kawagoe, S. Koyama, K. Matsui, H. Kumar, T. Kawai, S. Uematsu, O. Takeuchi, F. Takeshita, C. Coban, and S. Akira. 2008. TANK-binding kinase-1 delineates innate and adaptive immune responses to DNA vaccines. *Nature* 451: 725–729.
68. Wang, Z., M. K. Choi, T. Ban, H. Yanai, H. Negishi, Y. Lu, T. Tamura, A. Takaoka, K. Nishikura, and T. Taniguchi. 2008. Regulation of innate immune responses by DAI (DLM-1/ZBP1) and other DNA-sensing molecules. *Proc. Natl. Acad. Sci. USA* 105: 5477–5482.
69. Pulliainen, A. T., and C. Dehio. 2012. Persistence of *Bartonella* spp. stealth pathogens: from subclinical infections to vasoproliferative tumor formation. *FEMS Microbiol. Rev.* 36: 563–599.
70. Vermi, W., F. Facchetti, E. Riboldi, H. Heine, S. Scutera, S. Stornello, D. Ravarino, P. Cappello, M. Giovarelli, R. Badolato, et al. 2006. Role of dendritic cell-derived CXCL13 in the pathogenesis of *Bartonella henselae* B-rich granuloma. *Blood* 107: 454–462.
71. Schröder, G., R. Schuelein, M. Quebatte, and C. Dehio. 2011. Conjugative DNA transfer into human cells by the VirB/VirD4 type IV secretion system of the bacterial pathogen *Bartonella henselae*. *Proc. Natl. Acad. Sci. USA* 108: 14643–14648.
72. Stetson, D. B., and R. Medzhitov. 2006. Recognition of cytosolic DNA activates an IRF3-dependent innate immune response. *Immunity* 24: 93–103.




Supplemental Figure 1. PKC ϵ protein is not detectable in human neutrophils. (A) Cytoplasmic extracts prepared by nitrogen cavitation (see M&M) from neutrophils and autologous monocytes were either electrophoresed (50 μ g neutrophils, 30 μ g monocytes) or immunoprecipitated (500 μ g from neutrophils, 300 μ g from monocytes) with specific anti-PKC ϵ and TBK1. Samples were also immunoprecipitated with anti-rabbit IgG control antibodies. A representative immunoblot experiment out of three with similar results is shown. Panel (A) shows that PKC ϵ is not detectable in either cytoplasmic neutrophil cavitates (*upper* panel). On the other hand, TBK1, which has a MW comparable to that of PKC ϵ , is readily detected in neutrophil samples (*lower* panel), proving that our samples were correctly prepared. (B) Neutrophils and autologous monocytes were incubated with 100 ng/ml LPS or 100 U/ml IFN γ for 24 h. Whole cell extracts were then prepared, electrophoresed (50 μ g neutrophils, 30 μ g monocytes) and immunoblotted using antibodies specific for PKC ϵ , ISG15 and actin. Panels display one experiment representative of three with similar results. Data demonstrate that the levels of ISG15, but not PKC ϵ , protein are upregulated by IFN γ in both neutrophils and monocytes, as well as by LPS in monocytes only, in line with the fact that ISG15 is a “MyD88-independent/TRIF-dependent” gene.

Supplemental Figure 2




Supplemental Figure 2. Expression of transfected green fluorescent protein (GFP) in human neutrophils. (A) Human neutrophils were electroporated with a 15 µg GFP-encoding plasmid (pmaxGFP) as described in M&M and, after 6 h, cell images were captured by a fluorescence microscope, either in fluorescence (left panel), or in bright field (right panel), mode. (B) Neutrophils were electroporated with different amounts of pmaxGFP plasmid prior to measuring GFP expression levels after a 4 h-culture, by flow cytometry. Values within each plot indicate the percentage of GFP-positive neutrophils. A representative experiment out of three with similar results is shown. (C) Neutrophils were electroporated with 15 µg pmaxGFP plasmid and expression of GFP protein was detected after 4, 6 and 20 h by flow cytometry. Values (mean ± SE, n=3) indicate the percentage of GFP-positive neutrophils. (D) Neutrophils were treated as described for panel C to evaluate their apoptotic state after different incubation times, as measured by the propidium iodide staining method. Values are expressed as mean ± SE (n= 3).

Supplemental Figure 3



Supplemental Figure 3. Recognition of intracellular DNA by human neutrophils is independent from TLR9 or RNA polymerase III. (A) Neutrophils were pre-treated for 30 min with 5 μ g/ml chloroquine or 25 nM bafilomycin A1 and then either electroporated with 15 μ g of pEGFP (*left* and *center* panels), or incubated with 100 ng/ml LPS or 10 μ M R848 for 90 min (*right* panel). After 5 h (*left* and *center* panels), total RNA was extracted and analyzed for IFN β (*left* panels), TNF α (*right* and *center* panels) and GAPDH mRNA expression by RT-qPCR. (B) Neutrophils were pre-treated for 30 min with 30 μ M ML-60218 and then electroporated with either 15 μ g pEGFP or 15 μ g poly(dA:dT). After 5 h, total RNA was extracted and analyzed for IFN β and GAPDH mRNA expression by RT-qPCR. (C) HEK 293T cells were pre-treated for 2 h with different doses of ML-60218 and then transfected with 0.8 μ g/ml poly(dA:dT) complexed with lipofectamine. After 24 h, total RNA was extracted and analyzed for IFN β and β 2-microglobulin (β 2m) mRNA expression by RT-qPCR. For all panels, gene expression is depicted as MNE after GAPDH (A-B) or β 2m (C) normalization of triplicate reactions for each sample. Results are from one experiment representative of at least three independent ones.

Supplemental Figure 4



Supplemental Figure 4. Intracellular plasmid DNA activates IRF3 and favours the recruitment of LPS-activated NF-κB to the IFN β promoter in human neutrophils. (A) Upon stimulation of human neutrophils with LPS, p65/p50 NF-κB heterodimers, even if activated *via* the MyD88-dependent pathway, do not bind to the IFN β promoter since their binding sites are likely masked. No IRF3 activation occurs in LPS-stimulated neutrophils since the TRIF-dependent pathway is not functional. **(B)** Upon transfection, plasmid DNA binds to IFI16 and activates a signal transduction cascade that leads to IRF3 phosphorylation, dimerization and nuclear translocation. IRF3, in turn, is recruited to its binding sites within the IFN β promoter to induce IFN β transcription. Concomitantly, IRF3 presumably unmask NF-κB recognition sequences within the same IFN β promoter *via* activation of unknown, local chromatin remodelling events. **(C)** As a result, p65/p50 NF-κB heterodimers activated by LPS in DNA-transfected neutrophils become now able to bind the unmasked NF-κB recognition sequences and, cooperating with IRF3, promote the recruitment of more RNA polymerase II and consequently increase IFN β mRNA transcription.