



Explore what's possible with innovative
research tools

Discover the difference>



Pathogenic *Mycobacterium tuberculosis* Evades Apoptosis of Host Macrophages by Release of TNF-R2, Resulting in Inactivation of TNF- α

This information is current as
of April 20, 2021.

M. Katarzyna Balcewicz-Sablinska, Joseph Keane, Hardy
Kornfeld and Heinz G. Remold

J Immunol 1998; 161:2636-2641; ;
<http://www.jimmunol.org/content/161/5/2636>

References This article **cites 41 articles**, 22 of which you can access for free at:
<http://www.jimmunol.org/content/161/5/2636.full#ref-list-1>

Why *The JI*? Submit online.

- **Rapid Reviews! 30 days*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

**average*

Subscription Information about subscribing to *The Journal of Immunology* is online at:
<http://jimmunol.org/subscription>

Permissions Submit copyright permission requests at:
<http://www.aai.org/About/Publications/JI/copyright.html>

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://jimmunol.org/alerts>

The Journal of Immunology is published twice each month by
The American Association of Immunologists, Inc.,
1451 Rockville Pike, Suite 650, Rockville, MD 20852
Copyright © 1998 by The American Association of
Immunologists All rights reserved.
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



Pathogenic *Mycobacterium tuberculosis* Evades Apoptosis of Host Macrophages by Release of TNF-R2, Resulting in Inactivation of TNF- α ¹

M. Katarzyna Balcewicz-Sablinska,* Joseph Keane,[†] Hardy Kornfeld,[†] and Heinz G. Remold^{2*}

Infection by *Mycobacterium tuberculosis* (MTB) induces human alveolar macrophage (AM ϕ) apoptosis by a TNF- α -dependent mechanism. The apoptotic response is postulated to be a defense mechanism, limiting the growth of this intracellular pathogen. Consistent with that model, recent studies showed that the virulent MTB strain H37Rv induces substantially less AM ϕ apoptosis than the attenuated strain H37Ra. We now report that AM ϕ infection with either H37Rv or H37Ra induces comparable levels of TNF- α measured by ELISA but that TNF- α bioactivity is reduced in supernatants of H37Rv-infected AM ϕ . Differential release of soluble TNFR2 (sTNFR2), with formation of inactive TNF- α -TNFR2 complexes accounted for the difference in TNF- α bioactivity in these cultures. Release of sTNFR2 by H37Rv-infected AM ϕ was IL-10 dependent since it was inhibited by neutralizing anti-IL-10 Ab. Thus, the effect of TNF- α produced by AM ϕ following infection can be modulated by virulent MTB, using IL-10 as an upstream mediator. *The Journal of Immunology*, 1998, 161: 2636–2641.

Tuberculosis is the predominant cause of human mortality from infectious agents, with a death rate of 3 million individuals per year worldwide. The number of new cases for the decade 1990 to 2000 is estimated at 90 million (1). Moreover, the severity of this global medical problem has worsened in recent years due to emergence of drug-resistant *Mycobacterium tuberculosis* (MTB)³ strains (2).

The primary route of infection with MTB is via the lung. In the alveolus, AM ϕ are both the primary host cell for MTB and the first line of defense for the infected individual. Following infection of AM ϕ by MTB, a variety of cytokines are induced that orchestrate containment of the infection. Among these cytokines, TNF- α is involved in the protective immune response in mice (3). Production of TNF- α is also required for the induction of AM ϕ apoptosis in response to inoculation with MTB (4). Comparison in mice of different MTB strains correlated resistance against tuberculosis in vivo with susceptibility of infected M ϕ to apoptosis measured in vitro and in vivo (5). Apoptosis of human monocytes was shown to limit the growth of *Mycobacterium bovis* bacille Calmette-Guérin (BCG) (6) and *Mycobacterium avium* (7) in vitro. The latter studies shed light on the outcomes of host M ϕ apoptosis. First, apoptosis of the infected cells contains the mycobacteria, likely preventing spread of infection in vivo (8). Secondly, the process of host M ϕ apoptosis prevents growth of the mycobacteria (7). In-

deed, H37Rv, a well-characterized pathogenic MTB strain, was found to induce significantly less host M ϕ apoptosis than the closely related attenuated strain H37Ra (4), suggesting that abrogation of host M ϕ apoptosis might be a virulence-associated phenotype of MTB strains.

The present study was performed to delineate the mechanism by which virulent MTB interferes with the host M ϕ apoptotic response to infection. We identify inactivation of the cytokine TNF- α as a key event leading to decreased host M ϕ apoptosis. We also describe proximal events: the induction of IL-10 production by virulent MTB leading to release of soluble TNFR2 (sTNFR2) from the AM ϕ and inactivation of TNF- α by complex formation with sTNFR2. Together, these events comprise an apparent virulence mechanism operative in virulent and not in attenuated MTB.

Materials and Methods

Alveolar macrophages

AM ϕ were obtained from bronchoalveolar lavage fluid of healthy non-smoking volunteers with their informed consent, at Boston University Medical Center or Beth Israel-Deaconess Medical Center, under human study protocols approved by the respective institutional review boards. Lavage fluids and cells were filtrated through a single layer of sterile gauze, centrifuged (450 \times g, 100 min), and the cell pellet was suspended in RPMI 1640 medium (Sigma, St. Louis, MO) with 10% FCS, supplemented with cefotaxime (50 μ g/ml, Sigma) or penicillin and streptomycin (100 μ g/ml each, Sigma). Cells were plated on 24-well plates (Costar, Cambridge, MA) (5×10^5 cells/well) directly on the bottom of wells or on round glass coverslips for staining purposes. Nonadherent cells were removed by washing at 24 h, and fresh antibiotic-free medium was added. Cell morphology and differential counts were determined with light microscopy following cytocentrifugation and modified Giemsa staining (Dif-Quik, Sigma). Viability of adherent alveolar macrophages was examined in representative samples by trypan blue dye exclusion.

Mycobacterium tuberculosis

Stock cultures of MTB H37Ra and H37Rv (ATCC 25177 and ATCC 25618; American Type Culture Collection (ATCC), Manassas, VA) were stored frozen in 1-ml aliquots. Before inoculation of AM ϕ , the thawed mycobacteria were vortexed, sonicated for 15 s at 500 W, and allowed to stand for 10 min. The upper 500 μ l of the solution were used for experiments. Dilution of these dispersed mycobacterial preparations were made to yield a multiplicity of infection 1 to 5 organisms/cell. For each experiment, the adequacy of dispersion and the multiplicity of infection were

*Division of Rheumatology and Immunology, Department of Internal Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston MA 02115; and [†]Pulmonary Center, Boston University School of Medicine, Boston, MA 02118

Received for publication December 17, 1997. Accepted for publication April 30, 1998.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This work was supported by National Institutes of Health Grant PO1 HL43510.

² Address correspondence and reprint requests to Dr. Heinz G. Remold, Division of Rheumatology and Immunology, Department of Internal Medicine, Brigham and Women's Hospital and Harvard Medical School, Smith Building, Room 526B, 75 Francis Street, Boston MA, 02115.

³ Abbreviations used in this paper: MTB, *Mycobacterium tuberculosis*; AM ϕ , alveolar macrophages; sTNFR, soluble TNFR; TUNEL, terminal deoxynucleotidyl transferase-mediated dUTP-nick end labeling.

checked by acid-fast staining of the washed cells 4 h after inoculation. The rate of internalization and intracellular growth during 5 days after infection was similar for H37Ra and H37Rv as determined by electron microscopy, plate counting, and BACTEC (9).

Reagents, Abs, and cytokines

Human rIL-10, polyclonal anti-human IL-10, and polyclonal anti-human TNF- α Abs were obtained from Endogen (Cambridge, MA). Human rTNF- α was purchased from Genzyme (Cambridge, MA). Human rsTNFR2 and monoclonal anti-human sTNFR2 Ab were purchased from R&D Systems (Minneapolis, MN).

Measurement of TNF- α , IL-10, sTNFR, and TNF- α -sTNFR2 complexes

AM ϕ were cultured and challenged with MTB as described above, in triplicate cultures. Supernatants were harvested at the times indicated and passed through 0.22- μ m pore-size filter (Costar). The level of immunoreactive TNF- α and IL-10 was determined by ELISA using matched Ab pairs and cytokine standards (Genzyme and Endogen, respectively). The amount of sTNFR was estimated using commercial ELISA kits (R&D Systems) according to manufacturer's instructions.

TNF bioactivity was assessed by measuring cytotoxicity on L929 cells (10). In brief, L929 cell monolayers, cultured for 48 h on flat-bottom 96-well plates (0.2×10^5 cells/well), were overlaid with twofold serial dilutions of AM ϕ supernatants in RPMI 1640 + 10% FCS supplemented with actinomycin D to a final concentration of 1 μ g/ml.

After incubation at 37°C for 24 h the cells were washed with saline, stained with 0.05% crystal violet in 20% ethanol, and allowed to dry overnight. The stain was eluted from the cells with 100% methanol and read with a microtiter plate reader at an absorbance of 570 nm. Estimates of the concentrations of bioactive TNF in the supernatants were obtained by comparison with calibration curves established with a rTNF- α standard. TNF bioactivity in selected AM ϕ supernatant samples was inhibited by anti-TNF- α Ab, but not by control rabbit IgG, indicating that the cytotoxic activity in AM ϕ supernatants represents TNF- α .

TNF- α bound to TNFR2 was detected in supernatants of MTB-infected AM ϕ by a mixed Ab ELISA assay. The supernatants were added to 96-well polystyrene microtiter plates coated with a murine monoclonal Ab against sTNFR2 (part of TNFR2 ELISA; R&D Systems), and, after 2 h at room temperature, the plates were washed and incubated for 1 h at room temperature with biotinylated rabbit anti-human TNF- α Ab (0.2 μ g/ml, Genzyme). After washing, peroxidase-labeled streptavidin (Genzyme) was added. Peroxidase activity was determined by the addition of 3,3',5,5'-tetramethylbenzidine (Vector Laboratories, Burlingame, CA), and photo-spectrometry (450 nm) was performed using a microplate reader. Since the stability of TNF- α -sTNFR2 complexes strongly depends on the experimental conditions (11), calibrated dilutions of rTNF- α captured by immobilized anti-TNF- α Ab were used as an internal standard for the comparative measurements of TNFR2-bound TNF- α .

In situ DNA fragmentation analysis

DNA fragmentation was determined by the in situ TUNEL technique (12). The method is based on the specific binding of terminal deoxynucleotidyl transferase to 3'-OH ends of DNA. AM ϕ adherent to glass coverslips were fixed with 4% formaldehyde in PBS, rinsed with double distilled water, immersed in 0.5 U/ μ l of terminal transferase and 1 mM Biotin-16-dUTP in reaction buffer (Boehringer Mannheim, Indianapolis, IN) and incubated in a humid atmosphere at 37°C for 60 min. The reaction was terminated by transferring the coverslips to a solution of 300 mM NaCl, 30 mM sodium citrate, pH 7.2, for 15 min at room temperature. The coverslips were then rinsed with water, blocked with 2% BSA in PBS for 10 min, rinsed again with water, immersed in PBS for 5 min at room temperature, and immersed in avidin peroxidase (1:10) for 30 min at 37°C. The cells were then stained with 1% 3-amino, 9-ethyl carbazole (Sigma) in dimethyl formamide (Sigma) diluted 1:20 with 50 mM acetate buffer, pH 4.5, containing 0.018% hydrogen peroxide for 20 min. Slides of experimental and control cells subjected to in situ TUNEL were analyzed by light microscopy in a blinded fashion. A total of 600 cells per slide were evaluated for the presence of red-brown nuclear staining.

Statistical analysis

Results are expressed as mean \pm SD. Statistical analysis was performed employing SigmaStat Statistical Software (Jandel Scientific, San Rafael, CA), using the ANOVA with correction for multiple comparisons for analyses for multiple comparisons and paired *t* test for analysis of the data on IL-10 production by AM ϕ from individual donors.




FIGURE 1. Quantitation of TNF- α Ag (A), TNF- α bioactivity (B), and apoptosis (C) in cultures of AM ϕ infected with virulent and attenuated MTB. Shown are mean \pm SD of three representative experiments. Bars labeled "CTRL" represent uninfected AM ϕ . A, AM ϕ were infected with H37Rv or H37Ra, and culture supernatants were assayed at 6 and 24 h for TNF- α Ag by ELISA ($p > 0.4$ for 6 and 24 h). B, Culture supernatants at 6 and 24 h after infection were assayed for TNF- α bioactivity by measuring cytotoxicity for L929 mouse fibroblasts. Dilutions of rTNF- α were used as internal standard. The differences in TNF- α cytotoxicity between H37Ra- and H37Rv-infected cultures were statistically significant at 6 h and 24 h after infection ($p < 0.05$). C, Apoptosis of host AM ϕ at 2 and 4 days after infection measured by TUNEL assay and shown as percent of apoptotic cells. The differences in % apoptosis between H37Ra- and H37Rv-infected cultures were statistically significant at day 2 and day 4 ($p < 0.05$).

Results

Inactivation of TNF- α in AM ϕ cultures infected with virulent MTB

Using AM ϕ from healthy donors, we measured TNF- α levels by ELISA in culture supernatants after infection with the virulent MTB strain H37Rv or the attenuated strain H37Ra at 6 and 24 h. The immunoreactive TNF- α levels increased over time in both infections and were not significantly different (Fig. 1A). In contrast, TNF- α bioactivity measured at the same time by assessing L929 fibroblast cytotoxicity was lower in H37Rv-infected AM ϕ cultures than in H37Ra-infected AM ϕ cultures (Fig. 1B). Twenty-four hours after infection, the level of TNF- α bioactivity was 412 ± 75 pg/ml in supernatants from H37Rv-infected AM ϕ vs 1573 ± 189 pg/ml in supernatants from H37Ra-infected cells ($n = 3$, $p < 0.05$). Control experiments confirmed that the measurements of bioactivity reflected TNF- α activity since the addition of anti-TNF- α Ab (5 μ g/ml) completely inhibited L929 fibroblast cytotoxicity in both cultures, whereas irrelevant Ab had no effect (data not shown). Consistent with the previous report that apoptosis in MTB-infected AM ϕ resulted from TNF- α signaling (4), the level of AM ϕ apoptosis mirrored the pattern of TNF- α bioactivity




FIGURE 2. Levels of soluble TNFR1 in supernatants from AM ϕ inoculated with H37Rv, H37Ra, or uninfected AM ϕ and harvested at the indicated times. Shown are mean \pm SD of triplicate values from one representative experiment of three.

(Fig. 1C). In contrast to the measurement of TNF- α by ELISA or by using a cytotoxicity assay, apoptosis was determined at day 2 and 4 because optimal manifestation of apoptosis as measured by the TUNEL technique (12) was found to occur several days after inoculation of the M ϕ with mycobacteria (13). Together, these data suggest that the reduced apoptotic response of AM ϕ to infection by the virulent MTB strain H37Rv was due to the coinduction of a TNF- α -neutralizing activity.

Inactivation of TNF- α in AM ϕ infected with virulent MTB correlates with release of sTNFR2

To further dissect the events leading to attenuation of host M ϕ apoptosis, we tested whether release of sTNFR occurred following MTB infection as has been described in other systems (14–17). Although TNFR1 has been implicated as dominant in TNF- α -induced apoptosis (18), levels of sTNFR1 measured by ELISA were not significantly increased following MTB infection, and the levels did not differ with infection by the attenuated or virulent strain (at 48 h: $n = 7$, $p = 0.75$, Fig. 2). On the other hand, at 24 h the increase in sTNFR2 concentration in supernatants of H37Rv-infected AM ϕ was six times higher than in culture supernatants of AM ϕ infected with H37Ra (Fig. 3B). The difference in the TNFR2 levels was statistically significant at 6, 8, 24, 32, 48, and 96 h after infection ($n = 3$, $p < 0.05$ for all time points).

sTNFR2 inactivates TNF- α

To establish a causal relationship between levels of sTNFR2 and TNF- α bioactivity, we added recombinant sTNFR2 to cultures of AM ϕ infected with the attenuated MTB strain H37Ra and measured the bioactivity of TNF- α . Bioactivity measured by the L929 cell cytotoxicity assay was reduced by addition of exogenous sTNFR2 in a dose-dependent manner, and the reduction of cytotoxicity was statistically significant in cultures treated with 10 and 50 ng/ml of rTNFR2 compared with untreated controls ($n = 3$, $p < 0.05$; a representative experiment is shown in Fig. 4A). Apoptosis of host AM ϕ measured in the same cultures was also inhibited by sTNFR2 (Fig. 4B), and the amount of apoptosis was comparable to the level of TNF- α bioactivity. Addition of an irrelevant protein (bovine albumin) had no effect on either bioactivity.

The presence of TNF- α bound to sTNFR2 in supernatants of MTB-infected AM ϕ cultures was examined by a mixed ELISA assay. The TNF- α -sTNFR2 complexes were captured by immobilized Abs to sTNFR2 and quantified using Abs to TNF- α . Detectable levels of TNF- α -sTNFR2 complexes were found in cultures of AM ϕ infected with both MTB strains and were not found in




FIGURE 3. Levels of soluble TNFR2 in supernatants from AM ϕ inoculated with H37Rv, H37Ra, or uninfected AM ϕ and harvested at the indicated times. The experiment shown in A displays various early time points. The experiment shown in B covers a wider time range including a 96-h time point. For A and B, shown are mean \pm SD of triplicate values from one representative experiment of three. The differences in TNFR2 release by AM ϕ infected with H37Ra and H37Rv were statistically significant at 8, 32, and 96 h (A) and at 6, 24, and 48 h ($p < 0.05$).

uninfected AM ϕ . The levels of complexes in supernatants of AM ϕ infected with virulent H37Rv were about three times higher than in M ϕ infected with the attenuated strain H37Ra (Fig. 5).

IL-10 down-regulates AM ϕ apoptosis

Previous studies indicate that IL-10 has an anti-apoptotic function (19–21) and induces the release of sTNFR2 from several cell types (22, 23). To identify mechanisms regulating TNFR shedding from MTB-infected AM ϕ , we tested the effects of exogenous IL-10 and of neutralization of endogenous IL-10. Addition of anti-IL-10 Ab (10 μ g/ml) to AM ϕ cultures infected with the virulent MTB strain H37Rv significantly reduced the release of sTNFR2 ($n = 3$, $p < 0.05$; Fig. 6A) and increased M ϕ apoptosis ($n = 6$, $p < 0.05$, Fig. 6B). Conversely, addition of exogenous rIL-10 to AM ϕ cultures infected with the attenuated H37Ra decreased apoptosis of the host M ϕ in a dose-dependent manner (Fig. 6C). Reduction of apoptosis was statistically significant after addition of 5, 10, and 20 ng/ml IL-10 ($n = 6$, $p < 0.05$) to the infected cultures. Treatment of uninfected AM ϕ with IL-10 (10 ng/ml) was also found to increase sTNFR2 shedding from 56 ± 8 pg/ml to 160 ± 26 pg/ml at 6 h, and from 121 ± 38 pg/ml to 248 ± 45 pg/ml at 48 h of culture, and this difference was statistically significant ($n = 3$, $p < 0.05$). These findings indicate that the modulation of MTB-induced M ϕ apoptosis by sTNFR2 release may be regulated by IL-10 and suggest that the virulent strain H37Rv and the attenuated strain H37Ra differ in the induction of endogenous IL-10 secretion.

To demonstrate a role of IL-10 in modulation and induction of AM ϕ apoptosis by virulent and attenuated MTB strains, we measured IL-10 levels by ELISA in supernatants of AM ϕ cultures




FIGURE 4. Addition of srTNFR2 to AM ϕ cultures inoculated with attenuated MTB decreases cytotoxic activity of the supernatants (A) and apoptosis of host AM ϕ (B) in a dose-dependent fashion. One, 10, and 50 μ g/ml of srTNFR2 were added to AM ϕ cultures at the time of infection with H37Ra. Results from triplicate values of a typical experiment are shown. A, Cytotoxicity expressed as ng/ml TNF- α was measured at 24 h by assaying supernatants with L929 target cells. The reduction of cytotoxicity was statistically significant in cultures treated with 10 and 50 ng/ml rTNFR2 ($p < 0.05$). B, Apoptosis of host AM ϕ was measured at day 4 by the TUNEL assay. The reduction of % apoptosis was statistically significant in cultures treated with 1, 10, and 50 ng/ml rTNFR2 ($p < 0.05$).

inoculated with H37Rv and H37Ra. Uninfected M ϕ from all donors produced negligible levels of IL-10 (3 ± 1 pg/ml, $n = 12$). In contrast, H37Ra induced 58 ± 11 pg/ml, and H37Rv 103 ± 18 pg/ml IL-10 ($n = 12$, $p = 0.0092$, Fig. 7). Thus, the decreased induction of AM ϕ apoptosis by the virulent MTB strain H37Rv correlates directly with increased induction of IL-10.

Discussion

Infection with MTB is acquired via the lung. AM ϕ are the first line of defense against this infection, but mycobacteria can persist and




FIGURE 5. Concentration of TNF- α bound to sTNFR2 (representative for TNF- α -sTNFR2 complexes) in cultures of uninfected AM ϕ (CTRL) and AM ϕ infected with H37Ra or H37Rv. TNF- α -sTNFR2 complexes were assayed by mixed Ab ELISA (see *Materials and Methods*) of supernatants harvested 24 h after infection. The differences in the levels of TNF- α bound to TNFR2 in H37Ra- and H37Rv-infected cultures were statistically significant in all three experiments ($p < 0.05$ for all experiments).




FIGURE 6. Modulation of TNFR2 release (A) and apoptosis of host AM ϕ (B, C) in cell cultures infected with the virulent strain H37Rv after addition of anti-IL-10 Ab (10 μ g/ml) (A, B) and in cultures infected with attenuated H37Ra after addition of varying amounts of rIL-10 (C). TNFR2 was assayed at 24 h and AM ϕ apoptosis was assessed at 4 days. Shown are mean \pm SD of triplicate values from one representative experiment. The reduction of TNFR2 release (A) and the increase of % apoptosis (B) after addition of anti-IL-10 Ab is statistically significant ($p < 0.05$). Reduction of % apoptosis after addition of 5, 10, and 20 ng/ml IL-10 is also statistically significant ($p < 0.05$).

replicate within infected M ϕ . M ϕ respond to MTB by producing TNF- α , a pleiotropic cytokine critical for the protective immune response to tuberculosis (3), which elicits a wide spectrum of cellular responses including fever, tissue injury, and M ϕ activation (24). Recent studies showed that TNF- α induces also apoptosis (25). Host cell apoptosis is a defense strategy to limit the growth of certain intracellular pathogens, including viruses (26, 27). In the case of mycobacterial infection, this response has been shown to




FIGURE 7. IL-10 levels in supernatants of AM ϕ from 12 healthy individuals infected with H37Ra or H37Rv. IL-10 was measured by ELISA in supernatants harvested after 24 h. Values for cultures from individual donors are connected with a line. The mean of IL-10 concentration for all donors in supernatants from uninfected AM ϕ was 3 ± 1 pg/ml IL-10; this value is shown by the dashed line.

sequester bacilli within apoptotic cells and to restrict their replication directly (6), and by activation of uninfected M ϕ (7).

To investigate the possibility that virulent MTB might suppress host M ϕ apoptosis, we performed in vitro infection of normal human AM ϕ with a virulent and an attenuated strain of MTB. Our data show that inactivation of TNF- α and subsequent attenuation of apoptosis by the virulent strain H37Rv occurs by a unique mechanism. Compared with the attenuated strain H37Ra, H37Rv induces a quantitatively greater release of IL-10 from infected M ϕ . The enhanced IL-10 response causes increased accumulation of sTNFR2 in the supernatants, resulting in the inactivation of TNF- α by soluble receptor-ligand complex formation. sTNFR2 was found to inactivate TNF- α , since we showed that TNF- α -dependent cytotoxicity and apoptosis are significantly reduced by addition of rTNFR2 to H37Ra-infected AM ϕ cultures (Fig. 4). We have also attempted to demonstrate neutralization of endogenous sTNFR2 with anti-TNFR2 Abs and subsequent enhancement of apoptosis. In these experiments 10 μ g/ml anti-TNFR2 Abs were added to AM ϕ cultures at the time of inoculation with H37Rv. H37Rv-infected AM ϕ cultures to which no anti-TNFR2 Abs were added showed $39 \pm 5\%$ apoptosis at day 4 of culture in comparison with uninfected cultures ($9 \pm 1\%$ apoptosis, $p < 0.05$). In contrast, in the presence of anti-TNFR2 Abs, $49 \pm 3\%$ apoptosis ($n = 3$, $p < 0.05$) was observed. These experiments suggest that endogenous sTNFR2 is able to neutralize TNF- α , but do not rule out a stimulatory effect of these Abs on cell-associated TNFR2 and are therefore difficult to interpret. Furthermore, another way to prove involvement of sTNFR2 in TNF- α neutralization, specific blocking of TNFR2 release by metalloproteinase inhibitors, cannot be performed, because such inhibitors also affect processing of TNF- α (28).

The release of TNFR2 from the AM ϕ , which differs significantly in H37Rv- and H37Ra-infected AM ϕ , comprises an apparent virulence mechanism operative in pathogenic MTB.

This study indicates that a virulent MTB strain can modulate TNF- α activity by IL-10-dependent TNFR2 shedding. IL-10 has also been reported to prevent apoptosis of M ϕ during *Salmonella* infection (19), to prevent apoptosis of T cells in infectious mononucleosis (21), and to prevent apoptosis of germinal center B cells (20). In addition to attenuating apoptosis by increasing the release of sTNFR2, IL-10 might also down-regulate apoptosis by inhibiting IFN- γ and TNF- α production (29), and by down-regulating surface TNFR expression (23).

sTNFR2 release from the AM ϕ caused by IL-10 has been found to depend on the action of metalloproteinases (28). IL-10 has been found to down-regulate (30), to activate (31), and to have no effect (32) on the activity of metalloproteinases. In future studies we will examine the role of proteolytic enzymes in the IL-10-dependent release of TNFR2 from AM ϕ .

The cytokine responses of primary human AM ϕ to a MTB challenge typically demonstrate a high degree of individual variation, and a high IL-10 response might represent a host susceptibility factor to tuberculosis. On the other hand, the two individuals (of 12 individuals) whose AM ϕ did not respond to H37Rv inoculation with increased IL-10 levels (Fig. 7) might be part of a population with innate resistance to MTB. We are currently investigating this possibility.

AM ϕ apoptosis is induced by the action of TNF- α . TNF- α elicits cellular responses through its interaction with two distinct receptors, the 55-kDa TNFR1 and the 75-kDa TNFR2 (33). A distinctive difference between these two TNFR, which are both members of the TNF receptor superfamily including CD30, CD40, CD27, and Fas (34), is the presence of a death domain (DED) in TNFR1. Stimulation of TNFR1 results in activation of the apopto-

tic protease cascade (35, 36) leading to apoptosis. Inactivation of TNF- α by formation of a complex with sTNFR2 abrogates the induction of apoptosis upstream of the death-signaling cascade by elimination of the inducing cytokine signal. Related strategies are used by human herpes virus 4, which encodes a viral IL-10 homologue (37), parainfluenza virus, which induces IL-10 (38), and pox virus, which encodes a sTNFR1-like molecule (39).

sTNFR are naturally occurring inhibitors of TNF- α , which are normally present in the blood and urine (14–17). Serum concentrations of sTNFR increase significantly in a variety of disease states including tuberculosis and HIV-1 infection (40, 41). sTNFR2 were reported to be more prone to being released from AM ϕ than TNFR1 (42, 43), which is in agreement with our finding that MTB infection induced shedding of TNFR2, but not TNFR1, from AM ϕ . sTNFR2 are also more abundant than TNFR1 in the serum of patients with cancer and in the serum and synovial fluid of patients with rheumatoid arthritis (44, 45). This evidence suggests that sTNFR2 may be an important regulator of TNF- α activity in vivo. Our present investigations indicate that a virulent MTB strain can modulate TNF- α activity by IL-10-dependent shedding of sTNFR2.

The mycobacterial determinants that effect the unique capacity of virulent MTB to manipulate host TNF- α responses have yet to be defined. Identifying the basis of this multistep mechanism should provide further insight to the pathogenesis of human tuberculosis and may suggest novel approaches to therapy.

Note. A study describing reduction of TNF- α biological activity in supernatants from murine M ϕ infected with virulent *M. avium* strains with a possible role for sTNFR2 has recently come to our attention (46).

Acknowledgments

We thank Dr. Henry Koziel, Beth Israel-Deaconess Medical Center, Boston, MA, for supplying bronchoalveolar lavage fluid.

References

- Dolin, P. J., M. C. Ravigliano, and A. Kochi. 1994. Global tuberculosis incidence and mortality during 1990–2000. *Bull. W. H. O.* 72:213.
- Moore, M., I. M. Onorato, E. McCray, and K. G. Castro. 1997. Trends in drug-resistant tuberculosis in the United States, 1993–1996. *JAMA* 278:833.
- Flynn, J. L., M. M. Goldstein, J. Chan, K. J. Triebold, K. Pfeffer, C. J. Lowenstein, R. Schreiber, T. W. Mak, and B. R. Bloom. 1995. Tumor necrosis factor- α is required in the protective immune response against *Mycobacterium tuberculosis* in mice. *Immunity* 2:561.
- Keane, J., M. K. Balcewicz-Sablinska, H. G. Remold, G. L. Chupp, B. B. Meek, M. J. Fenton, and H. Kornfeld. 1997. Infection by *Mycobacterium tuberculosis* promotes human alveolar macrophage apoptosis. *Infect. Immun.* 65:298.
- Rojas, M., L. F. Barrera, G. Puzo, and L. F. Garcia. 1997. Differential induction of apoptosis by virulent *Mycobacterium tuberculosis* in resistant and susceptible murine macrophages: role of nitric oxide and mycobacterial products. *J. Immunol.* 159:1352.
- Molloy, A., P. Laochumroonvorapong, and G. Kaplan. 1994. Apoptosis, but not necrosis, of infected monocytes is coupled with killing of intracellular bacillus Calmette-Guérin. *J. Exp. Med.* 180:1499.
- Fratazzi, C., R. D. Arbeit, C. Carini, and H. G. Remold. 1997. Programmed cell death of *Mycobacterium avium* serovar 4-infected human macrophages prevents the mycobacteria from spreading and induces mycobacterial growth inhibition by freshly added, uninfected macrophages. *J. Immunol.* 158:4320.
- Fadok, V. A., J. S. Savill, C. Haslett, D. L. Bratton, D. E. Doherty, P. A. Campbell, and P. M. Henson. 1992. Different populations of macrophages use either the vitronectin receptor or the phosphatidylserine receptor to recognize and remove apoptotic cells. *J. Immunol.* 149:4029.
- Inderlied, C. B., L. S. Young, and J. K. Yamada. 1987. Determination of in vitro susceptibility of *Mycobacterium avium* complex isolates to antimycobacterial agents by various methods. *Antimicrob. Agents Chemother.* 31:1697.
- Flick, D. A., and G. E. Gifford. 1984. Comparison of in vitro cell cytotoxic assays for tumor necrosis factor. *J. Immunol. Methods* 68:167.
- Corti, A., C. Poesi, S. Merli, and G. Cassani. 1994. Tumor necrosis factor (TNF) α quantification by ELISA and bioassay: effects of TNF α -soluble TNF receptor (p55) complex dissociation during assay incubations. *J. Immunol. Methods* 177:191.
- Gavrieli, Y., Y. Sherman, and S. A. Ben-Sasson. 1992. Identification of programmed cell death in situ via specific labeling of nuclear DNA fragmentation. *J. Cell. Biol.* 119:493.

13. Gan, H., G. W. Newman, and H. G. Remold. 1995. Plasminogen activator inhibitor type 2 prevents programmed cell death of human macrophages infected with *Mycobacterium avium*, serovar 4. *J. Immunol.* 155:1304.
14. Novick, D., H. Engelmann, D. Wallach, and M. Rubinstein. 1989. Soluble cytokine receptors are present in normal human urine. *J. Exp. Med.* 170:1409.
15. Engelmann, H., D. Aderka, M. Rubinstein, D. Rotman, and D. Wallach. 1989. A tumor necrosis factor-binding protein purified to homogeneity from human urine protects cells from tumor necrosis factor toxicity. *J. Biol. Chem.* 264:11974.
16. Engelmann, H., D. Novick, and D. Wallach. 1990. Two tumor necrosis factor-binding proteins purified from human urine: evidence for immunological cross-reactivity with cell surface tumor necrosis factor receptors. *J. Biol. Chem.* 265:1531.
17. Neuner, P., G. Klosner, M. Pourmojib, F. Trautinger, and R. Knobler. 1994. Selective release of tumor necrosis factor binding protein II by malignant human epidermal cells reveals protection from tumor necrosis factor alpha-mediated cytotoxicity. *Cancer Res.* 54:6001.
18. Tartaglia, L. A., M. Rothe, Y. F. Hu, and D. V. Goeddel. 1993. Tumor necrosis factor's cytotoxic activity is signaled by the p55 TNF receptor. *Cell* 73:213.
19. Arai, T., K. Hiromatsu, H. Nishimura, Y. Kimura, N. Kobayashi, H. Ishida, Y. Nimura, and Y. Yoshikai. 1995. Endogenous interleukin 10 prevents apoptosis in macrophages during *Salmonella* infection. *Biochem. Biophys. Res. Commun.* 213:600.
20. Levy, Y., and J. C. Broutet. 1994. Interleukin-10 prevents spontaneous death of germinal center B cells by induction of the bcl-2 protein. *J. Clin. Invest.* 93:424.
21. Taga, K., J. Chretien, B. Cherney, L. Diaz, M. Brown, and G. Tosato. 1994. Interleukin-10 inhibits apoptotic cell death in infectious mononucleosis T cells. *J. Clin. Invest.* 94:251.
22. Seitz, M., P. Loetscher, B. Dewald, H. Towbin, H. Gallati, and M. Baggiolini. 1995. Interleukin-10 differentially regulates cytokine inhibitor and chemokine release from blood mononuclear cells and fibroblasts. *Eur. J. Immunol.* 25:1129.
23. Joyce, D. A., D. P. Gibbons, P. Green, J. H. Steer, M. Feldmann, and F. M. Brennan. 1994. Two inhibitors of pro-inflammatory cytokine release, interleukin-10 and interleukin-4, have contrasting effects on release of soluble p75 tumor necrosis factor receptor by cultured monocytes. *Eur. J. Immunol.* 24:2699.
24. Beutler, B., and A. Cerami. 1986. Cachectin and tumor necrosis factor as two sides of the same biological coin. *Nature* 320:584.
25. Laster, S. M., J. G. Wood, and L. R. Gooding. 1988. Tumor necrosis factor can induce both apoptotic and necrotic forms of cell lysis. *J. Immunol.* 141:2629.
26. Antoni, B. A., P. Sabbatini, A. B. Rabson, and E. White. 1995. Inhibition of apoptosis in human immunodeficiency virus-infected cells enhances virus production and facilitates persistent infection. *J. Virol.* 69:2384.
27. Vaux, D. L., G. Haeccker, and A. Strasser. 1994. An evolutionary perspective on apoptosis. *Cell* 76:777.
28. Crowe, P. D., B. N. Walter, K. M. Mohler, C. Otten-Evans, R. A. Black, and C. F. Ware. 1995. A metalloprotease inhibitor blocks shedding of the 80-kD TNF receptor and TNF processing in T lymphocytes. *J. Exp. Med.* 181:1205.
29. de Waal Malefyt, R., J. Abrams, B. Bennett, C. G. Figdor, and J. E. de Vries. 1991. Interleukin 10 (IL-10) inhibits cytokine synthesis by human monocytes: an autoregulatory role of IL-10 produced by monocytes. *J. Exp. Med.* 174:1209.
30. Lacraz, S., L. P. Nicod, R. Chicheportiche, H. G. Welgus, and J. M. Dayer. 1995. IL-10 inhibits metalloproteinase and stimulates TIMP-1 production in human mononuclear phagocytes. *J. Clin. Invest.* 96:2304.
31. Reitamo, S., A. Remitz, K. Tamai, and J. Uitto. 1994. Interleukin-10 modulates type I collagen and matrix metalloprotease gene expression in cultured human skin fibroblasts. *J. Clin. Invest.* 94:2489.
32. Kumar, R., Z. Dong, and I. J. Fidler. 1996. Differential regulation of metalloelastase activity in murine peritoneal macrophages by granulocyte-macrophage colony-stimulating factor and macrophage colony-stimulating factor. *J. Immunol.* 157:5104.
33. Bazzoni, F., and B. Beutler. 1996. The tumor necrosis factor ligand and receptor families. *N. Engl. J. Med.* 334:1717.
34. Smith, C. A., T. Farrah, and R. G. Goodwin. 1994. The TNF receptor superfamily of cellular and viral proteins: activation, costimulation, and death. *Cell* 76:959.
35. Muzio, M., A. M. Chinnaiyan, F. C. Kischkel, K. O'Rourke, A. Shevchenko, J. Ni, C. Scaffidi, J. D. Bretz, M. Zhang, R. Gentz, M. Mann, P. H. Kramer, M. E. Peter, and V. M. Dixit. 1996. FLICE, a novel FADD-homologous ICE/CED-3-like protease, is recruited to the CD95 (Fas/APO-1) death-inducing signaling complex. *Cell* 85:817.
36. Boldin, M. P., T. M. Goncharov, Y. V. Goltsev, and D. Wallach. 1996. Involvement of MACH, a novel MORT1/FADD-interacting protease, in Fas/APO-1- and TNF receptor-induced cell death. *Cell* 85:803.
37. Suzuki, T., H. Tahara, S. Narula, K. W. Moore, P. D. Robbins, and M. T. Lotze. 1995. Viral interleukin 10 (IL-10), the human herpes virus 4 cellular IL-10 homologue, induces local anergy to allogeneic and syngeneic tumors. *J. Exp. Med.* 182:477.
38. Sieg, S., C. King, Y. Huang, and D. Kaplan. 1996. The role of interleukin-10 in the inhibition of T-cell proliferation and apoptosis mediated by parainfluenza virus type 3. *J. Virol.* 70:4845.
39. Schreiber, M., L. Sedger, and G. McFadden. 1997. Distinct domains of M-T2, the myxoma virus tumor necrosis factor (TNF) receptor homolog, mediate extracellular TNF binding and intracellular apoptosis inhibition. *J. Virol.* 71:2171.
40. Kawaguchi, H., Y. Ina, S. Ito, S. Sato, Y. Sugiura, H. Tomita, N. Ogisu, K. Takada, M. Yamamoto, M. Morishita, and K. Yoshikawa. 1996. Serum levels of soluble tumor necrosis factor (TNF) receptors in patients with pulmonary tuberculosis. *Kekkaku* 71:259.
41. Zangerle, R., H. Gallati, M. Sarcelletti, G. Weiss, H. Denz, H. Wachter, and D. Fuchs. 1994. Increased serum concentrations of soluble tumor necrosis factor receptors in HIV-infected individuals are associated with immune activation. *J. Acquir. Immune Defic. Syndr.* 7:79.
42. Galve-de Rochemonteix, B., L. P. Nicod, and J. M. Dayer. 1996. Tumor necrosis factor soluble receptor 75: the principal receptor form released by human alveolar macrophages and monocytes in the presence of interferon gamma. *Am. J. Respir. Cell Mol. Biol.* 14:279.
43. Joyce, D. A., and J. H. Steer. 1996. IL-4, IL-10 and IFN-gamma have distinct, but interacting, effects on differentiation-induced changes in TNF-alpha and TNF receptor release by cultured human monocytes. *Cytokine* 8:49.
44. Aderka, D., H. Englemann, V. Hornik, Y. Skornick, Y. Levo, D. Wallach, and G. Kushtai. 1991. Increased serum levels of soluble receptors for tumor necrosis factor in cancer patients. *Cancer Res.* 51:5602.
45. Cope, A. P., D. Aderka, M. Doherty, H. Engelmann, D. Gibbons, A. C. Jones, F. M. Brennan, R. N. Maini, D. Wallach, and M. Feldmann. 1992. Increased levels of soluble tumor necrosis factor receptors in the sera and synovial fluid of patients with rheumatic diseases. *Arthritis Rheum.* 35:1160.
46. Eviks, I. S., and C. L. Emerson. 1997. Temporal effect of tumor necrosis factor-alpha on murine macrophages infected with *Mycobacterium avium*. *Infect. Immun.* 65:2100.