

Investigate small particles with unparalleled sensitivity  
**Amnis® CellStream®** Flow Cytometry System

For Research Use Only. Not for use in diagnostic procedures.



**Luminex®**  
complexity simplified.



## Cutting Edge: SIV Nef Protein Utilizes Both Leucine- and Tyrosine-Based Protein Sorting Pathways for Down-Regulation of CD4

This information is current as of October 26, 2021.

Patricia A. Bresnahan, Wes Yonemoto and Warner C. Greene

*J Immunol* 1999; 163:2977-2981; ;  
<http://www.jimmunol.org/content/163/6/2977>

**References** This article **cites 39 articles**, 18 of which you can access for free at:  
<http://www.jimmunol.org/content/163/6/2977.full#ref-list-1>

Why *The JI*? [Submit online.](#)

- **Rapid Reviews! 30 days\*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

*\*average*

**Subscription** Information about subscribing to *The Journal of Immunology* is online at:  
<http://jimmunol.org/subscription>

**Permissions** Submit copyright permission requests at:  
<http://www.aai.org/About/Publications/JI/copyright.html>

**Email Alerts** Receive free email-alerts when new articles cite this article. Sign up at:  
<http://jimmunol.org/alerts>

*The Journal of Immunology* is published twice each month by  
The American Association of Immunologists, Inc.,  
1451 Rockville Pike, Suite 650, Rockville, MD 20852  
Copyright © 1999 by The American Association of  
Immunologists All rights reserved.  
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



## Cutting Edge: SIV Nef Protein Utilizes Both Leucine- and Tyrosine-Based Protein Sorting Pathways for Down-Regulation of CD4<sup>1</sup>

Patricia A. Bresnahan,<sup>2\*‡</sup> Wes Yonemoto,<sup>2\*</sup> and Warner C. Greene<sup>3\*†‡</sup>

**The Nef protein is unique to primate lentiviruses and is closely linked to accelerated pathogenesis in both human and monkey hosts. Nef acts to down-regulate CD4 and MHC class I, two receptors important for immune function. A recent report demonstrated the presence of two tyrosine motifs in SIV Nef that contribute to its ability to down-regulate CD4 and to associate with clathrin adaptors. These tyrosine motifs are not present in HIV-1 Nef, which instead utilizes a leucine-based motif for its down-regulation of CD4. We now report that SIV Nef also contains a conserved leucine-based motif that contributes to CD4 down-regulation, functions to stimulate internalization, and contributes to the association of SIV Nef with clathrin adaptors AP-1 and AP-2. These results demonstrate that SIV Nef differs from HIV-1 Nef by its ability to use two parallel pathways of the protein-sorting machinery based on either tyrosine or leucine motifs. *The Journal of Immunology*, 1999, 163: 2977–2981.**

**B**ecause of the genetic and biological similarities between HIV-1 and SIV, SIV infection in macaques has emerged as a valuable animal model for HIV-1 infection in humans. Infection of macaques with live attenuated forms of SIV provides an important approach to assessing the safety and efficacy of a human HIV-1 vaccine (1, 2). Given the ability of both viruses to rapidly evolve, the identification of differences in the structure and function of viral gene products may not only determine the adequacy of the animal model, but also provide insights into functional motifs that are important for pathogenesis.

\*Gladstone Institute of Virology and Immunology, and Departments of <sup>†</sup>Medicine and <sup>‡</sup>Microbiology and Immunology, University of California, San Francisco, CA 94141

Received for publication April 26, 1999. Accepted for publication July 13, 1999.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup> This work was supported by National Institutes of Health Grants R01-AI28240-10 (to W.C.G.) and P30 MH59037 (University of California San Francisco-Gladstone Institute of Virology and Immunology Center for AIDS Research). P.A.B. is supported by Universitywide AIDS Research Program (UARP) Grant F98-GI-147.


<sup>2</sup> P.A.B. and W.Y. contributed equally to this work.

<sup>3</sup> Address correspondence and reprint requests to Dr. Warner C. Greene, Gladstone Institute of Virology and Immunology, P.O. Box 419100, San Francisco, CA 94141. E-mail address: wgreene@gladstone.ucsf.edu

The Nef proteins of HIV-1 and SIV are 27- to 32-kDa myristoylated proteins that are found in both the membranes and cytosol of infected host cells (reviewed in Ref. 3). While not strictly required for viral infection and replication, attenuation of Nef expression in either HIV-1 or SIV is associated with markedly delayed progression of disease (1, 4, 5). The absence of Nef also decreases the intrinsic infectivity of SIV and HIV-1 in cultured cells, a characteristic that likely underlies the diminished viral loads observed with Nef-deleted viruses in vivo (6–9). Nef also mediates the down-regulation of two key immune modulators, CD4 (10–14) and MHC class I (15–17) in host cells. However, the relationship of these functions of Nef to its role in viral pathogenesis remains poorly understood.

Several observations show that distinct subregions of Nef are required for down-regulation of CD4 and MHC class I (18). First, CD4 down-regulation requires the presence of a dileucine sequence in Nef, which acts as an internalization signal, (19–21) and two discontinuous regions, including the sequence W<sup>57</sup>L<sup>58</sup>, that may form a binding site for the CD4 receptor tail (22, 23). In addition, a diacidic motif at positions 155,156 in HIV-1 Nef was recently identified as a lysosomal sorting signal involved in CD4 down-regulation (24). In contrast, MHC class I down-regulation is not affected by mutation of the dileucine sequence, but instead requires a proline-rich domain (not required for CD4 down-regulation), an upstream cluster of acidic amino acids, and an N-terminal  $\alpha$  helical domain (16, 18, 25). Despite these genetically separable activities of Nef, its presence similarly leads to diminished CD4 and MHC class I expression involving increased endocytosis, blocked progression through the secretory pathway at the *trans* Golgi membrane, and enhanced sorting to the lysosome for degradation (11, 13, 17, 25).

Both HIV-1 and SIV Nef have recently been shown to form a complex with the clathrin adaptors AP-1 and AP-2 (19–21), which recruit clathrin to the Golgi membrane and the plasma membrane, respectively (26, 27). Both adaptors bind specific peptide motifs, including tyrosine-based sequences (Y-X-X- $\phi$ , where X may be any amino acid, and  $\phi$  is an amino acid with a hydrophobic side chain) and leucine-based sequences (L-X, where X may be L, I, M, V, or F) on the cytoplasmic portion of transmembrane proteins. Thus, these adaptors link transmembrane receptors with tyrosine- or leucine-based sorting signals to the clathrin-based vesicles involved in protein trafficking and compartmentalization. However, the two sorting signals are not entirely interchangeable. Tyrosine- and leucine-based motifs can be independently saturated (28), and



**FIGURE 1.** A, Structural comparison of HIV-1 Nef and SIV Nef, including the relative positions of two tyrosine-based internalization motifs at positions 28 and 39 in SIV Nef and a conserved leucine motif, at position 164,165 in HIV-1 Nef and 194,195 in SIV Nef. B, Relative expression of human CD4 on the surface of transfected 293 cells in the presence of wild-type and mutant forms of SIV Nef (lanes 3, 4, 5, and 6), HIV Nef (lane 7), or in the absence of Nef (lane 2). CD4 down-regulation was measured by FACS analysis, as previously described (2). Values are reported as the mean of three samples, with error bars representing the standard deviation. C, Expression of SIV Nef wild-type (lane 2), SIV Nef Y<sup>28G</sup>Y<sup>39G</sup> (lane 3), SIV Nef L<sup>194A</sup>M<sup>195A</sup> (lane 4), or SIV Nef Y<sup>28G</sup>Y<sup>39G</sup>/L<sup>194A</sup>M<sup>195A</sup> (lane 5) in transiently transfected human 293 cells. The SIV Nef proteins were detected by immunoblotting using an anti-SIV Nef rabbit antiserum (a gift from Dr. Paul Luciw). Lane 1 depicts the results obtained by immunoblotting lysate from mock-transfected cells.

the two motifs mediate different sorting reactions at the *trans* Golgi (29). The tyrosine- and leucine-based motifs thus appear to represent parallel but distinct sorting mechanisms within the cell.

As noted, different adaptor motifs are apparently employed by SIV Nef and HIV-1 Nef, raising the issue of whether these two viral proteins have evolved to use distinct and not entirely redundant protein sorting pathways. SIV utilizes two tyrosine-based motifs located near the N terminus (30) (Fig. 1A). In contrast, HIV-1 Nef lacks these tyrosine-based motifs and instead utilizes a highly conserved leucine-based motif located near the C terminus (Fig. 1A). Mutation of this motif leads to a loss of CD4 down-regulation and clathrin adaptor binding (19–21). We now describe the presence of a previously unrecognized leucine-based sorting motif in SIV Nef that is highly conserved in SIV Nef alleles. This leucine-based motif functions in a largely redundant manner with the previously reported tyrosine-based motifs to mediate CD4 down-regulation and assembly with clathrin adaptors. The tyrosine- and leucine-based motifs in SIV Nef contribute additively to the rate of internalization of a CD8-SIV Nef chimera, and mutation of both motifs is required to significantly diminish the ability SIV Nef to down-regulate CD4. These findings highlight the presence of a shared leucine-based trafficking motif in both SIV and HIV-1 Nef and the unique acquisition of additional tyrosine-based motifs in SIV Nef.

## Materials and Methods

### Abs and immunodetection

Rabbit serum to SIV Nef was provided by Dr. P. A. Luciw (University of California Davis, Davis, CA). OKT8 mAb specific for the human CD8 Ag was obtained from Dr. A. Weiss (University of California, San Francisco, CA). The following Abs were used for FACS analysis of transiently expressed cell-surface receptors: anti-CD4-Tricolor, anti-CD8-Tricolor (Caltag, South San Francisco, CA), anti-CD8-PE, and anti-CD25-FITC (Becton Dickinson, Mountain View, CA). Control cells used for determining background staining in the FACS analyses were transfected with empty expression vector only. Abs specific for AP-1 (clone 100/3, anti-adaptin  $\gamma$ ; Sigma, St. Louis, MO) or AP-2 (anti-adaptin  $\alpha$ ; Transduction Laboratories, Lexington, KY) were used for immunoblotting, with a HRP-conjugated goat-anti-mouse secondary Ab (Zymax; Zymed, South San Francisco, CA). Immunoprecipitations were performed as described previously (19).

### DNA constructions


The plasmids pSIV Nef and pSIV Nef Y<sup>28G</sup>Y<sup>39G</sup>, which express wild-type SIV Nef and a mutant of SIV Nef in which the tyrosine residues at positions 28 and 39 are replaced with glycines, were prepared by subcloning the SIV<sub>MAC239</sub> nef sequences from CMX-SIV<sub>MAC239</sub>Nef and CMX-SIV<sub>MAC239</sub> Nef Y<sup>28G</sup>Y<sup>39G</sup> (30) into the vector pcDNA3.1 (Invitrogen, San Diego, CA). To generate SIV<sub>MAC239</sub> Nef mutants, in which the leucine and methionine at positions 194 and 195 are replaced with alanines, the plasmids pSIV Nef L<sup>194A</sup>M<sup>195A</sup> and pSIV Nef Y<sup>28G</sup>Y<sup>39G</sup>/L<sup>194A</sup>M<sup>195A</sup> were generated from the plasmids pSIV Nef and pSIV Nef Y<sup>28G</sup>Y<sup>39G</sup>, respectively, by site-directed mutagenesis (Bio-Rad, Richmond, CA).

Plasmids expressing CD8-SIV Nef chimeras were prepared by amplifying wild-type and mutant SIV<sub>MAC239</sub> nef genes by PCR and inserting them into the pCN vector (31) as previously described (19). The resulting plasmids, pCD8-SIV Nef, pCD8-SIV Nef Y<sup>28G</sup>Y<sup>39G</sup>, pCD8-SIV Nef L<sup>194A</sup>M<sup>195A</sup>, and pCD8-SIV Nef Y<sup>28G</sup>Y<sup>39G</sup>/L<sup>194A</sup>M<sup>195A</sup>, all express chimeras in which the extracellular and transmembrane domains are derived from CD8, and the cytoplasmic domain consists of the entire sequence of SIV<sub>MAC239</sub> Nef.

## Results and Discussion

Comparison of SIV and HIV-1 Nef sequences revealed a highly conserved leucine in SIV Nef at position 194 corresponding to L164 in HIV-1 Nef (NL4-3 strain) (32) (Fig. 1A). An adjacent methionine at position 195 formed a consensus sequence for a leucine-based sorting signal. To assess the potential functional contribution of this conserved leucine-based motif to the SIV Nef-mediated down-regulation of CD4, the site was mutated by substitution of the leucine and methionine with alanines (L<sup>194A</sup>M<sup>195A</sup>). Additionally, glycine substitutions were introduced for Y<sup>28</sup> and Y<sup>39</sup> (Y<sup>28G</sup>Y<sup>39G</sup>) as previously described (30), and a composite mutant (Y<sup>28G</sup>Y<sup>39G</sup>/L<sup>194A</sup>M<sup>195A</sup>) was also prepared. Surprisingly, both the wild-type and Y<sup>28G</sup>Y<sup>39G</sup> forms of SIV Nef reduced the cell surface expression of CD4 to approximately the same extent (Fig. 1B). Introduction of the L<sup>194A</sup>M<sup>195A</sup> mutations resulted in a partial loss of CD4 down-regulation; however, the composite mutant (SIV Nef Y<sup>28G</sup>Y<sup>39G</sup>/L<sup>194A</sup>M<sup>195A</sup>) exhibited essentially no CD4 down-regulation. The partial loss of CD4 down-regulation by SIV Nef L<sup>194A</sup>M<sup>195A</sup> and SIV Nef Y<sup>28G</sup>Y<sup>39G</sup>/L<sup>194A</sup>M<sup>195A</sup> mutants could not be explained by poor expression, as all of the SIV Nef analogues were expressed at nearly the same steady state level (Fig. 1C). We conclude that the leucine-based motif in SIV Nef plays an important role in CD4 down-regulation, and that the N-terminal tyrosine-based motifs also play a role best revealed in the context of the composite mutation.

To further characterize the functional properties of both the tyrosine- and leucine-based motifs in SIV Nef, we constructed chimeric proteins with a truncated form of human CD8 comprising the extracellular and transmembrane domains fused to wild-type SIV Nef, or mutant SIV Nef sequences as a cytoplasmic domain (19, 31, 33). Steady state surface expression of the CD8-SIV Nef wild-type chimera was ~5-fold lower than that of CD8 (Fig. 2A), and the CD8-SIV Nef Y<sup>28G</sup>Y<sup>39G</sup> and CD8-SIV Nef L<sup>194A</sup>M<sup>195A</sup>




**FIGURE 2.** Steady state cell-surface expression of CD8 and CD8-SIV Nef chimeric proteins in transfected human 293 cells. *A*, Cell-surface expression was quantitated by FACS; positive cells were scored as brighter than 95% of mock-transfected cells. Error bars indicate the SD from the mean value of three samples. *B*, Relative expression of CD8-SIV Nef chimeras detected by immunoblotting with anti-SIV Nef Ab, as described in *Materials and Methods*.

chimeras were similarly low. In contrast, the composite mutant CD8-SIV Nef Y<sup>28G</sup>Y<sup>39G</sup>/L<sup>194A</sup>M<sup>195A</sup> was expressed at much higher levels at the cell surface. While all of the chimeras were comparably expressed (Fig. 2*B*), the composite mutant (CD8-SIV Nef Y<sup>28G</sup>Y<sup>39G</sup>/L<sup>194A</sup>M<sup>195A</sup>) was consistently expressed at lower levels on the cell surface than wild-type CD8. These findings suggest that a single mutation interrupting either the tyrosine- or leucine-based motifs in SIV Nef does not release the chimeric molecule from down-regulation. However, mutation of both motifs significantly impairs down-regulation, arguing that both play a role in intracellular protein sorting.

Expression of the CD8-Nef chimera does not demonstrate the partially impaired down-regulation activity found for SIV Nef L<sup>194A</sup>M<sup>195A</sup> (Fig. 1*B*). Experiments using SIV Nef sequences in *cis* and in *trans* detect different aspects of its association with the intracellular protein sorting machinery. In the *trans* assay of CD4 down-regulation, at least three different properties of SIV Nef are required: 1) membrane localization; 2) coupling to CD4, possibly through a linker protein; and 3) association with clathrin adaptors and possibly other components of the protein sorting machinery. In the *cis* assay, the Nef sequences are not required for membrane localization, nor is assembly with the CD4 cytoplasmic tail needed. Thus, while the chimeric proteins allow a more direct measurement of the effects of SIV Nef sequences on internalization rates, their overall trafficking in the cell is less complex than that found with the soluble form of Nef. Accordingly, the *trans*- and *cis*- down-regulation assays are not fully equivalent, although both can detect the functional interplay of SIV Nef with the protein sorting machinery.

Endocytosis of both CD4 and MHC class I molecules is enhanced in the presence of Nef. To determine the relative contributions of the tyrosine- and leucine-based motifs in SIV Nef to the stimulation of endocytosis, we measured the kinetics of internalization of each CD8-SIV Nef chimera (Fig. 3). Internalization was reduced to approximately half the rate of CD8-SIV Nef wild-type with mutation of either the tyrosine- or leucine-based motif. Mutation of both motifs (CD8-SIV Nef Y<sup>28G</sup>Y<sup>39G</sup>/L<sup>194A</sup>M<sup>195A</sup>) further reduced the rate of internalization. These results demonstrate




**FIGURE 3.** Both tyrosine motifs and leucine motifs contribute to the rate of CD8-SIV Nef internalization. Internalization rate of CD8-SIV Nef chimera in the absence of the N-terminal tyrosine motifs (CD8-SIV Nef Y<sup>28G</sup>Y<sup>39G</sup>), the C-terminal leucine-based motif (CD8-SIV Nef L<sup>194A</sup>M<sup>195A</sup>), or in the absence of both motifs (CD8-SIV Nef Y<sup>28G</sup>Y<sup>39G</sup>/L<sup>194A</sup>M<sup>195A</sup>). Error bars indicate the SD from the mean value of three samples.

that either motif can function independently as an internalization signal and that they both contribute additively to the overall rate of internalization.

We next investigated the effect of tyrosine- or leucine-based motifs on the ability of the CD8-SIV Nef chimeras to associate with the AP-1 and AP-2 adaptors in vivo. Mutation of either the tyrosine- or the leucine-based motif clearly diminished the association of the corresponding Nef analogues with both AP-1 and AP-2 (Fig. 4). The presence of the composite mutations (CD8-SIV Nef Y<sup>28G</sup>Y<sup>39G</sup>/L<sup>194A</sup>M<sup>195A</sup>) further reduced association with both adaptors. Thus, both motifs in SIV Nef contribute independently to an interaction with AP-1 and AP-2. The residual presence of AP-1 (13% of maximum signal) compared with AP-2 (0.2% of maximum signal, compare Figs. 4*A* and 4*B*, lanes 7) in complex with the composite mutant could reflect the contribution of Nef-associated proteins like NBP1 (34) in the immunoprecipitate that can themselves recruit clathrin adaptors through their own tyrosine- or leucine-based motifs.

While we detected relatively equivalent amounts of AP-1 associated with both HIV and SIV-1 Nef proteins, much less AP-2 was associated with HIV-1 Nef sequences (4% of SIV Nef signal) (Fig. 4*B*, lanes 3 and 4). These results could simply reflect fewer sorting



**FIGURE 4.** Coimmunoprecipitation of clathrin adaptors AP-1 (*A*) and AP-2 (*B*) with SIV Nef chimeras. Human 293 cells were mock transfected or transfected (lane 2) with CD8-HIV-1 Nef (lane 3) or CD8-SIV Nefs (lanes 4, 5, 6, and 7). Cell lysates were immunoprecipitated with anti-CD8 (clone OKT-8, a gift from Dr. A. Weiss) (lanes 2–8) or with control IgG of the same isotype (lane 1). Purified bovine adaptors (a gift from Dr. F. Brodsky, University of California, San Francisco, CA) were electrophoresed and immunoblotted as a control (lane 8). Relative percent signal was determined by densitometry analysis and normalized to CD8-SIV Nef wild type in both panels.

motifs for AP-2 in HIV-1 Nef (a single leucine motif) than in SIV Nef (two tyrosine motifs and a leucine motif), though the same sites appear involved in the association of AP-1, which is comparable. Alternatively, SIV Nef may form a more stable complex with AP-2 under these immunoprecipitating conditions than HIV-1 Nef. Studies of AP-2 interactions with sorting motifs suggest that regulatory mechanisms may stabilize these interactions in vivo (27, 35, 36). Further studies on binding affinities of HIV-1 Nef and SIV Nef for each of the adaptors are required to explore this issue.

Together, our results indicate that SIV Nef has evolved to utilize two apparently parallel pathways for CD4 receptor internalization involving both tyrosine- and leucine-based motifs. In contrast, HIV-1 Nef contains a trafficking signal for only one of these pathways. The significance of the two independent, parallel pathways for internalization is not clear. Several cellular receptors have been described to contain multiple internalization signals using both the tyrosine- and leucine-based signals for internalization. Of note, once the receptor is internalized, these signals may participate in distinct sorting processes (37–41). Thus, it is possible that these motifs are not entirely redundant, but are utilized differently for sorting to specific end compartments within the cell.

Tyrosine-based motifs are highly conserved in pathogenic SIV strains, suggesting that genetic selection pressure is applied to the tyrosine-based motif in Nef in the context of SIV in macaques. However, the leucine-based motif is highly conserved both in SIV Nef and HIV-1 Nef. Therefore, leucine-based motifs may be fundamental components of Nef activity in both HIV-1 and SIV, while tyrosine motifs may contribute to a distinct part of SIV pathogenesis in macaques. Whether down-regulation of CD4 by Nef is clearly linked to accelerated HIV-1 or SIV pathogenesis remains to be determined. It has recently been observed that Nef is not absolutely required for pathogenesis (2), but its absence in SIV or HIV-1 significantly prolongs the onset of AIDS in adults (1, 4).

Infection of macaques with attenuated SIV is a leading model for testing the effectiveness and the safety of various HIV-1 vaccines. Therefore, it is important to establish the functional equivalence of the two viruses and the relationship of their viral products to pathogenesis. The maintenance of both tyrosine- and leucine-based sorting motifs in SIV Nef, but only a leucine-based motif in HIV-1 Nef, could be an indication of functional distinctions. These differences, if present, may emerge with further examination of the function of HIV-1 Nef within the context of SIV-HIV chimeric viruses (SHIVs).

## Acknowledgments

We thank Drs. A. Weiss and P. Luciw for Abs, Drs. F. Brodsky and A. Wilde for purified bovine adaptors, and Drs. D. Trono and C. Aiken for SIVMAC239 clones. We also thank R. Gelezianas, S. Ferrell, and W. Xu for helpful discussion; J. Carroll and N. Shea for graphics preparation; S. Gonzales and C. Goodfellow for photography; and G. Howard and S. Ordway for editorial assistance.

## References

- Kestler, H. W., D. J. Ringler, K. Mori, D. L. Panicali, P. K. Sehgal, M. D. Daniel, and R. C. Derosiers. 1991. Importance of the *nef* gene for maintenance of high virus loads and for development of AIDS. *Cell* 65:651.
- Baba, T. W., V. Liska, A. H. Khimani, N. B. Ray, P. J. Dailey, D. Penninck, R. Bronson, M. F. Greene, H. M. McClure, L. N. Martin, and R. M. Ruprecht. 1999. Live attenuated, multiply deleted simian immunodeficiency virus causes AIDS in infant and adult macaques. *Nat. Med.* 5:194.
- Oldridge, J., and M. Marsh. 1998. Nef-an adaptor adaptor? *Trends Cell Biol.* 8:302.
- Deacon, N. J., A. Tsykin, A. Solomon, K. Smith, M. Ludford-Menting, D. J. Hooker, D. A. McPhee, A. L. Greenway, A. Ellett, C. Chartfield, et al. 1995. Genomic structure of an attenuated quasi species of HIV-1 from a blood transfusion donor and recipients. *Science* 270:988.
- Kirchoff, T., T. C. Greenough, D. B. Brettler, J. L. Sullivan, and R. C. Derosiers. 1995. Brief report: absence of intact *nef* sequences in a long-term survivor with nonprogressive HIV-1 infection. *N. Engl. J. Med.* 332:228.
- Chowers, M. Y., C. A. Spina, T. J. Kwok, N. J. Fitch, D. D. Richman, and J. C. Guatelli. 1994. Optimal infectivity in vitro of human immunodeficiency virus type 1 requires an intact *nef* gene. *J. Virol.* 68:2906.
- Chowers, M. Y., M. W. Pandori, C. A. Spina, D. D. Richman, and J. C. Guatelli. 1995. The growth advantage conferred by HIV-1 *nef* is determined at the level of viral DNA formation and is independent of CD4 downregulation. *Virology* 212:451.
- Miller, M. D., M. T. Warmerdam, I. Gaston, W. C. Greene, and M. B. Feinberg. 1994. The human immunodeficiency virus-1 *nef* gene product: a positive factor for viral infection and replication in primary lymphocytes and macrophages. *J. Exp. Med.* 179:101.
- Miller, M. D., M. T. Warmerdam, K. A. Page, M. B. Feinberg, and W. C. Greene. 1995. Expression of the human immunodeficiency virus type 1 (HIV-1) *nef* gene during HIV-1 production increases progeny particle infectivity independently of gp160 or viral entry. *J. Virol.* 69:579.
- Benson, R. E., A. Sanfridson, J. S. Ottinger, C. Doyle, and B. R. Cullen. 1993. Downregulation of cell surface CD4 expression by simian immunodeficiency virus *nef* prevents viral super infection. *J. Exp. Med.* 177:1561.
- Garcia, J. V., and A. D. Miller. 1991. Serine phosphorylation-independent down-regulation of cell-surface CD4 by *nef*. *Nature* 350:508.
- Guy, B., M. P. Kiény, Y. Riviere, C. L. Peuch, K. Dott, M. Girard, L. Mantagnier, and J. P. Lococq. 1987. HIV F/3' orf encodes a phosphorylated GTP-binding protein resembling an oncogene product. *Nature* 330:266.
- Aiken, C., J. Konner, N. R. Landau, M. E. Lenburg, and D. Trono. 1994. Nef induces CD4 endocytosis: requirement for a critical dileucine motif in the membrane-proximal CD4 cytoplasmic domain. *Cell* 76:853.
- Mariani, R., and J. Skowronski. 1993. CD4 down-regulation by *nef* alleles isolated from human immunodeficiency virus type I-infected individuals. *Proc. Natl. Acad. Sci. USA* 90:5549.
- Collins, K. L., B. K. Chen, S. A. Kalams, B. D. Walker, and D. Baltimore. 1998. HIV-1 Nef protein protects infected primary cells against killing by cytotoxic T lymphocytes. *Nature* 391:397.
- Le Gall, S., L. Erdtmann, S. Benichou, C. Berlioz-Torrent, L. Liu, R. Benarous, J.-M. Heard, and O. Schwartz. 1998. Nef interacts with the  $\mu$  subunit of clathrin adaptor complexes and reveals a cryptic sorting signal in MHC I molecules. *Immunity* 8:483.
- Schwartz, O., V. Marechal, S. Le Gall, F. Lemonnier, and J.-M. Heard. 1996. Endocytosis of major histocompatibility complex class I molecules is induced by the HIV-1 *nef* protein. *Nat. Med.* 2:338.
- Mangasarian, A., V. Pigué, J.-K. Wang, Y.-L. Chen, and D. Trono. 1999. Nef-Induced CD4 and major histocompatibility complex class I (MHC-I) down-regulation are governed by distinct determinants: N-Terminal  $\alpha$  helix and proline repeat of Nef selectively regulate MHC-I trafficking. *J. Virol.* 73:1964.
- Bresnahan, P. A., W. Yonemoto, S. Ferrell, D. Williams-Herman, R. Gelezianas, and W. C. Greene. 1998. A dileucine motif in HIV-1 Nef acts as an internalization signal for CD4 downregulation and binds the AP-1 clathrin adaptor. *Curr. Biol.* 8:1235.
- Craig, H. M., M. W. Pandori, and J. C. Guatelli. 1998. Interaction of HIV-1 Nef with the cellular dileucine-based sorting pathway is required for CD4 down-regulation and optimal viral infectivity. *Proc. Natl. Acad. Sci. USA* 95:11229.
- Greenberg, M., L. De Tulleo, I. Rapoport, J. Skowronski, and T. Kirchhausen. 1998. A dileucine motif in HIV-1 Nef is essential for sorting into clathrin-coated pits and for downregulation of CD4. *Curr. Biol.* 8:1239.
- Grzesiek, S., S. J. Stahl, P. T. Wingfield, and A. Bax. 1996. The CD4 determinant for downregulation by HIV-1 *nef* directly binds to *nef*: mapping of the *nef* binding surface by NMR. *Biochemistry* 35:10256.
- Hua, J., W. Blair, R. Truan, and B. R. Cullen. 1997. Identification of regions in HIV-1 *nef* required for efficient downregulation of cell surface CD4. *Virology* 231:231.
- Pigué, V., F. Gu, M. Foti, N. Demareux, J. Gruenberg, J.-L. Carpentier, and D. Trono. 1999. Nef-induced CD4 degradation: a diacidic-based motif in Nef functions as a lysosomal targeting signal through the binding of  $\beta$ -COP in endosomes. *Cell* 97:63.
- Greenberg, M. E., A. J. Iafate, and J. Skowronski. 1998. The SH3 domain-binding surface and an acidic motif in HIV-1 Nef regulate trafficking of class I MHC complexes. *EMBO J.* 17:2777.
- Robinson, M. S. 1997. Coats and vesicle budding. *Trends Cell Biol.* 7:99.
- Marks, M. S., L. Woodruff, H. Ohno, and J. S. Bonifacino. 1996. Protein targeting by tyrosine- and di-leucine-based signals: evidence for distinct saturable components. *J. Cell Biol.* 135:341.
- Marks, M. S., L. Woodruff, H. Ohno, and J. S. Bonifacino. 1996. Protein targeting by tyrosine- and di-leucine-based signals: evidence for distinct saturable components. *J. Cell Biol.* 135:341. Kirchhausen, T., J. S. Bonifacino, and H. Riezman. 1997. Linking cargo to vesicle formation: receptor tail interactions with coat proteins. *Curr. Opin. Cell Biol.* 9:488.
- Liu, S.-H., M. S. Marks, and F. M. Brodsky. 1998. A dominant-negative clathrin mutant differentially affects trafficking of molecules with distinct sorting motifs in the class II major histocompatibility complex (MHC) pathway. *J. Cell Biol.* 140:1023.
- Pigué, V., Y.-L. Chen, A. Mangasarian, M. Foti, J.-L. Carpentier, and D. Trono. 1998. Mechanism of Nef-induced CD4 endocytosis: Nef connects CD4 with the  $\mu$  chain of adaptor complexes. *EMBO J.* 17:2472.

31. Baur, A. S., E. T. Sawai, P. Dazin, W. J. Fanti, C. Cheng-Mayer, and B. M. Peterlin. 1994. HIV-1 Nef leads to inhibition or activation of T cells depending on its intracellular localization. *Immunity* 1:373.
32. Shugars, D. C., M. S. Smith, D. H. Glueck, P. V. Nantermet, F. Seillier-Moiseiwitsch, and R. Swanstrom. 1993. Analysis of human immunodeficiency virus type 1 *nef* gene sequences present in vivo. *J. Virol.* 67:4639.
33. Mangasarian, A., M. Foti, C. Aiken, D. Chin, J.-L. Carpentier, and D. Trono. 1997. The HIV-1 Nef protein acts as a connector with sorting pathways in the Golgi and at the plasma membrane. *Immunity* 6:66.
34. Lu, X., H. Yu, S.-H. Liu, F. M. Brodsky, and B. M. Peterlin. 1998. Interactions between HIV1 Nef and vacuolar ATPase facilitate the internalization of CD4. *Immunity* 8:647.
35. Rapoport, I., M. Miyazaki, W. Boo, B. Duckworth, L. C. Cantley, S. Shoelson, and T. Kirchhausen. 1997. Regulatory interactions in the recognition of endocytic sorting signals by AP-2 complexes. *EMBO J.* 16:2240.
36. Marks, M. S., H. Ohno, T. Kirchhausen, and J. S. Bonifacio. 1997. Protein sorting by tyrosine-based signals: adapting to the Ys and wherefores. *Trends Cell Biol.* 7:124.
37. Sorkin, A., and G. Carpenter. 1993. Interaction of activated EGF receptors with coated pit adaptins. *Science* 261:612.
38. Backer, J. M., S. E. Shoelson, M. A. Weiss, A. X. Hua, R. B. Cheatham, E. Haring, D. C. Cahill, and M. F. White. 1992. The insulin receptor juxtamembrane region contains two independent tyrosine/beta-turn internalization signals. *J. Cell Biol.* 118:831.
39. Chang, C. P., C. S. Lazar, B. J. Walsh, M. Komuro, J. F. Collawn, L. A. Kuhn, J. A. Tainer, I. S. Trowbridge, M. G. Farquhar, and M. G. Rosenfeld. 1993. Ligand-induced internalization of the epidermal growth factor receptor is mediated by multiple endocytic codes analogous to the tyrosine motif found in constitutively internalized receptors. *J. Biol. Chem.* 268:19312.
40. Johnson, K. F., and S. Kornfeld. 1992. The cytoplasmic tail of the mannose 6-phosphate/insulin-like growth factor-II receptor has two signals for lysosomal enzyme sorting in the Golgi. *J. Cell Biol.* 119:249.
41. Verhey, K. J., J. I. Yeh, and M. J. Birnbaum. 1995. Distinct signals in the GLUT4 glucose transporter for internalization and for targeting to an insulin-responsive compartment. *J. Cell Biol.* 130:1071.