


Ultra-pure antibodies for
in vivo research, targeting
immune checkpoints
and more.

EXPLORE

BioCell



 *The Journal of*
Immunology

Cutting Edge: TLR2-Deficient and MyD88-Deficient Mice Are Highly Susceptible to *Staphylococcus aureus* Infection

This information is current as of March 8, 2021.

Osamu Takeuchi, Katsuaki Hoshino and Shizuo Akira

J Immunol 2000; 165:5392-5396; ;

doi: 10.4049/jimmunol.165.10.5392

<http://www.jimmunol.org/content/165/10/5392>

References This article **cites 23 articles**, 11 of which you can access for free at:
<http://www.jimmunol.org/content/165/10/5392.full#ref-list-1>

Why *The JI*? Submit online.

- **Rapid Reviews! 30 days*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

**average*

Subscription Information about subscribing to *The Journal of Immunology* is online at:
<http://jimmunol.org/subscription>

Permissions Submit copyright permission requests at:
<http://www.aai.org/About/Publications/JI/copyright.html>

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://jimmunol.org/alerts>

The Journal of Immunology is published twice each month by
The American Association of Immunologists, Inc.,
1451 Rockville Pike, Suite 650, Rockville, MD 20852
Copyright © 2000 by The American Association of
Immunologists All rights reserved.
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



Cutting Edge: TLR2-Deficient and MyD88-Deficient Mice Are Highly Susceptible to *Staphylococcus aureus* Infection¹

Osamu Takeuchi,^{*†} Katsuaki Hoshino,^{*†} and Shizuo Akira^{2*†}

Toll-like receptor (TLR) family acts as pattern recognition receptors for pathogen-specific molecular patterns. We previously showed that TLR2 recognizes Gram-positive bacterial components whereas TLR4 recognizes LPS, a component of Gram-negative bacteria. MyD88 is shown to be an adaptor molecule essential for TLR family signaling. To investigate the role of TLR family in host defense against Gram-positive bacteria, we infected TLR2- and MyD88-deficient mice with *Staphylococcus aureus*. Both TLR2- and MyD88-deficient mice were highly susceptible to *S. aureus* infection, with more enhanced susceptibility in MyD88-deficient mice. Peritoneal macrophages from MyD88-deficient mice did not produce any detectable levels of cytokines in response to *S. aureus*. In contrast, TLR2-deficient macrophages produced reduced, but significant, levels of the cytokines, and TLR4-deficient macrophages produced the same amounts as wild-type cells, indicating that *S. aureus* is recognized not only by TLR2, but also by other TLR family members except for TLR4. *The Journal of Immunology*, 2000, 165: 5392–5396.

A bacterial infection is one of the major causes of death even in the present day. *Streptococcus aureus*, a Gram-positive extracellular growing bacteria, is a major source of mortality in medical facilities (1). It causes a wide range of infections from skin infection to life-threatening diseases including abscesses of various organs, pneumonia, osteomyelitis, endocarditis, arthritis, and sepsis.

Initial host defense against bacterial infection is executed by innate immunity, characterized by the use of germline-encoded receptors for pathogen recognition (2). In *Drosophila*, Toll family

plays a key role in antifungal and antibacterial host defense (3). Recently, mammalian homologues of Toll, designated as Toll-like receptors (TLRs),³ were discovered (4–6). Each TLR is a type I transmembrane receptor possessing an extracellular leucine-rich repeat and a cytoplasmic Toll/IL-1 receptor homology domain. They are expected to act as pattern recognition receptors, which distinguish pathogen-associated molecular patterns, such as LPS, peptidoglycan (PGN), and lipoprotein (2). Among TLR family members, both TLR2 and TLR4 have been shown to recognize bacterial components. A mutation in the *Tlr4* gene is responsible for the phenotype of the C3H/HeJ mouse strain, which is unresponsiveness to LPS, a component of the outer membrane of Gram-negative bacteria (7). Targeted disruption of the *Tlr4* gene resulted in abrogation of responses to LPS and lipoteichoic acid (8, 9). In contrast, TLR2 is implicated in the recognition of Gram-positive bacterial components, bacterial lipoproteins, and zymosan (10–15). TLR2-deficient mice displayed impaired cytokine production in response to *Staphylococcus aureus* PGN preparation and mycoplasmal lipopeptide (9, 16).

MyD88 is a cytoplasmic adaptor molecule essential for the signaling of IL-1R/TLR family. Ligand binding to IL-1R/TLR family results in the recruitment of MyD88 to Toll/IL-1 receptor domains of receptors, which bridges the signal to IL-1R-associated kinase. Ultimately, the activation of a transcription factor NF- κ B occurs and permits the transactivation of proinflammatory cytokine genes (17, 18). We have generated MyD88-deficient mice and shown that MyD88 is essential for cellular responses to IL-1, IL-18, and many bacterial cell wall components such as LPS, PGN, and lipopeptide (19–21).

C3H/HeJ mice are known to be highly sensitive to infection with Gram-negative bacteria, owing to the failure of LPS recognition (22). Although TLR2 participates in the recognition of Gram-positive bacteria, a substantial role of TLR2 in host defense against Gram-positive bacteria was still unclear. In the present study, we investigated the role of TLR2 and MyD88 in *S. aureus* infection using mutant mice deficient in these molecules.

Materials and Methods

Mice

The mutant mouse (F₂ interbred from 129/Ola \times C57BL/6) strains deficient in TLR2, TLR4, or MyD88 were generated by gene targeting as

^{*}Department of Host Defense, Research Institute for Microbial Diseases, Osaka University, Osaka, Japan; and [†]Core Research for Evolutional Science and Technology of Japan Science and Technology Corporation, Osaka, Japan

Received for publication May 31, 2000. Accepted for publication September 13, 2000.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This work was supported by grants from the Ministry of Education of Japan.

² Address correspondence and reprint requests to Dr. Shizuo Akira, Department of Host Defense, Research Institute for Microbial Diseases, Osaka University, 3-1 Yamada-oka, Suita, Osaka 565-0871, Japan. E-mail address: sakira@biken.osaka-u.ac.jp

³ Abbreviations used in this paper: TLR, Toll-like receptor; PGN, peptidoglycan.




FIGURE 1. Survival of TLR2-deficient mice ($n = 10$, \square) and wild-type mice ($n = 10$, \circ) after i.v. infection of 1×10^7 CFU of *S. aureus*. $p < 0.03$ in log rank test.




FIGURE 3. Survival of MyD88-deficient mice ($n = 10$, \circ) and wild-type mice ($n = 10$, \square) after i.v. infection of 1×10^7 CFU of *S. aureus*. $p < 0.0003$ in log rank test.

described previously (8, 9, 19). MyD88-deficient mice were backcrossed eight times with C57BL/6 mice. For *S. aureus* infection, groups of TLR2-deficient mice and wild-type littermates or MyD88-deficient mice (C57BL/6 background) and wild-type littermates were used. Age-matched groups of wild-type, TLR2-, TLR4-, and MyD88-deficient mice were used for the other experiment.

Bacteria and infection

S. aureus 834 strain (gift from A. Nakane, Hiroasaki University, Japan) was classified as coagulase type II, produced toxic shock syndrome toxin I and methicillin resistant (23), and it was prepared as described previously (23, 24). In brief, bacteria were cultured on trypticase soy agar (Becton Dickinson, Sparks, MD), inoculated with trypticase soy broth, and incubated for 15 h at 37°C. The bacteria were collected and resuspended by PBS. The concentration of resuspended cells was adjusted spectrophotometrically at 550 nm. Mice were given i.v. injections of 0.2 ml of bacterial solution containing 1×10^7 CFU of viable *S. aureus* and their survival was monitored for up to 14 days. *S. aureus* suspension (1×10^9 CFU/ml in PBS) was boiled for 30 min and used as heat-killed *S. aureus*.

Determination of the number of bacteria in blood and organs

Mice were given 1×10^7 CFU or 1×10^6 CFU of *S. aureus* as an i.v. infection, and they were killed 1, 2, or 5 days later. Spleens and kidneys

were dissected, homogenized, and diluted in 10-fold steps in sterile water containing 0.5% Triton X-100 (Nacalai Tesque, Kyoto, Japan). Blood was also diluted in water containing 0.5% Triton X-100. Bacterial CFU was determined by plating each dilution on trypticase soy agar and was cultured for 24 h at 37°C.

Preparation of peritoneal macrophages

Mice were i.p. injected with 2 ml of 4% thioglycolate (Difco, Detroit, MI). Three days later, peritoneal exudate cells were isolated from the peritoneal cavity by washing with ice-cold HBSS (Life Technologies, Rockville, MD). Cells were cultured for 2 h and washed with HBSS to remove non-adherent cells. Adherent monolayer cells were used as peritoneal macrophages. Peritoneal macrophages (5×10^7 /ml) were cultured in RPMI 1640 medium (Nacalai Tesque) supplemented with 10% FCS and stimulated with 1×10^7 CFU/ml of heat-killed *S. aureus* for 24 h. Concentration of TNF- α in culture supernatant was determined by ELISA (Genzyme Techno, Minneapolis, MN) and IL-6 concentration was also measured by ELISA (Endogen, Boston, MA).

Statistical analysis

Kaplan-Meier plots were conducted and the log rank test was used to test the differences in the survival between wild-type and TLR2-deficient or

FIGURE 2. Numbers of live *S. aureus* in TLR2-deficient and wild-type mice after injection of bacteria. TLR2-deficient mice ($n = 4$ or 5) and wild-type mice ($n = 4$ or 5) were infected i.v. with 1×10^7 CFU (A) or 1×10^6 CFU (B) *S. aureus*. The blood and organs were collected 1 and 2 days (A) or 2 and 5 days (B) after injection, and bacterial counts were determined by culture on agar. Data are represented as the mean number of CFU \pm SD. *, $p < 0.05$ in Mann-Whitney U test.





FIGURE 4. Numbers of live *S. aureus* in MyD88-deficient and wild-type mice after injection of bacteria. MyD88-deficient mice ($n = 3$ or 4) and wild-type mice ($n = 3$ or 4) were infected i.v. with 1×10^7 CFU (A) or 1×10^6 CFU (B) *S. aureus*. The blood and organs were collected 1 (A) or 2 days and 5 days (B) after injection, and bacterial counts were determined by culture on agar. Data are represented as the mean number of CFU \pm SD. *, $p < 0.05$ in Mann-Whitney *U* test.



MyD88-deficient mice. The significances of the difference between the groups in the numbers of bacteria and cytokine concentrations were tested using Mann-Whitney *U* test.

Results

Decreased survival of TLR2-deficient mice after infection with *S. aureus*

We have previously shown that TLR2 is required for the recognition of PGN from *S. aureus*. Moreover, TLR2-deficient macrophages did not produce TNF- α in response to cell wall preparation from *S. aureus* (9). To evaluate the in vivo role of TLR2 against *S. aureus* infection, wild-type and TLR2-deficient mice were infected i.v. with 1×10^7 CFU of *S. aureus* and their survival was monitored. As shown in Fig. 1, all wild-type mice survived for 8 days after *S. aureus* inoculation and 60% survived on day 14. In contrast, about 80% of TLR2-deficient mice succumbed to *S. aureus* and died within 8 days, and only 10% survived on day 14 ($p < 0.03$). However, when a low dose (1×10^6 CFU) of *S. aureus* was administered in the mice, the survival was not altered between

wild-type and TLR2-deficient mice (data not shown). To investigate whether this susceptibility was the result of altered bacterial distribution and growth in vivo, the bacterial numbers in the blood, spleen, and kidney of mice were determined at 1 and 2 days after 1×10^7 CFU of *S. aureus* infection. At both 1 and 2 days, the number of *S. aureus* in the blood and kidney were higher in TLR2-deficient mice, consistent with high morbidity and mortality of TLR2-deficient mice after infection with *S. aureus* (Fig. 2A). When the mice were inoculated with 1×10^6 CFU of *S. aureus*, a statistically significant difference was not observed in the bacterial numbers between wild-type and TLR2-deficient organs at both 2 and 5 days after inoculum (Fig. 2B).

High susceptibility of MyD88-deficient mice to *S. aureus* infection

MyD88 is an adaptor molecule essential for the signaling of the IL-1R/TLR family. We have previously shown that MyD88-deficient mice lacked responsiveness to IL-1, IL-18, and LPS (19, 20). *S. aureus* PGN-induced TNF- α production was also abrogated in MyD88-deficient macrophages, indicating the role of MyD88 in




FIGURE 5. Production of TNF- α and IL-6 in response to heat-killed *S. aureus* in TLR2-, TLR4-, and MyD88-deficient macrophages. Peritoneal macrophages from wild-type, TLR2-, TLR4-, and MyD88-deficient mice were cultured in the presence of 1×10^7 CFU/ml of heat-killed *S. aureus* for 24 h. Concentrations of TNF- α (A) and IL-6 (B) in the culture supernatants were measured by ELISA. The results are representative of three independent experiments and are shown as the mean \pm SEM of triplicate wells. *, $p < 0.05$ in Mann-Whitney *U* test. N.D., Not detected.

anti-Gram-positive bacterial host defense (21). Therefore, we infected MyD88-deficient mice with *S. aureus* and monitored their survival. After inoculation with 1×10^7 *S. aureus*, all MyD88-deficient mice succumbed to infection and died within 5 days, whereas 80% of wild-type mice survived for 14 days ($p < 0.0003$, Fig. 3). MyD88-deficient mice died more rapidly than TLR2-deficient mice. Next, the number of bacterial cells in the blood, spleen, and kidney were determined at 1 day of 1×10^7 CFU *S. aureus* infection. Bacterial numbers in the blood and kidney were increased in MyD88-deficient mice compared with wild-type mice (Fig. 4A). Even when the mice were inoculated with a lower dose of bacteria (1×10^6 CFU), the bacterial numbers in the organs from MyD88-deficient mice were significantly higher than those from wild-type mice (Fig. 4B). These results indicate that MyD88-deficient mice are more susceptible to *S. aureus* infection than TLR2-deficient mice.

Role of TLR2 and MyD88 in the cytokines production in response to heat-killed *S. aureus*

Furthermore, we examined the responsiveness of peritoneal macrophages from wild-type, TLR2-, and MyD88-deficient mice to heat-killed *S. aureus*. Thioglycolate-elicited peritoneal macrophages were cultured in the presence of 1×10^7 CFU/ml of *S. aureus* for 24 h, and concentrations of TNF- α and IL-6 in culture supernatant were measured. Macrophages from wild-type and TLR4-deficient mice produced almost the same amount of TNF- α in response to heat-killed *S. aureus*. TLR2-deficient macrophages produced a reduced, but significant level of TNF- α . In contrast, MyD88-deficient macrophages did not produce any detectable TNF- α (Fig. 5A). IL-6 production in response to *S. aureus* was also reduced in TLR2-deficient macrophages and abrogated in MyD88-deficient macrophages (Fig. 5B).

Discussion

The present study demonstrated the critical role of TLR2 and MyD88 against extracellular Gram-positive bacteria using knock-out mice. Both TLR2- and MyD88-deficient mice developed higher mortality compared with wild-type mice after inoculation with *S. aureus*. These are associated with a bacterial growth in the blood and kidney. We have previously shown that *S. aureus* PGN-induced cytokine production was dependent on TLR2 (9). Thus, failure of recognizing invading bacteria is responsible for high susceptibility in TLR2-deficient mice. It was reported that C3H/HeJ mice, possessing a point mutation in the *Tlr4* gene, are highly susceptible to infection with *Salmonella typhimurium*. LD₅₀ value of the bacteria for C3H/HeJ mouse strain was <2 , in contrast to the resistance of C3H/HeN mice to *S. typhimurium* infection (LD₅₀ = 1×10^4) (22). These observations indicate the distinct roles of TLR2 and TLR4 in discriminating invading pathogens.

MyD88-deficient mice were also highly susceptible to *S. aureus* infection and they were more susceptible to infection than TLR2-deficient mice. We have shown that MyD88-deficient mice display the defective response to many bacterial components and IL-1 family cytokines such as IL-1 and IL-18 (19). Both severely impaired bacterial recognition and failure of signaling mediated by IL-1 family cytokines may contribute to the susceptibility of MyD88-deficient mice to infection.

We have previously shown that *S. aureus* cell wall-mediated TNF- α production was fully TLR2 dependent (9). Therefore, it

was surprising that the production of proinflammatory cytokine was induced in response to heat-killed *S. aureus* in TLR2-deficient macrophages, although the level was reduced compared with that in wild-type macrophages. In contrast, bacteria-mediated TNF- α production was abrogated in MyD88-deficient cells. Although TLR4-deficient macrophages displayed impaired responsiveness to Gram-positive lipoteichoic acids, TLR4-deficient macrophages responded to heat-killed *S. aureus* to the same extent as wild-type cells, indicating that lipoteichoic acid on *S. aureus* cell surfaces may not significantly contribute to cellular activations. These results suggest that *S. aureus* is recognized not only by TLR2, but also by other TLR/IL-1R family members except for TLR4. A previous report demonstrated that the treatment with anti-TNF- α Ab to mice after inoculation of *S. aureus* increased the death rate (23). Our results also indicate the correlation between the *S. aureus*-induced TNF- α production and the resistance of mice to the infection.

Taken together, innate recognition of bacteria by TLR family members is quite important for eliminating invading bacteria. Particularly, TLR2 plays a crucial role in host defense against extracellular growing Gram-positive bacteria. Similarly, high mortality after *S. aureus* infection and complete abrogation of proinflammatory cytokine secretion in MyD88-deficient mice shows an essential role of MyD88 in resistance to Gram-positive bacterial infection.

Acknowledgments

We thank Dr. Akio Nakane for providing *S. aureus* 834 strain, Dr. Yoshihumi Takeda for helpful suggestions, and Dr. Hiroko Tsutsui for providing mice. We thank T. Aoki for excellent secretarial assistance and N. Tsuji and E. Nakatani for excellent technical assistance.

References

- Lowy, F. D. 1998. *Staphylococcus aureus* infections. *N. Engl. J. Med.* 339:520.
- Medzhitov, R., and C. A. Janeway Jr. 1997. Innate immunity: the virtues of a monoclinal system of recognition. *Cell* 91:295.
- Kopp, E. B., and R. Medzhitov. 1999. The Toll-receptor family and control of innate immunity. *Curr. Opin. Immunol.* 11:13.
- Medzhitov, R., P. Preston-Hurburt, and C. A. Janeway, Jr. 1997. A human homologue of the *Drosophila* Toll protein signals activation of adaptive immunity. *Nature* 388:394.
- Rock, F. L., G. Hardiman, J. C. Timans, R. A. Kastelein, and J. F. Bazan. 1998. A family of human receptors structurally related to *Drosophila* Toll. *Proc. Natl. Acad. Sci. USA* 95:588.
- Takeuchi, O., T. Kawai, H. Sanjo, N. G. Copeland, D. J. Gilbert, N. A. Jenkins, K. Takeda, and S. Akira. 1999. TLR6: a novel member of an expanding Toll-like receptor family. *Gene* 231:59.
- Poltorak, A., X. He, I. Smirnova, M.-Y. Liu, C. V. Huffel, X. Du, D. Birdwell, E. Alejos, M. Silva, C. Galanos, et al. 1998. Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in *Tlr4* gene. *Science* 282:2085.
- Hoshino, K., O. Takeuchi, T. Kawai, H. Sanjo, T. Ogawa, Y. Takeda, K. Takeda, and S. Akira. 1999. TLR4-deficient mice are hyporesponsive to LPS: evidence for TLR4 as the *Lps* gene product. *J. Immunol.* 162:3749.
- Takeuchi, O., K. Hoshino, T. Kawai, H. Sanjo, T. Ogawa, H. Takada, K. Takeda, and S. Akira. 1999. Differential roles of TLR2 and TLR4 in recognition of Gram-negative and Gram-positive bacterial cell wall components. *Immunity* 11:443.
- Schwandner, R., Dziarski, R., Wesche, H., Rothe, M., and Kirschning, C. J. Peptidoglycan- and lipoteichoic acid-induced cell activation is mediated by Toll-like receptor 2. *J. Biol. Chem.* 274:17406.
- Yoshimura, A., Lien, E., Ingalls, R. R., Tuomanen, E., Dziarski, R., and Golenbock, D. 1999. Recognition of Gram-positive bacterial cell wall components by the innate immune system occurs via Toll-like receptor 2. *J. Immunol.* 163:1.
- Brightbill, D. H., H. D. Libraty, R. S. Krutzik, B. R. Yang, T. J. Belisle, R. J. Bleharski, M. Maitland, V. M. Norgard, E. S. Plevy, T. S. Smale, et al. 1999. Host defense mechanisms triggered by microbial lipoproteins through Toll-Like receptors. *Science* 285:732.
- Aliprantis, O. A., R. B. Yang, R. M. Mark, S. Suggett, B. Devaux, D. J. Radolf, R. G. Klimpel, P. Godowski, and A. Zychlinsky. 1999. Cell activation and apoptosis by bacterial lipoproteins through Toll-like receptor-2. *Science* 285:736.

14. Lien, E., T. J. Sellati, A. Yoshimura, T. H. Flo, G. Rawadi, R. W. Finberg, J. D. Carroll, T. Espevik, R. R. Ingalls, J. D. Radolf and D. T. Golenbock. 1999. Toll-like receptor 2 functions as a pattern recognition receptor for diverse bacterial products. *J. Biol. Chem.* 274:33419.
15. Underhill, D. M., A. Ozinsky, A. M. Hajjar, A. Stevens, C. B. Wilson, M. Bassetti, and A. Aderem. 1999. The Toll-like receptor 2 is recruited to macrophage phagosomes and discriminates between pathogens. *Nature* 401:811.
16. Takeuchi, O., A. Kaufmann, K. Grote, T. Kawai, K. Hoshino, M. Morr, P. F. Muhlradt, and S. Akira. 2000. Preferentially the R-stereoisomer of the mycoplasmal lipopeptide macrophage-activating lipopeptide-2 activates immune cells through a Toll-like receptor 2- and MyD88-dependent signaling pathway. *J. Immunol.* 164:554.
17. Muzio, M., G. Natoli, S. Saccani, M. Levrero, and A. Mantovani. 1998. The human Toll signaling pathway: divergence of nuclear factor κ B and JNK/SAPK activation upstream of tumor necrosis factor receptor-associated factor 6 (TRAF6). *J. Exp. Med.* 187:2097.
18. Medzhitov, R., P. Preston-Hurlburt, E. Kopp, A. Stadlen, C. Chen, S. Ghosh, and C. A. Janeway, Jr. 1998. MyD88 is an adaptor protein in the hToll/IL-1 receptor family signaling pathways. *Mol. Cell* 2:253.
19. Adachi, O., T. Kawai, K. Takeda, M. Matsumoto, H. Tsutsui, M. Sakagami, K. Nakanishi, and S. Akira. 1998. Targeted disruption of the *MyD88* gene results in loss of IL-1- and IL-18-mediated function. *Immunity* 9:143.
20. Kawai, T., O. Adachi, T. Ogawa, K. Takeda, and S. Akira. 1999. Unresponsiveness of MyD88-deficient mice to endotoxin. *Immunity* 11:115.
21. Takeuchi, O., K. Takeda, K. Hoshino, O. Adachi, T. Ogawa and S. Akira. 2000. Cellular responses to bacterial cell wall components are mediated through MyD88-dependent signaling cascades. *Int. Immunol.* 12:113.
22. O'Brien, A. D., D. L. Rosenstreich, I. Scher, G. H. Campbell, R. P. MacDermott, and S. B. Formal. 1980. Genetic control of susceptibility to *Salmonella typhimurium* in mice: role of the LPS gene. *J. Immunol.* 124:20.
23. Nakane, A., M. Okamoto, M. Asano, M. Kohanawa, and T. Minagawa. 1995. Endogenous gamma interferon, tumor necrosis factor, and interleukin-6 in *Staphylococcus aureus* infection in mice. *Infect. Immun.* 63:1165.
24. Sasaki, S., T. Miura, S. Nishikawa, K. Yamada, M. Hirasue, and A. Nakane. 1998. Protective role of nitric oxide in *Staphylococcus aureus* infection in mice. *Infect. Immun.* 66:1017.