

Investigate small particles with unparalleled sensitivity
Amnis® CellStream® Flow Cytometry System

For Research Use Only. Not for use in diagnostic procedures.



Luminex®
complexity simplified.



Serum Amyloid P Component and C-Reactive Protein Mediate Phagocytosis Through Murine FcγRs

This information is current as of October 24, 2021.

Carolyn Mold, Hattie D. Gresham and Terry W. Du Clos

J Immunol 2001; 166:1200-1205; ;
doi: 10.4049/jimmunol.166.2.1200
<http://www.jimmunol.org/content/166/2/1200>

References This article **cites 37 articles**, 17 of which you can access for free at:
<http://www.jimmunol.org/content/166/2/1200.full#ref-list-1>

Why *The JI*? [Submit online.](#)

- **Rapid Reviews! 30 days*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

**average*

Subscription Information about subscribing to *The Journal of Immunology* is online at:
<http://jimmunol.org/subscription>

Permissions Submit copyright permission requests at:
<http://www.aai.org/About/Publications/JI/copyright.html>

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://jimmunol.org/alerts>

The Journal of Immunology is published twice each month by
The American Association of Immunologists, Inc.,
1451 Rockville Pike, Suite 650, Rockville, MD 20852
Copyright © 2001 by The American Association of
Immunologists All rights reserved.
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



Serum Amyloid P Component and C-Reactive Protein Mediate Phagocytosis Through Murine FcγRs¹

Carolyn Mold,* Hattie D. Gresham,*[†] and Terry W. Du Clos^{2†‡}

The pentraxins, serum amyloid P component (SAP) and C-reactive protein (CRP) are acute-phase serum proteins in mice and humans, respectively. Although SAP binds to DNA and chromatin and affects clearance of these autoantigens, no specific receptor for SAP has been identified. CRP is an opsonin, and we have shown that it binds to FcγR. Mice deficient in FcγR were used to assess the role of these receptors in phagocytosis by pentraxins using zymosan as a ligand. Phagocytosis of zymosan by bone marrow macrophages (BMM) was enhanced by opsonization with SAP or CRP. BMM from mice deficient in all three FcγR or in γ-chain ingested unopsonized zymosan, but phagocytosis of SAP- or CRP-opsonized zymosan was not enhanced. SAP binding to BMM from γ-chain-deficient mice was also greatly reduced, indicating little or no binding of SAP to FcγRII. SAP and CRP opsonized zymosan for phagocytosis by BMM from mice deficient in FcγRII or FcγRIII. SAP, but not CRP, opsonized zymosan for uptake by neutrophils that express only low levels of FcγRI. Together these results indicate that FcγRI and FcγRIII are receptors for SAP in the mouse. Opsonization of zymosan by CRP is mediated through FcγRI. Pentraxins are major proteins of the innate immune system and arose earlier in evolution than Igs. The use of FcγR by the pentraxins links innate and adaptive immunity and may have important consequences for processing, presentation, and clearance of the self-Ags to which these proteins bind. *The Journal of Immunology*, 2001, 166: 1200–1205.

Serum amyloid P component (SAP)³ is a major acute-phase reactant in the mouse (1). Baseline levels of SAP in the blood vary markedly among different mouse strains. However, all mice examined to date appear to undergo a very marked up-regulation of SAP in response to inflammation (2). SAP is the precursor of tissue amyloid P component, a constituent of all types of amyloid. However, it is unlikely that this is the primary function of this protein, as it tends to stabilize the pathogenic deposition of amyloid proteins.

SAP and C-reactive protein (CRP) are members of the pentraxin family of proteins. These proteins are characterized by cyclic pentameric structure, sequence homology, and calcium-dependent interactions with ligands (3). Like other molecules of the innate immune system, SAP and CRP recognize ligands from necrotic and apoptotic cells as well as determinants on microorganisms. SAP binds to microbial polysaccharides and matrix components through carbohydrate determinants including heparin, 6-phosphorylated mannose, 3-sulfated saccharides, and the 4,6-cyclic pyruvate acetal of galactose (4, 5). Other physiological ligands of SAP include DNA, chromatin, and histones in cell nuclei as well as in

solution (6). The interaction of SAP with nuclear Ags has been proposed to be a primary function of this molecule. This hypothesis is supported by findings in recently described SAP-deficient mice that spontaneously developed anti-nuclear Abs and glomerulonephritis (7). These studies, as well as earlier experiments in which SAP was injected or induced by an acute-phase stimulus (8), demonstrate a slower clearance of chromatin from the bloodstream in the presence of SAP. The reduced clearance rate of chromatin and nucleosomes in the presence of SAP was associated with increased uptake by the liver and decreased localization in the kidneys (8).

Despite these observations on the effects of SAP on chromatin clearance, there have been few studies of SAP binding to receptors on phagocytic cells and to our knowledge no demonstration that SAP can act as an opsonin. It has been reported that SAP binds to peritoneal macrophages from mice in a saturable manner (9). The binding was calcium dependent and inhibitable by mannose and mannose phosphates, leading the authors to conclude that the mannose 6-P receptor is the receptor for SAP. High- and low-affinity binding of SAP to human PMN has also been described (10). This binding was calcium dependent and could be inhibited by CRP and to a lesser extent by aggregated IgG.

We recently determined that CRP binds to both human and mouse receptors for IgG (FcγR) (11, 12). CRP has long been recognized as an opsonin, and like SAP it binds to both microbial determinants and components of damaged cells (13). Because of the similarities in structure and activities between SAP and CRP, we decided to determine whether SAP also binds to FcγR and to compare the receptors used by these two proteins for opsonization. We studied SAP and CRP opsonization of zymosan, because both SAP and CRP bind to zymosan and zymosan has been widely used in studies of phagocytosis.

Three classes of FcγR are present in mice (14). The high-affinity receptor for IgG, FcγRI, is expressed primarily on monocytes and macrophages in association with the γ-chain. Two low-affinity receptors for IgG have been described, FcγRII and FcγRIII. FcγRII is present on lymphocytes, macrophages, PMN, platelets, and mast

Department of *Molecular Genetics and Microbiology, [†]Veterans Affairs Medical Center, and [‡]Department of Medicine, University of New Mexico, Albuquerque, NM 87131

Received for publication June 23, 2000. Accepted for publication October 18, 2000.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This work was supported by the Veterans Administration and by National Institutes of Health Grant AI28358.

² Address correspondence and reprint requests to Dr. Terry W. Du Clos, Veterans Affairs Medical Center, Research Service 151, 1501 San Pedro SE, Albuquerque, NM 87108. E-mail address: tduclos@unm.edu

³ Abbreviations used in this paper: SAP, serum amyloid P component; BMM, bone marrow macrophages; CRP, C-reactive protein; GMFI, geometric mean fluorescence intensity; HSA, human serum albumin; PAB, PBS containing 0.1% BSA and 0.05% sodium azide; PEC, peritoneal exudate cells; PEGAM, PE-conjugated F(ab')₂ goat anti-mouse IgG; PI, phagocytic index; PMN, neutrophils; LCM, L cell conditioned medium.

cells. Macrophages from mice expressing only Fc γ RII bind but do not ingest IgG-coated E (15). Fc γ RIII is also associated with the γ -chain and is expressed as a transmembrane protein on PMN, macrophages, NK cells, and mast cells. Fc γ RIII, like Fc γ RI, is capable of mediating phagocytosis. The relative contributions of the different Fc γ R classes to inflammation and phagocytosis have recently been elucidated by the development of mice genetically deficient in Fc γ R (15–17). These deficient mice were used in the current study to examine the role of Fc γ R in phagocytosis of zymosan opsonized by CRP or SAP.

The results demonstrate opsonization of zymosan by both SAP and CRP and that phagocytosis of SAP- and CRP-opsonized zymosan proceeds through phagocytic Fc γ R. In the case of CRP, phagocytosis proceeds through Fc γ RI only. However, phagocytosis of SAP-opsonized zymosan is mediated by either Fc γ RI or Fc γ RIII. These studies identify the phagocytic SAP receptors in the mouse and provide a possible mechanism for the SAP-mediated clearance of chromatin by Kupffer cells. They further point to distinct differences in specificity of CRP and SAP for Fc γ R. These results may help elucidate the common and disparate activities of CRP and SAP in the inflammatory response.

Materials and Methods

Reagents and Abs

FITC-zymosan was obtained from Molecular Probes (Eugene, OR). PE-conjugated F(ab')₂ goat anti-mouse IgG (PEGAM) and PE-conjugated F(ab')₂ goat anti-rat IgG were purchased from Caltag (Burlingame, CA). The hybridoma cell line producing rat mAb 2.4G2 was purchased from American Type Culture Collection (Manassas, VA), and the mAb was purified from culture supernatants by affinity chromatography on protein G-agarose. The mAb SAP-5, a murine IgG2a anti-human SAP, was purchased from Sigma (St. Louis, MO) and was used as ascites at 5 μ g mAb/ml for immunofluorescence. Mouse IgG1 and IgG2a isotype control mAb were obtained from Sigma. Rat anti-mouse F4/80 (Serotec, Raleigh, NC) and Ly-6G (PharMingen, San Diego, CA) were used to identify macrophages and neutrophils (PMN) in peritoneal exudate cells (PEC) and bone marrow cell preparations. Mouse anti-sheep E Ab of the IgG2b isotype was obtained from Accurate Scientific (Westbury, NY). Sheep blood was obtained from Colorado Serum Company (Denver, CO). FMLP was obtained from Sigma.

Isolation of CRP and SAP

Human CRP was purified from pleural fluid by affinity chromatography and ion exchange chromatography as previously described (18). SAP was prepared as a side product of factor IX purification and generously provided to us by Dr. Walter Kisiel (Department of Pathology, University of New Mexico, Albuquerque, NM). Briefly, SAP was copurified with coagulation factor IX from Proplex by immunoaffinity chromatography as described (19). SAP was then separated from factor IX by Q Sepharose Fast Flow (Pharmacia, Piscataway, NJ) chromatography. Minor contaminants were removed from SAP by anion exchange chromatography on a Mono Q column using FPLC (Pharmacia, Piscataway, NJ) with a 0.15–0.5 M NaCl gradient in 20 mM Tris, pH 7.8 (20).

Mice

Male C57BL/6Ncr mice were purchased from the National Cancer Institute (Frederick, MD) and were used as controls for the γ -chain-deficient and Fc γ RIII-deficient mice. Male B6 \times 129 F₂J mice were purchased from The Jackson Laboratory (Bar Harbor, ME) and were used as controls for the Fc γ RII- and the γ -chain/Fc γ RII-deficient mice. Male γ -chain-deficient (15), Fc γ RII-deficient (17), and γ -chain/Fc γ RII-deficient mice were purchased from Taconic Farms (Westminster, NY). Fc γ RIII-deficient mice were purchased from The Jackson Laboratory. All mice were housed conventionally and were used between 3 and 5 mo of age. Paired cultures of bone marrow macrophages (BMM) from age-matched deficient and control mice were compared in each phagocytosis experiment. All animal studies were approved by the Veterans Affairs Animal Committee Review Board.

Cells

BMM were prepared as previously described (21). Briefly, mice were killed, femurs were isolated under sterile conditions, the ends of the femurs

were excised and bone marrow was flushed from the femurs in HBSS with 0.2% human serum albumin (HSA). Cells were resuspended in 12–15 ml DMEM with 2% FBS and 2% L cell conditioned medium (LCM, a source of M-CSF) and allowed to adhere to tissue culture dishes for 2 h. Nonadherent cells were removed and cultured in tissue culture flasks in DMEM, 2% FBS, 15% LCM for 7–10 days. Nonadherent cells were removed after overnight culture, and cultures were fed after 4 days. BMM were removed from tissue culture flasks with 10 mM EDTA for use in binding or phagocytosis assays. BMM were resuspended in MEM, 2% FBS, 15% LCM with 1 mM MnCl₂ for phagocytosis assays or in PBS containing 0.1% BSA and 0.05% sodium azide (PAB) for staining.

Bone marrow PMN were purified from C57BL/6- and γ -chain-deficient mice using NIM-2 isolation medium (Cardinal Associates, Santa Fe, NM). Bone marrow leukocytes were resuspended in RPMI 1640 with 10 mM HEPES, 5% FCS, and 1 mM MnCl₂ at 4 \times 10⁶ PMN/ml. The percentage of PMN was determined by cresyl violet staining and was 49% for wild-type and 48% for deficient mice.

Elicited PEC were obtained by peritoneal lavage of mice injected with 1 ml of 3% thioglycollate medium (Difco, Detroit, MI) 5 days prior to harvest. Cells were collected in 10 ml of HBSS without calcium and magnesium. PEC prepared in this way were >50% macrophages determined by staining with DiffQuik (Dade International, Miami, FL).

Opsonization

Sheep E were opsonized with mouse IgG2b or IgG2a monoclonal anti-sheep E as previously described (22), washed, and resuspended at 5 \times 10⁸/ml in dextrose-gelatin Veronal-buffered saline. FITC-zymosan was incubated for 30 min at 37°C with 100 μ g/ml SAP or CRP in HBSS containing CaCl₂ and MgCl₂, washed, and resuspended at 4 \times 10⁸/ml in the same buffer.

Phagocytosis assays

To measure phagocytosis 4 \times 10⁵ BMM or PMN in 100 μ l culture medium were centrifuged briefly (1 min at 200 \times g) with opsonized FITC-zymosan (2 \times 10⁶/tube for BMM and 4 \times 10⁶/tube for PMN) and incubated for 30 min at 37°C. The fluorescence of the noningested yeast particles was quenched by the addition of trypan blue to a final concentration of 0.02%. Phagocytosis was scored visually on a Zeiss Axiovert fluorescent microscope. The results are expressed as the phagocytic index (PI) (number of ingested yeast cells or E per 100 BMM or PMN). For E-IgG2b or E-IgG2a, 7.5 \times 10⁵/tube were added, and noningested E were lysed prior to analyzing phagocytosis by phase microscopy.

The relative PI was calculated by dividing the PI of opsonized zymosan by the PI of unopsonized zymosan. Mean values of different mouse strains were compared by *t* tests.

SAP and CRP binding assays

BMM were released from culture flasks with EDTA, washed, and incubated with the indicated concentrations of SAP in PAB for 1 h on ice. Cells were washed twice with PAB and incubated with mAb SAP-5 or rat anti-mouse F4/80. Following a 30-min incubation on ice, cells were washed twice and incubated for 30 min with PE-conjugated secondary Abs. Cells were washed twice, resuspended in PAB, and analyzed on a Becton Dickinson FACScalibur flow cytometer (Mountain View, CA). For inhibition experiments, heat-aggregated human IgG (63°C for 45 min) was added to cells during the incubation with SAP. Forward and side scatter characteristics of BMM were determined by F4/80 binding and used to collect cells for fluorescent analysis.

Zymosan was washed with HBSS (containing calcium and magnesium), counted, and distributed into tubes. Zymosan (10⁶ particles/tube) was incubated with the indicated concentrations of CRP or SAP for 30 min at 37°C. Zymosan was washed and binding determined by incubation with 5 μ g/ml mAb (2C10 for CRP and SAP-5 for SAP) for 30 min on ice. After washing, zymosan was incubated with PEGAM and analyzed on a Becton Dickinson FACScalibur flow cytometer.

Fluorescence was analyzed using CellQuest software (Becton Dickinson). Fluorescence data were collected on a log scale, and data are reported as the difference in geometric mean fluorescent intensity (GMFI) in the presence of CRP or SAP compared with mAb and PEGAM alone. Binding curves were generated by nonlinear regression analysis using GraphPad Prism software (GraphPad Software, San Diego, CA).

Results

SAP and CRP binding to zymosan

As SAP closely resembles CRP in both structural and binding characteristics and because we have recently demonstrated that

CRP binds to Fc γ R, we decided to test whether SAP was capable of enhancing the phagocytosis of zymosan. Although previous studies have demonstrated the ability of both CRP and SAP to bind to zymosan, the optimal concentrations for binding of the two pentraxins were unknown. Therefore, preliminary studies were performed to examine the binding of SAP and CRP to zymosan (Fig. 1). SAP and CRP bound to zymosan with similar avidity. The total amount of pentraxin bound at saturation was comparable. The normal concentration of SAP in human serum is $\sim 50 \mu\text{g/ml}$ (23), and CRP concentrations range from $<5 \mu\text{g/ml}$ at baseline to $>200 \mu\text{g/ml}$ during an acute-phase response (24). For subsequent experiments, FITC-zymosan was opsonized with $100 \mu\text{g/ml}$ SAP or CRP.

Enhanced phagocytosis of SAP-opsonized zymosan requires Fc γ R

Pretreatment of FITC-zymosan with different concentrations of SAP enhanced its phagocytosis by BMM in a dose-dependent manner (data not shown). Maximal enhancement of phagocytosis was seen at $\sim 50 \mu\text{g/ml}$ SAP, consistent with the binding data. To determine whether Fc γ R were involved in the phagocytosis of particles opsonized by SAP, BMM from various strains of Fc γ R-deficient mice were tested. BMM from mice deficient in all three Fc γ R were cultured in parallel with BMM from control mice. FITC-zymosan was opsonized with $100 \mu\text{g/ml}$ of SAP or buffer, and phagocytosis was measured (Fig. 2). Uptake of zymosan by BMM from wild-type mice was enhanced following opsonization with SAP. BMM from Fc γ R-deficient mice ingested unopsonized zymosan as efficiently as wild-type BMM, but opsonization with SAP did not enhance ingestion. Significant phagocytosis of unopsonized zymosan by BMM occurred due to the uptake of zymosan through other receptors. However, SAP enhanced the level of phagocytosis of zymosan by BMM from wild-type but not Fc γ R-deficient mice in three experiments.

To determine which Fc γ R were most important for the uptake of opsonized zymosan, three additional strains of mice were tested. First, phagocytosis of zymosan was examined in γ -chain-deficient mice. These mice do not express Fc γ RI or Fc γ RIII, both of which require the γ -chain for expression. Macrophages from γ -chain-deficient mice express only Fc γ R_{II} and do not phagocytose E-IgG (15). As expected, these mice also showed markedly decreased phagocytosis of SAP-opsonized zymosan (Fig. 3). The ability of CRP to opsonize FITC-zymosan was examined in the same assay. Consistent with our previous identification of Fc γ RI and Fc γ R_{II} as CRP receptors in the mouse (12), CRP opsonized zymosan for phagocytosis by wild-type but not γ -chain-deficient BMM.

The phagocytosis of SAP- and CRP-opsonized FITC-zymosan was also examined using BMM from Fc γ R_{II}-deficient mice. En-




FIGURE 1. SAP and CRP binding to zymosan. Zymosan was incubated with the indicated concentrations of SAP or CRP for 30 min at 37°C , washed, and binding was detected using anti-SAP mAb SAP-5 or anti-CRP mAb 2C10 and PEGAM. The GMFI was determined by flow cytometry. Results from a representative experiment are shown.




FIGURE 2. SAP does not opsonize zymosan for phagocytosis by BMM from Fc γ R-deficient mice. FITC-zymosan was incubated with $100 \mu\text{g/ml}$ SAP or HBSS for 30 min at 37°C and then washed once with HBSS. Opsonized zymosan was then incubated with BMM at a ratio of 4:1. BMM and FITC-zymosan were centrifuged to enhance contact and incubated for 30 min at 37°C . Trypan blue was added to quench the fluorescence of uningested FITC-zymosan. Phagocytosis was scored by fluorescent microscopy. Results are expressed as the number of FITC-zymosan per 100 BMM (PI). Results shown are representative of four separate experiments done with paired wild-type and deficient BMM.

hanced phagocytosis of FITC-zymosan mediated by either SAP or CRP was unchanged compared with wild-type controls (Fig. 4).

A statistical comparison of the ability of SAP to opsonize zymosan for uptake by BMM from normal and Fc γ R-deficient mice is shown in Table I.

SAP binding to mouse macrophages requires γ -chain expression

SAP binding to wild-type and γ -chain-deficient BMM was determined by flow cytometry using mAb SAP-5 and PEGAM (Fig. 5). The apparent K_D of the interaction of SAP with wild-type BMM determined by nonlinear regression analysis of the binding curve was $\sim 7.5 \times 10^{-7} \text{ M}$. BMM from γ -chain-deficient mice bound much less SAP than did wild-type BMM. SAP binding to F4/80-positive PEC from γ -chain-deficient mice was also greatly reduced compared with the same PEC from wild-type mice (data not shown). The binding of $50 \mu\text{g/ml}$ SAP to PEC was inhibited




FIGURE 3. SAP and CRP do not opsonize zymosan for phagocytosis by BMM from γ -chain-deficient mice. FITC-zymosan was incubated with $100 \mu\text{g/ml}$ SAP or CRP or with HBSS for 30 min at 37°C and then washed once with HBSS. Opsonized zymosan was then incubated with BMM at a ratio of 4:1. BMM and FITC-zymosan were centrifuged to enhance contact and incubated for 30 min at 37°C . Trypan blue was added to quench the fluorescence of uningested FITC-zymosan. Phagocytosis was scored by fluorescent microscopy. Results are expressed as the number of FITC-zymosan per 100 BMM (PI). A representative experiment (of three) is shown.




FIGURE 4. Fc γ RII-deficient mice show no decrease in the level of phagocytosis of FITC-zymosan opsonized by SAP or CRP. FITC-zymosan was incubated with 100 μ g/ml SAP or CRP or with HBSS for 30 min at 37°C and then washed once with HBSS. Opsonized zymosan was then incubated with BMM at a ratio of 4:1. BMM and FITC-zymosan were centrifuged to enhance contact and incubated for 30 min at 37°C. Trypan blue was added to quench the fluorescence of uningested FITC-zymosan. Phagocytosis was scored by fluorescent microscopy. Results are expressed as the number of FITC-zymosan per 100 BMM (PI). A representative experiment (of two) is shown.

>95% in the presence of 100 μ g/ml aggregated IgG, further supporting a requirement for Fc γ R in SAP binding. These results differ from previous binding studies using CRP in which significant binding to peritoneal macrophages from γ -chain-deficient mice was observed indicating binding to Fc γ RII (12). Thus, SAP shows dose-dependent and saturable binding to mouse macrophages with a moderate affinity. This interaction requires one or both receptors associated with the γ -chain. Because SAP is multivalent and its binding to microbial cells occurs through highly repetitive determinants on the surface, the physiological interaction between SAP and cellular receptors is likely to involve multiple receptors resulting in increased avidity of binding.

Enhanced phagocytosis of SAP- and CRP-opsonized zymosan can be mediated by Fc γ RI

Both Fc γ RI and Fc γ RIII are capable of mediating phagocytosis *in vivo*, and there is evidence that Fc γ RIII may be more efficient than Fc γ RI (25). To examine the role of Fc γ RI, the ability of BMM derived from Fc γ RIII-deficient mice to phagocytose zymosan was examined. As shown in Fig. 6 and Table I, BMM from these mice showed little or no decrease in their ability to phagocytose zymosan opsonized by SAP or CRP. These results suggest that the phagocytosis of zymosan via SAP and CRP can be mediated by Fc γ RI. Whether Fc γ RIII also contributes to the uptake of zymosan opsonized with SAP or CRP cannot be determined from this experiment. Soluble IgG1, IgG2a, and the rat mAb 2.4G2 to mouse

Table I. Comparison of phagocytosis of SAP-opsonized zymosan between mouse strains

Mouse Strain	Relative PI ^a (Mean)	SEM	n	p ^b
C57BL/6	1.97	0.23	5	–
γ -chain knockout	1.20	0.14	3	0.0055
Fc γ RIII knockout	1.68	0.11	3	0.31
B6 \times 129	1.71	0.09	4	–
Double knockout	0.62	0.14	4	<0.0001
Fc γ RII knockout	1.81	0.26	2	0.59

^a The relative PI (PI SAP-zymosan/PI unopsonized zymosan) for SAP-opsonized zymosan compared to unopsonized zymosan. PI for unopsonized zymosan were not significantly different for γ -chain knockout or double knockout mice compared to control mice.

^b Values of p were determined by t tests between the deficient strain and the appropriate control strain.




FIGURE 5. BMM from γ -chain-deficient mice show decreased SAP binding. BMM from C57BL/6 mice or γ -chain-deficient mice were incubated with SAP in PAB for 60 min on ice. SAP binding was detected using mAb SAP-5 and PEGAM. The increase in GMFI relative to cells incubated with mAb and PEGAM is shown.

Fc γ RII/III all failed to inhibit SAP-mediated phagocytosis by wild-type BMM (data not shown).

The ability of Fc γ RI to mediate phagocytosis of CRP- or SAP-opsonized zymosan was also examined by selective digestion of Fc γ RI from BMM with trypsin. Trypsin has been shown to selectively remove Fc γ RI from mouse macrophages (26). BMM were treated with 20 μ g/ml trypsin for 30 min at 37°C, and phagocytosis of zymosan, opsonized zymosan and E-IgG2a was tested (26). The phagocytosis of E-IgG2a was decreased by 70%, indicating a loss of Fc γ RI, whereas ingestion of unopsonized zymosan was unchanged, indicating that general phagocytic responses were not affected. With this loss of Fc γ RI, the phagocytosis of CRP-zymosan was decreased by 55% and the phagocytosis of SAP-zymosan was decreased by 24%. Thus, CRP-mediated opsonization of zymosan appeared to be dependent on the presence of Fc γ RI, whereas SAP-mediated opsonization was less affected. As Fc γ RII is not a phagocytic receptor on mouse BMM, it appears that phagocytosis of SAP-opsonized zymosan may proceed through either Fc γ RI or Fc γ RIII.

Enhanced phagocytosis of SAP- but not CRP-opsonized zymosan can be mediated by Fc γ RIII

As Fc γ RI α -chain-deficient mice have only recently been produced and were not available for these studies, we tested the phagocytosis of zymosan by PMN, which express low levels of




FIGURE 6. Fc γ RIII-deficient mice show no decrease in the level of phagocytosis of FITC-zymosan opsonized by SAP or CRP. FITC-zymosan was incubated with 100 μ g/ml SAP or CRP or with HBSS for 30 min at 37°C and then washed once with HBSS. Opsonized zymosan was then incubated with BMM at a ratio of 4:1. BMM and FITC-zymosan were centrifuged to enhance contact and incubated for 30 min at 37°C. Trypan blue was added to quench the fluorescence of uningested FITC-zymosan. Phagocytosis was scored by fluorescent microscopy. Results are expressed as the number of FITC-zymosan per 100 BMM (PI). A representative experiment (of three) is shown.

Fc γ RI, to further examine whether SAP- or CRP-opsonized zymosan could be taken up through Fc γ RIII. Bone marrow PMN bound very low levels of monomeric IgG2a and did not ingest E-IgG2a, indicating a lack of functional Fc γ RI on these cells (data not shown). PMN were isolated from bone marrow and incubated with SAP- or CRP-opsonized zymosan or E-IgG2b as a positive control for Fc γ RIII-mediated phagocytosis. The ability of SAP and CRP to opsonize zymosan for phagocytosis by PMN from wild-type and γ -chain-deficient mice was tested. Phagocytosis was also examined in the presence of FMLP, which enhances phagocytosis through Fc γ R (22). As shown in Fig. 7, ingestion of E-IgG2b and SAP-zymosan by PMN was greater for wild-type compared with γ -chain-deficient mice, indicating that phagocytosis was mediated by Fc γ RIII. In contrast to the results with BMM (Fig. 3), CRP-zymosan was poorly ingested by PMN, and there was no difference between PMN from control and γ -chain-deficient mice. This finding is consistent with the inability of CRP to bind to mouse Fc γ RIII and indicates that there is insufficient Fc γ RI present to mediate phagocytosis. Similar results were seen in the presence of FMLP, with higher levels of phagocytosis for E-IgG2b and SAP-zymosan, but no increase in the uptake of CRP-zymosan or unopsonized zymosan. Thus these results indicate that SAP can opsonize zymosan for phagocytosis through either Fc γ RIII (Fig. 7) or Fc γ RI (Fig. 6).

Discussion

The major findings reported here are first that SAP is an efficient opsonin for zymosan and second that SAP-mediated phagocytosis is dependent on the presence of receptors for IgG on mouse phagocytic cells. Another significant finding reported for the first time here is that opsonization for phagocytosis by CRP occurs through Fc γ R. Although CRP-enhanced phagocytosis was previously demonstrated in murine (27) and human phagocytes (28) a unique re-




FIGURE 7. SAP enhances phagocytosis of FITC-zymosan by PMN from C57BL/6 but not γ -chain-deficient mice. Bone marrow PMN were isolated on NIM-2 medium and resuspended in medium. In the FMLP-stimulated group, FMLP was added at 100 μ M. FITC-zymosan was incubated with 100 μ g/ml SAP or CRP for 30 min at 37°C and then washed once with HBSS. Opsonized FITC-zymosan or E-IgG2b was then added to PMN at a ratio of 10:1. The tubes were centrifuged to enhance contact and incubated for 30 min at 37°C. Uningested E-IgG2b were lysed. Trypan blue was added to quench the fluorescence of uningested FITC-zymosan. Phagocytosis was scored by microscopy. Results are expressed as the number of FITC-zymosan or E-IgG2b per 100 PMN (PI). A representative experiment is shown.

ceptor was felt to be responsible (27). These studies suggest that both CRP and SAP are opsonins that act through specific binding through Fc γ R on phagocytic cells.

SAP has been demonstrated to bind to a wide variety of ligands *in vitro* including heparin, laminin (29), heparan sulfate (30), fibronectin, and C4b binding protein (31) as well as a number of microbial polysaccharides (4, 5). The binding site is thought to be the calcium-dependent ligand binding site that specifically recognizes phosphoethanolamine. The relative physiological significance of these interactions is not known. SAP also binds to chromatin and DNA in the circulation (6). Recently, SAP has been shown to be an important mediator of the clearance and processing of chromatin (8). The development of autoantibodies to chromatin in SAP-deficient mice suggests an important role for this protein in the prevention of autoimmunity (7).

Saturable binding of SAP to phagocytic cells was previously demonstrated but the nature of the receptor was not identified (9, 10). Contrary to CRP, for which evidence of opsonic activity was presented early on, studies of the ability of SAP to enhance phagocytosis of bacterial or fungal pathogens are limited.

We have previously shown that CRP interacts with Fc γ RI and Fc γ RII in the mouse (12). Therefore, we wanted to determine whether the receptor for SAP was also an Fc γ R. The present results confirm that both SAP and CRP opsonize zymosan for phagocytosis by mouse BMM and that Fc γ R are required. The interaction of SAP with mouse Fc γ R differs from that of CRP. Both CRP and SAP are capable of binding Fc γ RI and Fc γ RI may be important for phagocytosis through both pentraxins. However, SAP was capable of enhancing phagocytosis in PMN, which express only low levels of Fc γ RI and in BMM after selective removal of Fc γ RI by trypsin, suggesting that Fc γ RIII is sufficient for SAP-mediated opsonization. Furthermore, SAP-mediated phagocytosis was enhanced by FMLP treatment of PMN. FMLP preferentially enhances phagocytosis through Fc γ R as opposed to complement receptors (22). This finding supports the role of Fc γ R in opsonization by SAP. Ongoing studies to be presented separately suggest that similar differences in Fc γ R binding specificity occur in the human as well. It is likely that these differences in receptor specificity will be reflected in different activities of SAP and CRP *in vivo*.

SAP binding was γ -chain dependent and inhibited by aggregated IgG, consistent with binding to Fc γ RI and Fc γ RIII. Both of these receptors can invoke a stimulatory response by the cell and are therefore likely to enhance the inflammatory response in the mouse. CRP, but not SAP, binds to Fc γ RII as well (12). Fc γ RII in mice is known to provide an anti-inflammatory signal through the recruitment of the inositol phosphatase Src homology 2 domain-containing inositol phosphatase (SHIP) (32). Although CRP binds to Fc γ RII, no increase in uptake of CRP-opsonized zymosan by BMM was observed in Fc γ RII-deficient mice. This is in contrast to the enhanced phagocytosis of E-IgG2b by PEC from Fc γ RII-deficient mice (33). These results may be explained by the interaction of CRP with Fc γ RI, whereas phagocytosis of E-IgG2b is predominantly mediated by Fc γ RIII.

The interaction of SAP with phagocytic receptors, its ability to activate complement, and its abundance in the serum suggest that SAP may play a role in host defense as well. SAP has been shown to bind a number of human pathogens, especially fungal organisms. Singh et al. (34) reported that SAP enhanced the ability of mouse macrophages to kill *Listeria monocytogenes*. This listericidal activity was not associated with enhanced ingestion of the bacteria by SAP or with SAP binding to *Listeria*. Thus the nature of this effect was not determined. CRP was previously shown to provide protection from infection with *Streptococcus pneumoniae* (35).

It is now known that both CRP and SAP bind to Fc γ R in the human and the mouse. The evolutionary significance of these findings is of considerable interest as the pentraxins appeared earlier in evolution than the Igs. The pentraxins are present in the horseshoe crab, whereas the Igs do not appear until the jawed fishes (36). Therefore, it is possible that Fc receptors preceded the appearance of Ig and served as receptors for pentraxins.

The interaction of CRP and SAP, which are both capable of interacting with altered self and with microorganisms, identify the pentraxins as innate immune recognition molecules. However, the finding that they interact with Fc γ R bridges the gap between innate and acquired immunity. This connection may have served to facilitate the development of the adaptive immune system. It remains to be determined whether SAP and CRP are most important for the recognition of foreign Ags as exemplified by the protection afforded mice by CRP (35) or whether their role in regulating autoimmunity as suggested by studies of CRP and SAP in down-regulating the autoimmune response in systemic lupus erythematosus is more important (7, 37).

The function of SAP has long remained unknown despite extensive studies of its binding activities, its interaction with the complement system, and its role in the pathogenesis of amyloidosis. The finding that SAP can specifically interact with Fc γ R should help to delineate the *in vivo* functions of this ancient member of the innate immune system.

Acknowledgments

We thank Dr. Walter Kisiel (University of New Mexico, Albuquerque, NM) for providing SAP, Larry Potempa (ImmTech, Evanston, IL) for mAb 2C10, and Ruobing Xiao for excellent technical assistance.

References

- Pepys, M. B., M. L. Baltz, K. Gomer, J. S. Davis, and M. Doenhoff. 1979. Serum amyloid P component is an acute phase reactant in mouse. *Nature* 278:259.
- Mortensen, R. F., K. Beisel, N. J. Zeleznik, and P. T. Le. 1983. Acute phase reactants of mice II. Strain dependence of serum amyloid P-component (SAP) levels and response to inflammation. *J. Immunol.* 130:885.
- Gewurz, H., C. Mold, J. Siegel, and B. Fiedel. 1982. C-reactive protein and the acute phase response. *Adv. Int. Med.* 27:345.
- Hind, C. R. K., P. M. Collins, D. Renn, R. B. Cook, D. Caspi, M. L. Baltz, and M. B. Pepys. 1984. Binding specificity of serum amyloid P component for the pyruvate acetal of galactose. *J. Exp. Med.* 159:1058.
- Loveless, R. W., G. Floyd-O'Sullivan, J. G. Raynes, C.-T. Yuen, and T. Feizi. 1992. Human serum amyloid P component is a multispecific adhesive protein whose ligands include 6-phosphorylated mannose and the 3-sulphated saccharides galactose, N-acetylglucosamine and glucuronic acid. *EMBO J.* 11:813.
- Pepys, M. B., and P. J. G. Butler. 1987. Serum amyloid P component is the major calcium-dependent specific DNA binding protein of the serum. *Biochem. Biophys. Res. Commun.* 148:308.
- Bickerstaff, M. C. M., M. Botto, W. L. Hutchinson, J. Herbert, G. A. Tennent, A. Bybee, D. A. Mitchell, H. T. Cook, P. J. G. Butler, M. J. Walport, and M. B. Pepys. 1999. Serum amyloid P component controls chromatin degradation and prevents antinuclear autoimmunity. *Nat. Med.* 5:694.
- Burlingame, R. W., M. A. Volzer, J. Harris, and T. W. Du Clos. 1996. The effect of acute phase proteins on clearance of chromatin from the circulation of normal mice. *J. Immunol.* 156:4783.
- Siripont, J., J. M. Tebo, and R. F. Mortensen. 1988. Receptor-mediated binding of the acute phase reactant mouse serum amyloid P-component (SAP) to macrophages. *Cell. Immunol.* 117:239.
- Landsmann, P., O. Rosen, M. Pontet, M. Pras, D. Levaratowsky, E. G. Shephard, and M. Fridkin. 1994. Binding of human serum amyloid P component (hSAP) to human neutrophils. *Eur. J. Biochem.* 223:805.

- Bharadwaj, D., M. P. Stein, M. Volzer, C. Mold, and T. W. Du Clos. 1999. The major receptor for C-reactive protein on leukocytes is Fc γ RII. *J. Exp. Med.* 190:585.
- Stein, M. P., C. Mold, and T. W. Du Clos. 2000. C-reactive protein binding to murine leukocytes requires Fc γ receptors. *J. Immunol.* 164:1514.
- Du Clos, T. W. 1997. The interaction of C-reactive protein and serum amyloid P component with nuclear antigens. *Mol. Biol. Rep.* 23:253.
- Ravetch, J. V. 1994. Fc receptors: rubor redux. *Cell* 78:553.
- Takai, T., M. Li, D. Sylvestre, R. Clynes, and J. V. Ravetch. 1994. FcR γ chain deletion results in pleiotropic effector cell defects. *Cell* 76:519.
- Hazenbos, W. L. W., J. E. Gessner, F. M. A. Hofhuis, H. Kuipers, D. Meyer, A. F. M. Heijnen, R. E. S. M. Schmidt, P. J. A. Capel, M. Daeron, J. G. J. van de Winkel, and J. S. Verbeek. 1996. Impaired IgG-dependent anaphylaxis and Arthus reaction in Fc γ RIII (CD16) deficient mice. *Immunity* 5:181.
- Takai, T., M. Ono, M. Hikida, H. Ohmori, and J. V. Ravetch. 1995. Augmented humoral and anaphylactic responses in Fc γ RII-deficient mice. *Nature* 379:346.
- Du Clos, T. W., L. Zlock, and L. L. Marnell. 1991. Definition of a C-reactive protein binding determinant on histones. *J. Biol. Chem.* 266:2167.
- Spanier, T. B., C. O. Mehmet, O. P. Minanov, R. Simantov, W. Kisiel, D. M. Stern, E. A. Rose, and A. M. Schmidt. 1998. Heparinless cardiopulmonary bypass with active-site blocked factor IXA: a preliminary study on the dog. *J. Thor. Cardiovasc. Surg.* 115:1179.
- Saunero-Nava, L., J. E. Coe, C. Mold, and T. W. Du Clos. 1992. Hamster female protein binding to chromatin, histones and DNA. *Mol. Immunol.* 29:837.
- Roach, T., S. Slater, M. Koval, E. C. White, M. McFarland, M. Okumura, M. Thomas, and E. Brown. 1997. CD45 regulates Src family member kinase activity associated with macrophage integrin-mediated adhesion. *Curr. Biol.* 7:408.
- Gresham, H. D., C. J. Ray, and F. X. O'Sullivan. 1991. Defective neutrophil function in the autoimmune mouse strain: potential role of transforming growth factor- β . *J. Immunol.* 146:14611.
- Nelson, S. R., G. A. Tennent, D. Sethi, P. E. Gower, F. W. Ballardie, S. Amatayakul-Chantler, and M. B. Pepys. 1991. Serum amyloid P component in chronic renal failure and dialysis. *Clin. Chim. Acta* 200:191.
- Claus, D. R., A. P. Osmand, and H. Gewurz. 1976. Radioimmunoassay of human C-reactive protein and levels in normal sera. *J. Lab. Clin. Med.* 87:120.
- Fossati-Jimack, L., A. R. L. Ioan-Facsinay, Y. Chicheportiche, N. Watanabe, T. Saito, F. M. A. Hofhuis, J. E. Gessner, C. Schiller, R. E. Schmidt, T. Honjo, J. S. Verbeek, and S. Izui. 2000. Markedly different pathogenicity of four immunoglobulin G isotype-switch variants of an antierthrocyte autoantibody is based on their capacity to interact *in vivo* with the low-affinity Fc γ receptor III. *J. Exp. Med.* 191:1293.
- Unkeless, J. C. 1977. The presence of two Fc receptors on mouse macrophages: evidence from a variant cell line and differential trypsin sensitivity. *J. Exp. Med.* 145:931.
- Zahedi, K., J. M. Tebo, J. Siripont, G. F. Klimo, and R. F. Mortensen. 1989. Binding of human C-reactive protein to mouse macrophages is mediated by distinct receptors. *J. Immunol.* 142:2384.
- Mortensen, R. F., A. P. Osmand, T. F. Lint, and H. Gewurz. 1976. Interaction of C-reactive protein with lymphocytes and monocytes: complement-dependent adherence and phagocytosis. *J. Immunol.* 117:774.
- Zahedi, K. 1997. Characterization of the binding of serum amyloid P to laminin. *J. Biol. Chem.* 272:2143.
- Hamazaki, H. 1987. Ca²⁺-mediated association of human serum amyloid P component with heparan sulfate and dermatan sulfate. *J. Biol. Chem.* 262:1456.
- De Beer, F. C., M. L. Baltz, A. Holford, A. Feinstein, and M. B. Pepys. 1981. Fibronectin and C4-binding protein are selectively bound by aggregated amyloid P component. *J. Exp. Med.* 154:1134.
- Ono, M., S. Bolland, P. Tempst, and J. V. Ravetch. 1996. Role of the inositol phosphatase SHIP in negative regulation of the immune system by the receptor Fc γ RIB. *Nature* 383:263.
- Clynes, R., J. S. Maizes, R. Guinamard, M. Ono, T. Takai, and J. V. Ravetch. 1999. Modulation of immune complex-induced inflammation *in vivo* by the coordinate expression of activation and inhibitory Fc receptors. *J. Exp. Med.* 189:179.
- Singh, P. P., F. Gervais, E. Skamene, and R. F. Mortensen. 1986. Serum amyloid P component-induced enhancement of macrophage listericidal activity. *Infect. Immun.* 52:688.
- Mold, C., S. Nakayama, T. J. Holzer, H. Gewurz, and T. W. Du Clos. 1981. C-reactive protein is protective against *Streptococcus pneumoniae* infection in mice. *J. Exp. Med.* 154:1703.
- Robey, F. A., and T.-Y. Liu. 1981. Limulin: A C-reactive protein from *Limulus polyphemus*. *J. Biol. Chem.* 256:969.
- Du Clos, T. W., L. Zlock, P. S. Hicks, and C. Mold. 1994. Decreased autoantibody levels and enhanced survival of (NZB \times NZW) F₁ mice treated with C-reactive protein. *Clin. Immunol. Immunopath.* 70:22.