

This information is current as
of February 28, 2021.

Persistent Decreases in Blood Plasmacytoid Dendritic Cell Number and Function Despite Effective Highly Active Antiretroviral Therapy and Increased Blood Myeloid Dendritic Cells in HIV-Infected Individuals

Jihed Chehimi, Donald E. Campbell, Livio Azzoni, Darlene
Bacheller, Emmanouil Papasavvas, Ghassen Jerandi, Karam
Mounzer, Jay Kostman, Giorgio Trinchieri and Luis J.
Montaner

J Immunol 2002; 168:4796-4801; ;
doi: 10.4049/jimmunol.168.9.4796
<http://www.jimmunol.org/content/168/9/4796>

References This article **cites 32 articles**, 12 of which you can access for free at:
<http://www.jimmunol.org/content/168/9/4796.full#ref-list-1>

Why *The JI*? [Submit online.](#)

- **Rapid Reviews! 30 days*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

**average*

Subscription Information about subscribing to *The Journal of Immunology* is online at:
<http://jimmunol.org/subscription>

Permissions Submit copyright permission requests at:
<http://www.aai.org/About/Publications/JI/copyright.html>

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://jimmunol.org/alerts>

Persistent Decreases in Blood Plasmacytoid Dendritic Cell Number and Function Despite Effective Highly Active Antiretroviral Therapy and Increased Blood Myeloid Dendritic Cells in HIV-Infected Individuals¹

Jihed Chehimi,* Donald E. Campbell,[†] Livio Azzoni,* Darlene Bacheller,[†] Emmanouil Pappasavvas,* Ghassen Jerandi,* Karam Mounzer,[‡] Jay Kostman,[‡] Giorgio Trinchieri,[§] and Luis J. Montaner^{2*}

Dendritic cells (DC) have an instrumental role in the activation and function of both innate and adaptive immune responses. In humans, at least two distinct DC subsets have been characterized based on phenotypic markers: the myeloid DC (MDC) and the plasmacytoid DC (PDC). Both subsets are critical producers of cytokines (IL-12 for MDC and type I/II IFNs for PDC) and are functionally different. We show in this study that HIV⁺ individuals have a significant decrease in the number of the Lin⁻HLA-DR⁺CD123⁺ and BDCA-2⁺ PDC compared with uninfected donors ($p = 0.0001$). HIV⁺ individuals also have a sustained impairment in viral-induced IFN- α production ($p < 0.0001$). The decrease of the PDC subsets did not correlate with CD4 count or viral load and was not reversed in subjects under virally suppressive treatment, suggesting an irreversible change after infection. By contrast, the absolute number and median frequency of MDC in HIV-infected individuals were similar to those observed in uninfected controls, while a significant decrease was present in subjects with >5000 HIV-1 copies/ml. The inverse association with viral load of the MDC number, but not of IFN- α secretion or the number of PDC, suggests a role for MDC in viral control. Our data suggest that DC subsets are differentially reconstituted during the immune recovery associated with antiviral therapy. The persistent impairment of certain DC subsets may result in a sustained defect in DC-mediated innate immune functions despite an effective treatment regimen. *The Journal of Immunology*, 2002, 168: 4796–4801.

Human immunodeficiency virus infection is associated with an activation of immune responses, followed by a gradual loss of immune function and an increased susceptibility to opportunistic infections (O.I.).³ Patients with HIV infection at both early and late stages show immunoregulatory defects that precede CD4 T cell depletion (1, 2). While the loss of adaptive HIV-specific immune responses is an area of active investigation in AIDS research, the potential role of the innate immune response is largely unexplored. Of interest is the relationship between dendritic cell (DC) subsets, disease progression, viral replication, and immune reconstitution in the setting of suppressive highly active antiretroviral therapy (HAART) regimens.

Members of a distinct family of bone marrow-derived cells, DC represent <1% of total cells in peripheral blood and in lymphoid and some nonlymphoid tissues (3). In addition to participating in the regulation of adaptive responses, DC have a direct role in enhancing innate immune responses by regulating NK and macrophage functions (4). Recently, at least two distinct human DC subsets, myeloid DC (MDC) and plasmacytoid DC (PDC), have been characterized (5–8). MDC, also referred to as DC1 or APC type 1 (APC1), express CD11c and CD1c Ags and are associated with Ag uptake, T cell activation, dendritic morphology, stimulation of MLRs, and ability to secrete IL-12 in response to bacterial stimuli (9–11). PDC, also referred to as lymphoid DC, DC2, APC2, or IFN- α -producing cells, express CD123, BDCA-2, and BDCA-4 Ags and are characterized by a lymphoid morphology, modest Ag-presenting potential, and high virus-induced IFN- α secretion (7, 9, 10, 12, 13). Dysfunction and potential loss of DC subsets in HIV-1 infection could be associated with decreased T cell activation (Ag-specific or allogeneic responses) and secretion of type 1 cytokines (e.g., IL-12, IL-15, and IFN- α) that may be in part responsible for the patients' increased susceptibility to O.I. (14–16). A selective loss of IFN- α production was associated with impairment of accessory cell function in HIV infection (16, 17). A decreased number of MDC was shown in late-stage patients and in primary HIV-1 infection (18, 19), and more recently loss of PDC, defined by CD123 expression, was described in patients with high viral loads or in those with AIDS that develop O.I. or cancer (20, 21) and in primary HIV-1 infection (19). Among the limited studies that have measured effects of therapy on circulating peripheral blood DC, viral suppression following monotherapy with AZT has been associated with an increase in number of MDC and MLR

*HIV Immunopathogenesis Laboratory, Wistar Institute, and [†]Children's Hospital of Philadelphia, Division of Immunologic and Infectious Diseases, Philadelphia, PA 19104; [‡]Philadelphia Field Initiation Group for HIV Trials, Philadelphia, PA 19103; and [§]Laboratory for Immunological Research, Schering-Plough, Dardilly, France

Received for publication January 7, 2002. Accepted for publication February 22, 2002.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ A portion of this work has been presented at the National Institutes of Health Workshop on Innate Immunity (June 12–13, 2001, Gaithersburg, MD). This work was supported by the Philadelphia Foundation (Robert I. Jacobs Fund), M. Stengel-Miller, H. S. Miller, Jr., AIDS Funds from the Commonwealth of Pennsylvania, and National Institutes of Health Grants AI47760, AI44304, and AI34412.

² Address correspondence and reprint requests to Dr. Luis J. Montaner, HIV Immunopathogenesis Laboratory, Wistar Institute, 3601 Spruce Street, Room 480, Philadelphia, PA 19130. E-mail address: montaner@mail.wistar.upenn.edu

³ Abbreviations used in this paper: O.I., opportunistic infection; DC, dendritic cell; HAART, highly active antiretroviral therapy; MDC, myeloid DC; PDC, plasmacytoid DC.

activity (22), and more recently a moderate increase in PDC after HAART (19). However, it remains unknown whether DC functions may be recovered following suppressive HAART and whether the MDC and PDC subsets are differentially regulated and associated with viral control in chronically HIV-infected subjects. The recent availability of new DC markers for MDC and PDC subsets (23) allows a more accurate detection, quantitation, and characterization of DC subsets in HIV infection.

To analyze the potential alteration in DC function in circulating blood during HIV infection, we evaluated phenotypic and biological properties of two major subsets of DC from a cross-sectional cohort of HIV-infected donors and uninfected control subjects. Our data show that MDC number is inversely associated with viral load in untreated chronic HIV-1-infected patients and increases under HAART, in contrast to a sustained loss of PDC (CD123⁺ or BDCA-2⁺) subset and a concurrent decrease in IFN- α secretion.

Materials and Methods

Patients and PBMC isolation

Consenting chronically HIV-seropositive patients ($n = 65$) at various stages of disease were enrolled in this study. All patients were from the Jonathan Lax Immune Disorder Clinic (Philadelphia Field Initiation Group for HIV Trials). CD4 T cell counts range from 9 to 1,347 (mean of 501 cells/mm³), and viral load from <50 to 199,000 copies/ml (mean of 15,091 copies/ml). Twenty-one patients were not receiving therapy, and 44 were under HAART. Ninety-six percent of virally suppressed patients (viral load <50 copies/ml) were under HAART. None of the patients presented any O.I. or HIV-related neoplasms. Healthy HIV-1-seronegative donors from the Wistar Institute Blood Donor Program were included as control subjects. Institutional Review Board approval (from the Wistar Institute and Philadelphia Field Initiation Group for HIV Trials) and informed consent were obtained before blood donation. Blood was processed within 2–3 h from drawing. All reagents used were selected for their low levels of endotoxin contamination. PBMC were separated on Ficoll-Paque (Amersham Pharmacia Biotech, Uppsala, Sweden) density gradient and resuspended in RPMI 1640 medium supplemented with 10% heat-inactivated FBS, L-glutamine, and antibiotics.

Whole blood phenotypic analysis of MDC and PDC/IFN- α -producing cell subsets

A total of 200 μ l blood was lysed in ammonium chloride (Pharml; BD PharMingen, San Diego, CA) at room temperature and washed twice in FACS buffer (PBS, 2% FBS, 0.5% BSA, and 0.02% sodium azide). Cell pellets were incubated in 200 μ l FACS buffer with adequate amounts of mAb at room temperature for 20 min. The following mAbs were used: lineage mixture (Lin1-FITC), HLA-DR-PerCP, CD123-PE, and CD11c-allophycocyanin or PE, from BD Immunocytometry Systems (San Jose, CA). PE-conjugated BDCA-2 Ab was purchased from Miltenyi Biotec (Auburn, CA). Irrelevant isotype-matched (IgG1, IgG2b) mAbs were used

in each experiment. MDC were identified by Lin⁻HLA-DR⁺CD11c⁺ and PDC by Lin⁻HLA-DR⁺CD123⁺ and by the newly described mAb BDCA-2 (HLA-DR⁺BDCA-2⁺) that identify a novel human DC Ag on peripheral blood PDC (23). Cells were washed twice with FACS buffer, fixed, and analyzed in a FACSCalibur with CellQuest software (BD Immunocytometry Systems). The fluorescence of 150,000–200,000 events was accumulated for analysis. When indicated, the absolute number of circulating blood DC (MDC and PDC) was calculated using the percentage of cells with respect to the lymphocyte and monocyte absolute counts, as determined by an automated differential blood count.

IFN- α production

PBMC (2.5×10^6 /well) from controls and patients were cultured in 24-well plates with irradiated cell-free viral supernatant (5HA Influenza-PR8 or 10^5 PFU HSV-1-NS) for 18 h. As control for Influenza-PR8, allantoic fluid was used and did not induce any detectable amount of IFN- α (data not shown). Cell-free supernatants were harvested and tested by a commercial ELISA using matched Ab pair for IFN- α and performed according to the manufacturer (Endogen, Woburn, MA). Absorbance was measured on an automatic ELISA reader. Sensitivity of the assay was ~8–12 pg/ml.

Mixed lymphocyte reaction

MLR was performed as described by Young and Steinman (24). A total of 2×10^5 plastic nonadherent mononuclear cells isolated from the umbilical cord blood of healthy term neonates by Ficoll-Paque density gradient centrifugation was used as responders. Responding cells were cultured in triplicate for 5 days with 1×10^5 /well irradiated allogeneic stimulator cells in 96-well plates in 200 μ l medium. Cultures were pulsed with [³H]TdR (1 μ Ci/well; Amersham, Arlington Heights, IL) for 16 h. Nuclei were collected into fiberglass filter paper (Packard Instrument, Meriden, CT). Radioactivity in the filter paper was quantified as cpm by use of a direct beta counter 9600 (Packard Instrument). Results were expressed as stimulation index (SI) determined as follows: mean cpm in cultures containing responding and stimulator cells/mean cpm in culture containing responder cells alone. A SI of 3 and above was considered positive.

Statistical analysis

This was performed with JMP 4.0 software (SAS Institute, Cary, NC). Data were compared using paired or unpaired nonparametric tests, as appropriate. Statistical significance between HIV⁻ and HIV⁺ was performed by the Wilcoxon tests. Correlation and association between CD4 counts, viral load, and other parameters were performed by the Spearman's rank test. When needed, patients were stratified according to their CD4 T cell counts (<250, 250–500, >500), viral load (<50 copies/ml, 50–5000 copies/ml, and >5000 copies/ml), and combination antiretroviral therapy. All p values are two sided.

Results and Discussion

PDC and MDC subsets in HIV infection

We analyzed the phenotype and functions of the two major peripheral blood DC subsets (MDC and PDC) in a cross-sectional cohort of healthy uninfected controls ($n = 53$) and chronically




FIGURE 1. Representative phenotypic analysis of MDC and PDC subsets. Whole blood (200 μ l) assay is used. Multicolor staining for MDC (Lin⁻HLA-DR⁺CD11c⁺) and for PDC (Lin⁻HLA-DR⁺CD123⁺; HLA-DR⁺BDCA-2⁺) is shown. After excluding dead cells based on light scatter gating, 150,000–200,000 events were acquired. *A*, PBMC from HIV⁻ and HIV⁺ donors gated on HLA-DR^{high}, and of Lin-1⁻, DC are identified based on CD123 (PDC) or CD11c (MDC) expression. *B*, The same donors stained with HLA-DR and BDCA-2.




FIGURE 2. Phenotypic analyses of MDC and PDC subsets. Whole blood (200 μ l) assay is used, and multicolor staining is performed as described in *Materials and Methods*. Each dot represents a single donor, and horizontal bars represent the median. Statistical significance between controls and patients is shown at the top of the figure.

HIV-infected individuals ($n = 65$). Cells were quantified by multiparametric flow cytometry on whole blood (Fig. 1 shows a representative analysis). Our patient cohort consisted of HAART-treated ($n = 44$) and untreated individuals ($n = 21$), with a CD4⁺ T cell number ranging from 9 to 1,347/mm³ and a viral load from <50 to 199,000 copies/ml. The median frequency of PDC in control individuals was 0.29% (7,250 cells/ml) for CD123⁺ subset and 0.37% (8,250 cells/ml) for BDCA-2⁺ subset, whereas that of MDC was 0.27% (6,625 cells/ml). PDC were significantly ($p < 0.001$) decreased in patients (0.10%, 3,288 CD123⁺ cells/ml; 0.09%, 2,612 BDCA-2⁺ cells/ml) as compared with healthy controls (Fig. 2 and Table I). The proportions of PDC (CD123⁺ and BDCA-2⁺) were highly correlated in both control ($r = 0.55$, $p = 0.007$, $n = 21$) and HIV-infected subjects ($r = 0.64$, $p < 0.001$, $n = 33$), indicating that these two phenotypes (CD123⁺ and BDCA-2⁺) identify largely overlapping cell subsets. Unlike PDC, the median frequency of MDC in HIV-infected patients (0.27%, 6,100 cells/ml) was similar to that observed in uninfected controls (0.27%, 6,625 cells/ml; Fig. 2 and Table I). Additional DC subsets analyzed include the newly described BDCA-3⁺ subset (23), which was found to be decreased in patients as compared with uninfected controls (0.03%, 1,000 cells/ml vs 0.07%, 1,750 cells/ml, $p = 0.0034$, data not shown). However, because the nature of the BDCA-3⁺ subset is still poorly understood, further functional characterization of this subset is needed before its role in the context of an immune response can be interpreted.




FIGURE 3. IFN- α production. A total of 2.5×10^6 PBMC from HIV⁻ ($n = 36$) and HIV⁺ ($n = 48$) was exposed to irradiated Influenza-PR8 (6HA) or HSV-1-NS (10^5 PFU) for 18 h. IFN- α was measured in the cell-free supernatants by IFN- α -specific ELISA. Each dot represents a single donor, and horizontal bars represent the median. Statistical significance between controls and patients is shown.

IFN- α secretion and MLR response as functional correlates of PDC and MDC subsets

To gain further insight into the functional outcome of the decrease in circulating PDC, we evaluated the production of IFN- α by healthy uninfected controls ($n = 36$) and HIV-infected individuals ($n = 48$) in response to Influenza-PR8 or HSV-1-NS strain exposure. IFN- α has a potent antiviral and antitumor activity and, within peripheral blood, it is selectively produced by PDC following enveloped viral stimulation (7, 11, 20). Consistent with a decreased number of circulating PDC in HIV infection, PBMC from HIV-infected patients produced on average 20-fold less IFN- α than PBMC from control donors when stimulated in vitro with either Influenza-PR8 or HSV-1-NS ($p < 0.0001$, Fig. 3). Moreover, a positive correlation was found between PDC frequencies and IFN- α production by PBMC in response to either virus ($Rho = 0.8$, $p < 0.0001$ for CD123, and $Rho = 0.699$, $p < 0.0001$ for BDCA-2, respectively).

We analyzed whether in HIV-infected patients APC-specific activities associated with MDC were affected by testing the ability of PBMC to stimulate proliferation of allogeneic T cells. We compared in PBMC from 14 HIV-infected patients and in 8 healthy controls the number of both DC subsets by phenotypical analysis and their associated functional properties. As shown in Table II, even though the PDC compartment was impaired in the HIV-infected patients (as indicated by their decreased frequency and IFN- α secretion), MDC frequency and the allostimulatory function of PBMC from the patients were comparable with those of the

Table I. Characterization of DC subsets in peripheral blood of HIV-infected individuals^a

DC Subsets	HIV ⁻	HIV ⁺	<i>p</i> Values ^b
MDC	0.27 (0.17–0.47) ^c	0.27 (0.13–0.40) ^d	NS
PDC (CD123 ⁺)	0.29 (0.13–0.42) ^c	0.10 (0.05–0.18) ^d	<0.0001
PDC (BDCA-2 ⁺)	0.37 (0.15–0.50) ^e	0.09 (0.05–0.14) ^f	<0.0001

^a Values represent percentage of positive cells and are expressed as median (25th–75th percentile).

^b Statistical significance between HIV⁻ and HIV⁺.

^c $n = 53$ donors tested.

^d $n = 65$ donors tested.

^e $n = 21$ donors tested.

^f $n = 33$ donors tested.

Table II. Differential functional responses of MDC and PDC in HIV-infected individuals

Patients	MDC ^a	PDC ^a	Influenza IFN- α ^b	HSV-1 IFN- α ^b	MLR SI ^c	CD4 T Cells/mm ³	Viral Load (copies/ml)
P183	0.31	0.12	117	98	8	220	50
P187	0.3	0.11	278	158	11.6	388	50
P269	0.55	0.25	388	497	11.4	89	50
P271	0.3	0.11	202	139	11.8	622	50
P272	ND	0.19	309	201	9.6	188	50
P273	0.31	0.1	116	189	11.8	459	50
P275	0.29	0.17	375	218	10.4	598	50
P277	0.21	0.2	98	87	7.6	175	73
P270	0.14	0.12	189	209	9	283	112
P278	0.44	0.18	178	221	12.6	340	132
P185	0.26	0.09	90	ND	9.6	289	2,480
P276	0.41	0.19	312	234	10	1,050	7,150
P184	0.09	0.2	89	220	2.5	313	28,300
P186	0.22	0.19	289	401	7	600	48,600
All patients (n = 14)	0.29 \pm 0.03	0.16 \pm 0.013	217 \pm 29	221 \pm 31	9.5 \pm 0.7	401 \pm 67	6,228 \pm 3,835
All controls (n = 8)	0.30 \pm 0.035	0.32 \pm 0.03	1,459 \pm 370	1,113 \pm 306	9.22 \pm 0.6		
p values ^d	NS	p = 0.002	p < 0.0003	p < 0.0008	NS		

^a Results represent percentage of positive cells \pm SE. PDC are defined by HLA-DR⁺CD123⁺Lin⁻.

^b Viral stimulated IFN- α measured in picograms per milliliter \pm SE.

^c SI \pm SE (SI as defined in *Materials and Methods*).

^d Statistical significance between HIV⁻ and HIV⁺ was performed using the Wilcoxon tests.

controls. Moreover, no significant difference was observed in the MLR assays between virally suppressed (<50 copies/ml) and viremic patients (>50 copies/ml). Our results, indicating that MDC from HIV-infected patients are not impaired in their ability to stimulate allogeneic T cells, are in agreement with earlier reports (25, 26) and with our previous observation using enriched population of total DC (27). Interestingly, we found a positive correlation between the frequency of the MDC subset and the MLR response ($r = 0.70$, $p = 0.0073$) consistent with a role for the MDC subset in this functional response. Taken together, these results strongly suggest a dichotomy between PDC and MDC in our cohort of HIV-infected individuals: the depletion of PDC is associated with impairment of IFN- α production, while MDC are maintained and retain functional allogeneic responses.

Lack of correlation between CD4 counts, viral load, and PDC number

We compared patients grouped according to CD4 count or viral load, as described in *Materials and Methods*, to address the question of whether the PDC and MDC subsets were affected by CD4 T cell counts, viral load, or antiretroviral therapy. Statistical analysis revealed no correlation between CD4 T cell counts or viral load and PDC frequency (defined by expression of CD123⁺ or BDCA-2⁺), albeit their number in the patients was significantly decreased in comparison with healthy uninfected controls. Importantly, analysis of PDC in virally suppressed patients as compared with uninfected controls revealed a sustained decrease in both cell frequency and IFN- α secretion (Tables II and III), indicating that

the loss of PDC is not reversed by antiretroviral treatment. IFN- α production was significantly associated with CD4 counts only when the analysis was restricted to viremic patients with >50 HIV copies/ml (data not shown). In contrast to a recent report in which increased levels of IFN- α secretion were described following HAART (28), impaired IFN- α production was observed in treated patients with complete viral suppression (<50 copies/ml) as well as in patients with persistent viral replication (>50 copies/ml, $p < 0.0001$ when compared with controls; Table III). The differences in the results reported in this study and the one mentioned above (28) could depend on specific characteristics of the patient cohorts studied: e.g., none of the patients used in this study had a history of O.I. or oncological diseases. It should also be noted that our interpretation of a sustained decrease in IFN- α secretion (by ELISA) rests on a direct comparison with uninfected controls, while the above mentioned study addressed recovery based on investigator-assigned thresholds of IFN- α secretion (by bioassay) within specific HIV⁺ populations (28). Taken together, our results suggest that a loss of PDC may be an early event in HIV infection with a limited capacity for reconstitution following antiretroviral therapy, as reflected by persistent low frequency of PDC in subjects under treatment with <50 HIV copies/ml and high CD4 T cell count.

Although the exact mechanism responsible for the PDC deficiency remains to be defined, our observations argue against a predominant role of PDC in the control of HIV replication in chronic infection, as proposed by others (20, 21, 28). Of interest, we noticed that six of the HIV-infected subjects on HAART had PDC frequencies in the normal range ($\geq 0.29\%$) and only two of these individuals, with CD4 counts

Table III. DC subsets and viral load in HIV-infected individuals^a

DC Subsets/IFN- α	HIV (<50 copies/ml)	HIV (50–5000 copies/ml)	HIV (>5000 copies/ml)	HIV ⁻
MDC	0.31 (0.28–0.46), n = 25	0.26 (0.14–0.67), n = 15	0.13 (0.10–0.22), n = 24 ^b	0.27 (0.17–0.47), n = 53
PDC (CD123 ⁺)	0.11 (0.08–0.18), n = 26	0.09 (0.03–0.17), n = 15	0.09 (0.04–0.19), n = 24, NS	0.29 (0.13–0.42), n = 53
PDC (BDCA-2 ⁺)	0.08 (0.02–0.18), n = 12	0.11 (0.07–0.17), n = 9	0.09 (0.06–0.13), n = 12, NS	0.37 (0.15–0.50), n = 21
Influenza-IFN- α	107 (49–325), n = 18	80 (26–180), n = 14	93 (36–189), n = 16, NS	1468 (626–2094), n = 34
HSV-1-IFN- α	164 (57–233), n = 18	54 (10–179), n = 14	84 (33–173), n = 16, NS	912 (672–1790), n = 35
CD4 counts	584 (279–694)	378 (312–607)	425 (271–600)	

^a Results shown are measured in percentage of positive cells and picograms per milliliter for IFN- α , and values are expressed as median (25th–75th percentile).

^b Statistical significance within the MDC subset is as follows: <50 copies/ml vs >5000 copies/ml, $p < 0.0032$, and 50–5000 copies/ml vs >5000 copies/ml, $p = 0.029$, respectively.

of 1246 and 651 cells/mm³, had normal IFN- α responses. These data suggest that retention of the PDC subset may be possible in a small subset of patients highly responsive to antiretroviral therapy, as it was previously described for the long-term nonprogressor patients (20). Additional experiments isolating the PDC will be needed to determine whether a decrease in IFN- α secretion is due to a decrease in frequency of PDC or whether a functional impairment of this subset is also present.

Our data confirm a recent report by Feldman et al. (29) and support the association in HIV-infected patients between a decrease in IFN- α secretion and a decrease in the CD123⁺ DC subset, and extend it with the use of additional markers such as BDCA-2.

CD4 counts, viral load, and frequency of MDC

The overall frequency of MDC in patients remained stable with similar cell percentage as compared with uninfected controls (Fig. 2 and Table I). No correlation was found between MDC frequency and CD4⁺ T cell counts (data not shown). However, virally suppressed patients were found to have a significantly higher number of MDC (0.31%; 7020 cells/ml) than patients with a viral load of >5000 copies/ml (0.13%; 2636 cells/ml, $p < 0.0032$; Table III). An association between a decrease of MDC frequency with an increase of viral load was further indicated by a significant negative correlation between these variables ($Rho = -0.5$, $p < 0.0001$, $n = 65$). The inverse association between MDC and viral load was also sustained when the analysis was restricted to a subset of 21 patients not receiving treatment ($Rho = -0.68$, $p = 0.0007$). It is important to note that the extensive use of antiviral treatment in our cohort may have biased our study against the possibility to detect the previously reported decreases in MDC during HIV infection (18), specifically the prevalence of HIV-infected subjects with <5000 copies/ml that did not show a significant decrease in MDC frequency compared with healthy controls. Overall, these results are consistent with a decreased MDC frequency only at high viral loads that can be reversed following antiretroviral treatment in contrast to the early and persistent deficiency in the PDC subset measured in the same patients, as described above. A longitudinal analysis of the consequences of HAART-mediated suppression on each DC subset (i.e., recovery of number and function as a correlate of viral suppression) is needed to confirm our cross-sectional findings.

We show for the first time a differential depletion between the frequency and function of MDC and PDC subsets when analyzed in parallel under suppressive therapy. Our data complement previous observations describing the depletion of MDC and PDC in viremic or end-stage HIV-infected patients (20, 21, 28). However, our studies indicate a lack of full DC reconstitution upon HAART that may contribute to a sustained immune impairment in these subjects. While we document a phenotypic and functional preservation in the MDC subset in the virally suppressed patients and HAART-responding patients, alteration in the PDC subset appears to be an early event and associated with a sustained impairment of IFN- α production. The consequence of the lack of recovery of IFN- α production and PDC number on the innate and adaptive immune functions of the patients remains to be defined. For example, a decrease in viral-induced IFN- α secretion may also affect in part the immune reconstitution of NK cell function. Additional experiments should assess whether the loss of this subset from circulation reflects a cytopathic effect caused by the virus through direct infection of the PDC subset (30) or through viral-induced apoptosis, as described with measles virus infection (31, 32). A role for interactions between HIV-1 and PDC subsets is supported by the susceptibility of IL-3-cultured Lin⁻HLA-DR⁺CD11c⁻ cells to

be infected in vitro by HIV-1 (30). It is also of interest to relate our findings to current treatment guidelines for HIV infection, as a delay in treatment of patients with viral loads above 5000 copies/ml, irrespective of CD4 T cell count, would not prevent a decrease of MDC and PDC that may bear on immune function in general.

Acknowledgments

We are grateful to the patient population from the Jonathan Lax Immune Disorder Clinic (Philadelphia Field Initiation Group for HIV Trials). We thank Dr. Harvey Friedman (Division of Infectious Diseases, University of Pennsylvania, Philadelphia, PA) for providing HSV strain NS. We thank Deborah Davis (Wistar Blood Donor Program), Serena Black, Kim Hart, Jane Shull, and the clinical staff at Philadelphia Field Initiation Group for HIV Trials for their assistance in providing blood samples.

References

- Levy, J. A. 1993. Pathogenesis of human immunodeficiency virus infection. *Microbiol. Rev.* 57:183.
- Fauci, A. S. 1993. Immunopathogenesis of HIV infection. *J. Acquired Immune Defic. Syndr.* 6:655.
- Steinman, R. M., G. Kaplan, M. D. Witmer, and Z. A. Cohn. 1979. Identification of a novel cell type in peripheral lymphoid organs of mice. V. Purification of spleen dendritic cells, new surface markers, and maintenance in vitro. *J. Exp. Med.* 149:1.
- Banchereau, J., and R. M. Steinman. 1998. Dendritic cells and the control of immunity. *Nature* 392:245.
- Steinman, R. M., and K. Inaba. 1999. Myeloid dendritic cells. *J. Leukocyte Biol.* 66:205.
- Banchereau, J., B. Pulendran, R. Steinman, and K. Palucka. 2000. Will the making of plasmacytoid dendritic cells in vitro help unravel their mysteries? *J. Exp. Med.* 192:F39.
- Banchereau, J., F. Briere, C. Caux, J. Davoust, S. Lebecque, Y. J. Liu, B. Pulendran, and K. Palucka. 2000. Immunobiology of dendritic cells. *Annu. Rev. Immunol.* 18:767.
- Liu, Y. J., H. Kanzler, V. Soumelis, and M. Gilliet. 2001. Dendritic cell lineage, plasticity and cross-regulation. *Nat. Immun.* 2:585.
- Chehimi, J., S. E. Starr, H. Kawashima, D. S. Miller, G. Trinchieri, B. Perussia, and S. Bandyopadhyay. 1989. Dendritic cells and IFN- α -producing cells are two functionally distinct non-B, non-monocytic HLA-DR⁺ cell subsets in human peripheral blood. *Immunology* 68:488.
- Starr, S. E., S. Bandyopadhyay, V. Shanmugam, N. Hassan, S. Douglas, S. J. Jackson, G. Trinchieri, and J. Chehimi. 1993. Morphological and functional differences between HLA-DR⁺ peripheral blood dendritic cells and HLA-DR⁺ IFN- α producing cells. *Adv. Exp. Med. Biol.* 329:173.
- Sallusto, F., and A. Lanzavecchia. 2000. Understanding dendritic cell and T-lymphocyte traffic through the analysis of chemokine receptor expression. *Immunol. Rev.* 177:134.
- Perussia, B., V. Fanning, and G. Trinchieri. 1985. A leukocyte subset bearing HLA-DR antigens is responsible for in vitro α interferon production in response to viruses. *Nat. Immun. Cell Growth Regul.* 4:120.
- Siegal, F. P., N. Kadowaki, M. Shodell, P. A. Fitzgerald-Bocarsly, K. Shah, S. Ho, S. Antonenko, and Y. J. Liu. 1999. The nature of the principal type 1 interferon-producing cells in human blood. *Science* 284:1835.
- Lopez, C., P. A. Fitzgerald, and F. P. Siegal. 1983. Severe acquired immune deficiency syndrome in male homosexuals: diminished capacity to make interferon- α in vitro associated with severe opportunistic infections. *J. Infect. Dis.* 148:962.
- Lopez, C., P. A. Fitzgerald, F. P. Siegal, S. Landesman, J. Gold, and S. E. Krown. 1984. Deficiency of interferon- α generating capacity is associated with susceptibility to opportunistic infections in patients with AIDS. *Ann. NY Acad. Sci.* 437:39.
- Siegal, F. P., C. Lopez, P. A. Fitzgerald, K. Shah, P. Baron, I. Z. Leiderman, D. Imperato, and S. Landesman. 1986. Opportunistic infections in acquired immune deficiency syndrome result from synergistic defects of both the natural and adaptive components of cellular immunity. *J. Clin. Invest.* 78:115.
- Ferbas, J., J. Navratil, A. Logar, and C. Rinaldo. 1995. Selective decrease in human immunodeficiency virus type 1 (HIV-1)-induced α interferon production by peripheral blood mononuclear cells during HIV-1 infection. *Clin. Diagn. Lab. Immunol.* 2:138.
- Grassi, F., A. Hosmalin, D. McIlroy, V. Calvez, P. Debre, and B. Autran. 1999. Depletion in blood CD11c-positive dendritic cells from HIV-infected patients. *AIDS* 13:759.
- Pacanowski, J., S. Kahi, M. Baillet, P. Lebon, C. Deveau, C. Goujard, L. Meyer, E. Oksenhendler, M. Sinet, and A. Hosmalin. 2001. Reduced blood CD123⁺ (lymphoid) and CD11c⁺ (myeloid) dendritic cell numbers in primary HIV-1 infection. *Blood* 98:3016.
- Soumelis, V., I. Scott, F. Gheyas, D. Bouhour, G. Cozon, L. Cotte, L. Huang, J. A. Levy, and Y. J. Liu. 2001. Depletion of circulating natural type 1 interferon-producing cells in HIV-infected AIDS patients. *Blood* 98:906.
- Donaghy, H., A. Pozniak, B. Gazzard, N. Qazi, J. Gilmour, F. Gotch, and S. Patterson. 2001. Loss of blood CD11c⁺ myeloid and CD11c⁻ plasmacytoid dendritic cells in patients with HIV-1 infection correlates with HIV-1 RNA virus load. *Blood* 98:2574.

22. Gompels, M., S. Patterson, M. S. Roberts, S. E. Macatonia, A. J. Pinching, and S. C. Knight. 1998. Increase in dendritic cell numbers, their function and the proportion uninfected during AZT therapy. *Clin. Exp. Immunol.* 112:347.
23. Dzionek, A., A. Fuchs, P. Schmidt, S. Cremer, M. Zysk, S. Miltenyi, D. W. Buck, and J. Schmitz. 2000. BDCA-2, BDCA-3, and BDCA-4: three markers for distinct subsets of dendritic cells in human peripheral blood. *J. Immunol.* 165:6037.
24. Young, J. W., and R. M. Steinman. 1988. Accessory cell requirements for the mixed-leukocyte reaction and polyclonal mitogens, as studied with a new technique for enriching blood dendritic cells. *Cell. Immunol.* 111:167.
25. Cameron, P. U., U. Forsum, H. Tepler, A. Granelli-Piperno, and R. M. Steinman. 1992. During HIV-1 infection most blood dendritic cells are not productively infected and can induce allogeneic CD4⁺ T cells clonal expansion. *Clin. Exp. Immunol.* 88:226.
26. Cameron, P., M. Pope, A. Granelli-Piperno, and R. M. Steinman. 1996. Dendritic cells and the replication of HIV-1. *J. Leukocyte Biol.* 59:158.
27. Chehimi, J., W. Z. Ho, and S. E. Starr. 1996. HIV infection of human peripheral blood, cord blood and tonsillar dendritic cells. In *Perspectives in Drug Discovery and Design*, Vol. 5. P. S. Anderson, G. L. Kenyon, and G. R. Marshall, eds. ESCOM, Leiden, The Netherlands, p. 103.
28. Siegal, F. P., P. Fitzgerald-Bocarsly, B. K. Holland, and M. Shodell. 2001. Interferon- α generation and immune reconstitution during antiretroviral therapy for human immunodeficiency virus infection. *AIDS* 15:1603.
29. Feldman, S., D. Stein, S. Amrute, T. Denny, Z. Garcia, P. Kloser, Y. Sun, N. Megjugorac, and P. Fitzgerald-Bocarsly. 2001. Decreased interferon- α production in HIV-infected patients correlates with numerical and functional deficiencies in circulating type 2 dendritic cell precursors. *Clin. Immunol.* 101:201.
30. Patterson, S., A. Rae, N. Hockey, J. Gilmour, and F. Gotch. 2001. Plasmacytoid dendritic cells are highly susceptible to human immunodeficiency virus type 1 infection and release infectious virus. *J. Virol.* 75:6710.
31. Fugier-Vivier, I., C. Servet-Delprat, P. Rivallier, M. C. Rissoan, Y. J. Liu, and C. Rabourdin-Combe. 1997. Measles virus suppresses cell-mediated immunity by interfering with the survival and functions of dendritic and T cells. *J. Exp. Med.* 186:813.
32. Servet-Delprat, C., P. O. Vidalain, O. Azocar, F. Le Deist, A. Fischer, and C. Rabourdin-Combe. 2000. Consequences of Fas-mediated human dendritic cell apoptosis induced by measles virus. *J. Virol.* 74:4387.