

Protein Detection by Sequencing

Uncover What Makes Each Cell Unique With Antibodies, Panels, and Multiomics Software



BioLegend®

Learn More



Lipopolysaccharide (LPS)-Binding Protein Mediates LPS Detoxification by Chylomicrons

This information is current as of April 17, 2021.

Anita C. E. Vreugdenhil, Corine H. Rousseau, Thomas Hartung, Jan Willem M. Greve, Cornelis van 't Veer and Wim A. Buurman

J Immunol 2003; 170:1399-1405; ;
doi: 10.4049/jimmunol.170.3.1399
<http://www.jimmunol.org/content/170/3/1399>

References This article **cites 41 articles**, 16 of which you can access for free at:
<http://www.jimmunol.org/content/170/3/1399.full#ref-list-1>

Why *The JI*? [Submit online.](#)

- **Rapid Reviews! 30 days*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

**average*

Subscription Information about subscribing to *The Journal of Immunology* is online at:
<http://jimmunol.org/subscription>

Permissions Submit copyright permission requests at:
<http://www.aai.org/About/Publications/JI/copyright.html>

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://jimmunol.org/alerts>

The Journal of Immunology is published twice each month by
The American Association of Immunologists, Inc.,
1451 Rockville Pike, Suite 650, Rockville, MD 20852
Copyright © 2003 by The American Association of
Immunologists All rights reserved.
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



Lipopolysaccharide (LPS)-Binding Protein Mediates LPS Detoxification by Chylomicrons¹

Anita C. E. Vreugdenhil,* Corine H. Rousseau,* Thomas Hartung,[†] Jan Willem M. Greve,* Cornelis van 't Veer,* and Wim A. Buurman^{2*}

Chylomicrons have been shown to protect against endotoxin-induced lethality. LPS-binding protein (LBP) is involved in the inactivation of bacterial toxin by lipoproteins. The current study examined the interaction among LBP, chylomicrons, and bacterial toxin. LBP was demonstrated to associate with chylomicrons and enhance the amount of LPS binding to chylomicrons in a dose-dependent fashion. In addition, LBP accelerated LPS binding to chylomicrons. This LBP-induced interaction of LPS with chylomicrons prevented endotoxin toxicity, as demonstrated by reduced cytokine secretion by PBMC. When postprandial circulating concentrations of chylomicrons were compared with circulating levels of low density lipoprotein, very low density lipoprotein, and high density lipoprotein, chylomicrons exceeded the other lipoproteins in LPS-inactivating capacity. Furthermore, highly purified lipoteichoic acid, an immunostimulatory component of Gram-positive bacteria, was detoxified by incubation with LBP and chylomicrons. In conclusion, our results indicate that LBP associates with chylomicrons and enables chylomicrons to rapidly bind bacterial toxin, thereby preventing cell activation. Besides a role in the detoxification of bacterial toxin present in the circulation, we believe that LBP-chylomicron complexes may be part of a local defense mechanism of the intestine against translocated bacterial toxin. *The Journal of Immunology*, 2003, 170: 1399–1405.

Besides a crucial role in the transport of dietary lipids, chylomicrons appear to reinforce the host defense against LPS, a major constituent of the outer membrane of Gram-negative bacteria (1, 2). LPS provokes an inflammatory response in the infected host by virtue of its ability to bind membrane factors present on the surface of macrophages and other cells essential to host defense. Once the response to Gram-negative bacteria is initiated, there is a need to inactivate and eliminate LPS to minimize potential deleterious effects of an unregulated response. Circulating chylomicrons were previously demonstrated to be potent inhibitors of LPS activity in vivo and to prevent endotoxin-induced mortality (1, 2). The metabolic fate of LPS was shown to be strongly affected by complex formation with chylomicrons. Whereas the association of LPS with high density lipoprotein (HDL)³ decreases its clearance from the circulation (3, 4), chylomicrons enhance the plasma clearance and hepatic uptake of LPS, with shunting of LPS to hepatocytes and away from hepatic macrophages (2, 5). Harris et al. (6) recently presented data indicating that both the low density lipoprotein (LDL) receptor and LDL receptor-related protein participate in the rapid internalization of

chylomicron-bound LPS by hepatocytes. Consequently, LPS is delivered into the bile, where it is inactivated by bile salts (7). Lipoproteins other than chylomicrons were previously found to prevent a cellular response by binding of LPS (8), and recent studies by Kitchens et al. (9) demonstrated that LPS already bound to monocytes is released in response to lipoproteins, thereby reducing cytokine secretion. Besides an enhanced clearance of LPS from the circulation, such mechanisms may also be applicable to chylomicrons and may account for the chylomicron-induced increase in survival during endotoxemia. It was demonstrated that plasma constituents other than lipoproteins are required for maximal binding of LPS to lipoproteins (10) and for lipoproteins to achieve maximal prevention of endotoxin-induced death (1), suggesting that a constituent of plasma is required for the binding of endotoxin to lipoproteins.

The acute phase protein, LPS-binding protein (LBP), plays an important role in the reduction of LPS activity by catalyzing the transfer of monomerized LPS from micelles into circulating lipoproteins (11). The enhanced concentration of circulating LBP during an acute phase response was recently demonstrated to be crucial, since it diminishes the transfer of LPS to monocytes and reduces cytokine secretion (12). LBP also induces lipoprotein inactivation of lipoteichoic acid (LTA), a component of most Gram-positive bacteria (13). Although initial studies focused on the LPS-detoxifying function of HDL (14), later studies demonstrated that in serum LPS predominantly binds to very low density lipoprotein (VLDL) and LDL, which also results in the inactivation of LPS (15–17). We recently demonstrated that LBP circulates in complex with apolipoprotein B (apoB)-containing lipoproteins, i.e., LDL and VLDL (18). LBP complexed with LDL and VLDL strongly enhances LPS binding to these lipoproteins, which suggests an important role for LBP/lipoprotein complexes in the scavenging of endotoxin present in the circulation (18). The association of LBP with LDL and VLDL appears to result in part from the high affinity of LBP for apoB. This prompted us to study whether LBP also cooperates

*Nutrition and Toxicology Research Institute Maastricht, Department of General Surgery, Maastricht University, Maastricht, The Netherlands; and [†]Department of Biochemical Pharmacology, University of Konstanz, Konstanz, Germany

Received for publication April 11, 2002. Accepted for publication November 15, 2002.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This work was supported by a grant from the Dutch Digestive Disease Foundation (Nieuwegein, The Netherlands) and Numico Research B.V. (Wageningen, The Netherlands).

² Address correspondence and reprint requests to Dr. Wim A. Buurman, Department of General Surgery, Maastricht University, P.O. Box 616, 6200 MD Maastricht, The Netherlands. E-mail address: w.buurman@ah.unimaas.nl

³ Abbreviations used in this paper: HDL, high density lipoprotein; apo, apolipoprotein; LBP, LPS-binding protein; LDL, low density lipoprotein; LTA, lipoteichoic acid; pAb, polyclonal Ab; TMB, tetramethylbenzidine; VLDL, very low density lipoprotein.

with the apoB-containing chylomicrons in detoxification of the bacterial toxins LPS and LTA. Consequently, we studied the interaction of LBP with chylomicrons and the effect of this interaction on toxin binding and activity.

Materials and Methods

Reagents

LPS from *Escherichia coli*, serotype 055:B5, was purchased from Sigma-Aldrich (St. Louis, MO). Polyclonal Abs (pAb) to human LBP were obtained by immunizing rabbits with purified human LBP. Protein A-purified anti-LBP IgG was biotinylated following standard procedures. Anti-human LBP mAb HM14 was obtained by immunizing mice with LBP following classical procedures (18). BSA (containing <1 ng LPS/mg) was derived from Sigma-Aldrich. Human rTNF- α was provided by BASF/Knoll Antigens (Ludwigshafen, Germany). Since LTA preparations available to date are highly contaminated by LPS (19), we used a recently introduced preparation method for LTA (20) that allows study of the effects of LPS-free, biologically active LTA. LTA, 99% pure and containing <30 pg of LPS/mg LTA, was prepared from *Staphylococcus aureus*.

Blood samples

Human venous blood was collected from healthy volunteers after an overnight fast and 2.5 h postprandially. To prepare serum, blood was allowed to clot for 1 h at room temperature. Fresh serum was used for isolation of lipoproteins.

Lipoprotein isolation

Chylomicrons were isolated from pooled postprandial human serum (six donors) by ultracentrifugation according to the procedure described by Terpstra et al. (21). Briefly, dry potassium bromide (770 mg) and sucrose (50 mg) together with 2 ml of serum were placed in the bottoms of ultra clear centrifuge tubes (Beckman Instruments, Brea, CA). These components were carefully mixed with a spatula. A discontinuous gradient was made with the serum mixture at a density of 1.250 g/ml, two potassium bromide solutions with densities of 1.225 and 1.100 g/ml, respectively, and, on top, a water density of 0.998 g/ml. The tubes were placed in an SW41 rotor and centrifuged in a Beckman XL-80 ultracentrifuge (Beckman Instruments) at 30,000 rpm and 20°C for 40 min. The chylomicrons were collected by aspiration of the top 1-ml fraction and were stored at -80°C until use. HDL, LDL, and VLDL were isolated from pooled fresh human sera (six healthy donors) by a 22-h single spin density gradient ultracentrifugation at 40,000 rpm at 17°C according to the method reported by Terpstra et al. (22) using a Beckman XL-80 ultracentrifuge with an SW40 rotor. A discontinuous gradient was constructed from 2 ml of serum adjusted to a density of 1.250 g/ml with potassium bromide and sucrose, two NaCl/potassium bromide solutions with densities of 1.225 and 1.100 g/ml, respectively, and a water density of 0.998 g/ml. All solutions contained 0.1 mg/ml EDTA. The VLDL, LDL, and HDL fractions were collected by aspiration, dialyzed extensively against PBS at 4°C, and stored at -80°C until use. The lipoprotein fractions used contained <4 ng/ml of LBP, as assessed by LBP-specific ELISA (23).

Purification of LBP

LBP was isolated from human plasma by selective affinity immunosorption as described previously (18). In short, anti-hLBP mAb HM-14 was cross-linked to cyanogen bromide-activated Sepharose (Amersham Pharmacia Biotech, Piscataway, NJ) according to the manufacturer's instructions. Normal human plasma provided by the local blood bank was applied to the anti-LBP column, and unbound proteins were washed out with 0.5 M MgCl. Bound LBP was eluted with 0.1 M glycine-HCl buffer (pH 2.5).

Preparation of biotin-labeled LBP and LPS

LPS and LBP were biotinylated using hydrazide-biotin (Pierce, Rockford, IL) as described previously (18). Briefly, LBP and LPS were oxidized with sodium periodate to generate aldehydes from vicinal diols present on saccharide units. The products were then made to react with the hydrazide functionalities present on the biotin, forming stable hydrazone linkages. Linking biotin to the saccharide units in the LBP glycoprotein using hydrazide-biotin did not alter the biologic activity of LBP (data not shown).

Assays for evaluating the association of LBP and LPS to chylomicrons

To study the binding of LBP and LPS to chylomicrons, an in vitro assay was used. Ninety-six-well plates were coated with chylomicrons for 2 h at

37°C. Nonspecific binding sites were blocked by 2-h incubation with 5% BSA in PBS. Plates were washed with PBS. A concentration range of biotin-labeled LBP diluted in 0.1% BSA in PBS was incubated with the immobilized chylomicrons for 1.5 h at 37°C. In control experiments 10 μ g/ml of anti-human LBP pAb or aspecific pAb was added together with LBP. Bound LBP was detected using peroxidase-conjugated streptavidin and tetramethylbenzidine (TMB; Kirkegaard & Perry Laboratories, Gaithersburg, MD).

The effect of LBP association with chylomicrons on the capacity of these lipoproteins to interact with LPS was evaluated. To this end, chylomicrons were coated to plates, and nonspecific binding sites were blocked with 5% BSA in PBS. A concentration range of LBP was incubated for 1.5 h and was allowed to associate with the chylomicrons at 37°C. Unbound LBP was washed away, and biotin-labeled LPS was incubated with the chylomicron-LBP complexes for 1 h at 37°C. In control experiments 10 μ g/ml of anti-human LBP pAb was incubated for 1 h at 37°C with the LBP-chylomicron complexes, and biotin-labeled LPS was added together with anti-LBP Ab. After washing the plates, bound LPS was detected using streptavidin-peroxidase and TMB. The LPS binding capacity of the LBP-chylomicron complexes was compared with that of chylomicrons free of LBP.

Labeling of LPS with BODIPY and measurement of fluorescence

LPS from *E. coli* serotype 055:B5 was labeled with the fluorophore BODIPY FL SE (Molecular Probes, Eugene, OR) according to the manufacturer's instructions. Fluorescent-labeled LPS can be used to study the binding and transfer of LPS in real time (24). Time-dependent changes in fluorescence caused by the addition of LBP and chylomicrons were measured with a Spectra Max Gemini XS microplate fluorometer (Molecular Devices, Sunnyvale, CA), with emission at 518 nm and excitation at 485 nm.

Cell isolation and stimulation

For experiments designed to determine the effect of chylomicron binding of toxin derived from Gram-negative bacteria as well as from Gram-positive bacteria on their biologic activity, cytokine induction by toxin incubated with chylomicrons in the presence or the absence of LBP was assessed. PBMC were isolated from buffy coats of healthy volunteers, provided by the local blood bank, by buoyant density centrifugation on Lymphoprep (Nycomed, Oslo, Norway). Cells were washed five times in HBSS to remove LBP and other plasma proteins and then were suspended in macrophage serum-free medium (Life Technologies, Paisley, U.K.) supplemented with 100 U/ml penicillin and 100 μ g/ml streptomycin (Life Technologies). LPS and LTA were preincubated with LBP, LBP and lipoproteins, or lipoproteins alone as indicated in 96-well, flat-bottom tissue culture plates (Costar, Cambridge, MA) for 24 or 0.5 h at 37°C. Cells were added immediately after isolation at 10⁵ cells/well. Control cultures were incubated with medium alone. The cell cultures were incubated for 4 h at 37°C in 5% CO₂. Supernatants were collected and stored at -20°C until use.

TNF- α and IL-8 ELISA

Cytokine concentrations in the cell culture supernatants were determined using sandwich ELISA for TNF- α (25) and IL-8 (26). In short, 96-well Immunomaxisorp plates (Nunc, Roskilde, Denmark) were coated overnight at 4°C with cytokine-specific mAb. TNF- α in the supernatants was detected with polyclonal rabbit anti-human TNF- α , followed by peroxidase-conjugated, goat anti-rabbit IgG (Jackson ImmunoResearch Laboratories, West Grove, PA). IL-8 was detected by biotinylated polyclonal rabbit anti-human IL-8, followed by peroxidase-conjugated streptavidin. TMB was used as a substrate for peroxidase. Photospectrometry was performed at 450 nm.

Results

LBP associates with chylomicrons

Recently, LBP was shown to be predominantly complexed with apoB-containing lipoprotein particles, i.e., LDL and VLDL. This interaction was found to be at least in part mediated by an interaction of LBP with apoB (18). These findings prompted us to investigate the interaction of LBP with another triacylglyceride-rich apoB-containing lipoprotein, i.e., chylomicrons. To study the binding of LBP to isolated chylomicrons, an in vitro assay was

used that revealed a concentration-dependent binding of biotinylated LBP to chylomicrons (Fig. 1). Binding of LBP to chylomicrons was completely blocked by a pAb against LBP (Fig. 1), indicating the specificity of this interaction. Addition of aspecific pAb did not affect the results. The data strongly suggest that LBP and chylomicrons form a complex.

LBP associated with chylomicrons enhances the LPS binding capacity of chylomicrons

LBP was demonstrated to transfer LPS into lipoproteins, thereby enhancing its detoxification (11). To elucidate whether LBP complexed to chylomicrons is able to enhance LPS binding to chylomicrons, an *in vitro* assay was used. LPS binding to chylomicrons was compared with binding of LPS to LBP-chylomicron complexes. LPS was observed to interact with chylomicrons in the absence of LBP (Fig. 2A). However, LBP associated with chylomicrons strongly enhanced the LPS binding capacity of chylomicrons (Fig. 2A). The LBP-induced up-regulation of LPS binding to chylomicrons was dose dependent. This effect of LBP was blocked by pAb against LBP (Fig. 2B). LPS binding to chylomicrons in the absence of LBP was not affected by the presence of pAb against LBP, which excludes that this binding was caused by small amounts of LBP present in the chylomicron fractions. Overall, the data imply that LBP associated with chylomicrons was functionally active in transferring LPS to chylomicrons.

LBP accelerates the transfer of BODIPY-LPS into chylomicrons

To confirm the ability of LBP to mediate transfer of LPS to chylomicrons and to monitor the time kinetics of this interaction, we used BODIPY-LPS donor micelles and chylomicrons as acceptor particles. Addition of chylomicrons to BODIPY-LPS in the absence of LBP resulted in a dequenching of the fluorescence, which is indicative for dissociation of LPS from the micelles and binding to chylomicrons. Incubation with LBP together with chylomicrons resulted in a pronounced enhancement of the rate of dequenching of BODIPY-LPS (Fig. 3). This acceleration of LPS binding to




FIGURE 1. LBP associates with chylomicrons. Chylomicrons isolated from postprandial human serum by ultracentrifugation were immobilized to 96-well plates and incubated with biotinylated LBP (●). As a control a pAb against human LBP was added together with LBP (■). Bound LBP was detected by peroxidase-conjugated streptavidin and TMB. Binding of LBP to the lipoproteins was expressed as the mean \pm SD of the OD₄₅₀ of four wells of a representative experiment repeated twice after correction for background binding of biotin-labeled LBP to coated plates in the absence of chylomicrons. A dose-dependent association of LBP with chylomicrons was observed.




FIGURE 2. LBP associated with chylomicrons enhances the interaction with LPS. *A*, Immobilized chylomicrons were preincubated with a concentration range of LBP overnight at 37°C; unbound LBP was removed by washing the plates, followed by addition of biotinylated LPS to the LBP-chylomicron complexes. Bound LPS was detected using peroxidase-conjugated streptavidin. Binding of LPS to chylomicrons is expressed as the mean \pm SD of the OD₄₅₀ of four wells of a representative experiment repeated twice after correction for background binding of biotin-labeled LPS to coated plates in the absence of chylomicrons. LBP complexed to the chylomicrons enhanced LPS binding dose-dependently. *B*, Immobilized chylomicrons were preincubated either with or without 1 μ g/ml overnight at 37°C (□). Unbound LBP was removed. In controls, 10 μ g/ml of anti-human LBP pAb was added for 1 h (▨). Biotinylated LPS was added to the LBP-chylomicron complexes. Binding of LPS was assessed, and values are expressed as described in *A*. The anti-human LBP pAb completely blocked LBP-mediated binding of LPS to chylomicrons.

chylomicrons was LBP dose dependent. These findings are in line with our observations described above and indicate that LBP catalyzes LPS binding to chylomicrons and, more importantly, accelerates LPS binding to chylomicrons.

Chylomicrons detoxify LPS in a LBP dose-dependent fashion

Next, we studied whether the observed enhanced LPS binding to chylomicrons induced by LBP results in abrogation of the bioactivity of LPS. Therefore, we exposed PBMC to LPS preincubated with chylomicrons and different concentrations of LBP. TNF- α and IL-8 release by PBMC after a 4-h incubation period were measured. Fig. 4 shows that preincubation of LPS with chylomicrons and LBP clearly reduced the LPS-evoked TNF- α and IL-8




FIGURE 3. LBP accelerates the binding of LPS to chylomicrons. BODIPY-LPS was incubated with PBS alone (curve A), with 1 and 10 μg of LBP/ml (curves B and C), with chylomicrons (curve D), with chylomicrons and 1 μg of LBP/ml (curve E), and with chylomicrons and 10 μg of LBP/ml (curve F). The fluorescence at 518 nm was recorded over time. The figure is representative of two separate experiments. LBP was found to dose-dependently accelerate the binding of LPS to chylomicrons.

secretion by PBMC. This LPS-detoxifying capacity of chylomicrons was found to depend on the concentration of LBP present. In the absence of chylomicrons, LBP enhanced the LPS-induced cell activation dose-dependently, as expected. In the presence of increasing LBP concentrations (100 ng/ml, 1.7 nM), however, this effect of LBP declined, as described by others (27). A high concentration of LPS (100 ng/ml, 40 nM) was used in these experiments to allow stimulation of cells in the absence of LBP. In addition, pilot experiments showed that the capacity to induce cytokine secretion by 10 ng/ml LPS was completely abrogated in the presence of chylomicrons and 5 ng/ml LBP (data not shown).

To study the effect of the chylomicron concentration on inhibition of LPS-induced cytokine secretion, 100 ng/ml of LPS was preincubated with 100 ng/ml of LBP and different concentrations of chylomicrons (Fig. 5). The LPS-detoxifying function of chylomicrons was concentration dependent. Moreover, 50% of the postprandial circulating level of chylomicrons abrogated the cytokine response induced by 100 ng/ml of LPS, while a profound reduction of LPS activity was also seen at low chylomicron concentrations.

Prior studies described a maximal endotoxin inactivation by lipoproteins after 24 h of preincubation (17). However, using BODIPY-LPS dequenching experiments we found a rapid binding of LPS to chylomicrons in the presence of LBP (Fig. 3). To study whether this rapid LBP-mediated binding of LPS to chylomicrons results in detoxification of LPS, chylomicrons were preincubated with LBP and LPS only for 0.5 h, and PBMC were added. Fig. 6 demonstrates that in the presence of LBP, chylomicrons are capable of reducing LPS activity within 0.5 h.

Postprandial chylomicron levels exceed the LPS-neutralizing capacity of circulating levels of HDL, LDL, and VLDL

It was demonstrated that lipoproteins have the ability to neutralize LPS in an LBP-dependent fashion (11). To determine to which extent postprandial chylomicron levels contribute to LPS neutralization by lipoproteins, the LPS-detoxifying capacity of lipoproteins was compared at concentrations equated with their physiologic plasma ratio. For chylomicrons, this is in regard to postprandial levels. All lipoproteins were diluted to 12.5% of their plasma concentration and preincubated with LPS and LBP. The




FIGURE 4. Chylomicrons reduce LPS toxicity in an LBP dose-dependent fashion. LPS (100 ng/ml) was preincubated with 0, 10, 50, and 100 ng/ml of LBP in the presence (●) or the absence (■) of isolated chylomicrons (12.5% of circulating levels) for 24 h at 37°C. The remaining biological activity of LPS was determined by incubation with PBMC for 4 h and measurements of TNF- α (A) and IL-8 (B) secretion in the supernatants. Values represent the mean \pm SD of three wells. The experiment shown is representative of three separate experiments. LBP catalyzed the LPS neutralization by chylomicrons.

ability of the remaining bioactive LPS to induce TNF- α was determined. In the absence of lipoproteins, 100 ng/ml of LPS together with 50 ng/ml of LBP led to a strong increase in TNF- α induction (Fig. 7). This TNF- α secretion was reduced by all tested lipoproteins. However, the most pronounced reduction of TNF- α secretion was observed when LPS was incubated with chylomicrons and LBP.

Chylomicrons reduce LTA-induced TNF secretion

The incidence of septic shock caused by Gram-positive bacteria has risen markedly in the last decade (28, 29). LTA, a component of the membrane of Gram-positive bacteria, can elicit the release of proinflammatory cytokines from immune cells (30). We determined whether chylomicrons in cooperation with LBP are also able to inhibit the stimulation of PBMC by LTA. Fig. 8 demonstrates that LBP induces detoxification of LTA by chylomicrons dose-dependently. These data suggest that besides a role in combating Gram-negative bacteria, LBP-chylomicron complexes may be of importance in the host defense against Gram-positive bacteria.




FIGURE 5. Chylomicrons neutralize LPS in a dose-dependent fashion. A range of chylomicron concentrations and 100 ng/ml LPS were incubated in the presence of 100 ng/ml of LBP. The remaining biological activity of LPS was assessed by incubation with PBMC for 4 h and measurement of IL-8 (●) and TNF-α (■) secretion. Values represent the mean ± SD of three wells. The experiment shown is representative of three separate experiments. Chylomicrons dose-dependently reduced LPS activity in the presence of LBP.




FIGURE 7. LPS-neutralizing capacity of chylomicrons compared with other circulating lipoproteins. Lipoproteins were isolated from human serum by ultracentrifugation. HDL, LDL, VLDL, and chylomicrons were diluted to 12.5% of their plasma concentrations and incubated with 100 ng/ml of LPS and 50 ng/ml of LBP for 24 h at 37°C. PBMC were added, and the remaining LPS activity was assessed by measuring the cytokine secretion of the cells after 4 h. Values represent the mean ± SD of three wells. The experiment shown is representative of three separate experiments. The LPS-neutralizing capacity of the chylomicrons exceeded the capacity of LDL, VLDL, and HDL to neutralize LPS.

Discussion

In the present study we obtained evidence for a functional interaction of LBP with chylomicrons. The association of LBP with chylomicrons was shown to enhance the LPS binding capacity of chylomicrons. Most importantly, chylomicrons and LBP cooperate in reducing LPS toxicity, resulting in decreased cytokine and chemokine secretion by PBMC. Moreover, chylomicron-LBP complexes were demonstrated to prevent cell activation by LTA, a cell wall component of Gram-positive micro-organisms. These data imply that LBP is an important cofactor for chylomicrons to execute their function as scavengers of bacterial toxin.

Consistent with others (31), we found that isolated chylomicrons bind LPS in the absence of LBP (Figs. 2 and 3). ApoB (18), and apoE (5), present in chylomicrons, were previously demonstrated to bind LPS and may account for LPS binding to chylomicrons. Binding of LPS to the phospholipids in the membrane of the chylomicrons under these conditions is not likely, since plasma proteins, such as phospholipid transfer protein (32), cholesteryl ester transfer protein, and LBP, were demonstrated to be required for intercalation of LPS into phospholipid membranes (33). In line with these findings, plasma factors were reported to be necessary for enhancement of the LPS binding capacity of chylomicrons and maximal prevention of LPS-induced death by i.v. administrated




FIGURE 6. LPS is rapidly neutralized by chylomicron-LBP complexes. LPS (100 ng/ml) was preincubated with (▨) or without (□) chylomicrons (12.5% of circulating levels) and 100 ng/ml of LBP for a 0.5-h period. Biologically active LPS remaining after 0.5 h was assessed by the addition of PBMC and measurement of cytokine secretion. Values represent the mean ± SD of three wells. The experiment shown is representative of three separate experiments. LPS neutralization by chylomicron-LBP complexes was evident after 0.5-h incubation.




FIGURE 8. LBP induces detoxification of LTA by chylomicrons. LTA (20 μg/ml) was preincubated with (▨) or without (□) chylomicrons (5% of the circulating level) and 0, 10, 50, and 100 ng/ml LBP for 24 h. The biologically active LTA remaining was assessed by the addition of PBMC and measurement of TNF secretion. Values represent the mean ± SD of three wells. The experiment shown is representative of three separate experiments. LBP and chylomicrons cooperate in detoxification of LTA.

lipoproteins (1). We now describe that LBP binds to chylomicrons and enhances the LPS-chylomicron interaction dose-dependently. Moreover, LBP was observed to accelerate LPS binding to chylomicrons. These data indicate that LBP is an important plasma factor involved in LPS binding to chylomicrons.

Harris et al. (2) reported that infusion of LPS together with chylomicrons compared with infusion of LPS alone resulted in lower levels of circulating TNF, the cytokine known to mediate much of the LPS toxicity *in vivo*. Moreover, they suggested that an enhanced uptake of LPS by hepatocytes and clearance of LPS from the circulation contribute to the reduced cytokine levels in such treated animals. In line with the reduction of TNF levels by chylomicrons observed *in vivo* by these investigators, we now found evidence for a direct LBP-mediated inhibiting effect of chylomicrons on LPS-induced cytokine secretion by PBMC *in vitro*. On the basis of our findings we postulate that the LPS-LBP-chylomicron interaction may participate in the reduced TNF levels *in vivo* by both enhanced detoxification of LPS and enhanced clearance of LPS complexed to chylomicrons from the circulation. Although in this study we determined the effect of preincubated LPS-chylomicron complexes on cellular activation, the capability of chylomicrons to bind LPS may also enable these lipoproteins to release LPS already bound to monocytes, as described for other lipoproteins (9). The finding that infusion of chylomicrons up to 30 min after the administration of endotoxin improved survival in rodents (2, 34) makes such a mechanism plausible.

To date, studies of LPS-inactivating properties of chylomicrons have been focused on circulating chylomicrons, and therapeutic application of chylomicrons was performed by systemic infusion of chylomicrons isolated from plasma or lymph (1, 2, 5, 7). Chylomicrons present in the circulation may cooperate with both hepatic and gut-derived LBP and detoxify circulating bacterial toxin. Besides detoxification of toxin in the circulation by LBP-chylomicron complexes, it can be hypothesized that this mechanism may be part of a local defense mechanism of the intestine against translocated bacterial toxin. Chylomicrons are produced by enterocytes, and interestingly, recent evidence was found for the basolateral secretion of LBP by the intestinal epithelium in response to cytokines (35). It is tempting to speculate from the data presented here that LBP and chylomicrons, both secreted by the intestinal mucosa, cooperate in local neutralization of LPS. Such a mechanism may occur in the subepithelial space or in the lymph and possibly prevents further damage of bacterial toxin to the intestinal epithelial barrier and entrance of biological active toxin into the circulation. A local function is only relevant when LPS detoxification by chylomicron-LBP complexes occurs rapidly. The data showed that LBP strongly accelerated the transfer of LPS to chylomicrons (Fig. 3), and in PBMC stimulation experiments we observed that LPS detoxification by chylomicrons was pronounced after 0.5 h of preincubation (Fig. 6). These observations support the potential relevance of a local role for chylomicrons and LBP in the prevention of LPS-induced damage. Such a mechanism is of special interest in the intestine, since this organ is loaded with high amounts of bacteria and their toxic products that under pathologic conditions are thought to translocate and then play an important role in inflammatory and infectious complications, such as multi-organ failure and sepsis (36).

Employing highly purified, essentially LPS-free LTA, we show that the phenomenon also extends to Gram-positive bacteria. LBP was not required for activation of PBMC by 20 $\mu\text{g/ml}$ of LTA, although LTA binds to this acute phase protein (37). Our data suggest that LBP and chylomicrons cooperate in the detoxification of LTA.

Enteral feeding is required for chylomicron secretion, and our findings may therefore explain at least in part the reported beneficial effects of oral feeding in critically ill patients (38). The physiological advantages of enteral nutrition as opposed to both starvation and parenteral nutrition are more and more accepted. Meta analysis has demonstrated that in a large variety of patients, including patients suffering from pancreatitis, ulcerative colitis, Crohn's disease, surgery, trauma, or multiorgan failure, enteral feeding was associated with a lower risk of infection compared with starvation and parenteral nutrition (38). Early postoperative enteral feeding was found to reduce septic morbidity (39) and hospital stay (40). However, in most centers patients are restricted from early postoperative enteral feeding and from enteral feeding during their stay in the intensive care unit. Such abstinence of enteral feeding may cause a decline in the capacity of the body to detoxify bacterial toxin. We consider that enteral feeding of critically ill patients exploits a physiologic mechanism to enhance the endogenous chylomicron production, potentially leading to an enhanced presence of LPS-inactivating LBP-chylomicron complexes in the circulation and, possibly more important, in the intestinal subepithelial space and lymph. The chylomicron levels attained by oral feeding seem to be of relevance, since postprandial levels of chylomicrons are more potent in LPS neutralization than physiologic levels of other lipoprotein fractions (HDL, LDL, and VLDL) *in vitro*. We are not able to conclude from the data whether this greater inhibiting capacity of chylomicrons is due to a higher relative affinity for LPS, higher LBP or LPS binding capacity, or a molar excess of circulating chylomicrons compared with the other lipoprotein classes. Since even low concentrations of chylomicrons exert a significant effect on LPS activity, and postprandial chylomicron levels rise acutely and drop slowly over hours thereafter (41), the effect of feeding is most likely to persist for several hours.

Overall, in this study we obtained evidence for a cooperative function between LBP and chylomicrons in the detoxification of toxin from both Gram-positive and Gram-negative microorganisms. Enhanced binding of LPS to LBP-chylomicron complexes in the circulation may prevent cytokine production and most likely enhances their clearance from the circulation. We further speculated that the production of LBP-chylomicron complexes in the gut may represent a natural defense mechanism against endotoxemia of enteric origin. Enteral feeding and subsequent chylomicron production may be a preventive or even therapeutic intervention in patients who undergo abdominal surgery or major cardiovascular surgery, since in those patients bacterial translocation across the epithelial barrier of the gut is thought to play an important role in postoperative complications (36).

References

- Harris, H. W., C. Grunfeld, K. R. Feingold, and J. H. Rapp. 1990. Human very low density lipoproteins and chylomicrons can protect against endotoxin-induced death in mice. *J. Clin. Invest.* 86:696.
- Harris, H. W., C. Grunfeld, K. R. Feingold, T. E. Read, J. P. Kane, A. L. Jones, E. B. Eichbaum, G. F. Bland, and J. H. Rapp. 1993. Chylomicrons alter the fate of endotoxin, decreasing tumor necrosis factor release and preventing death. *J. Clin. Invest.* 91:1028.
- Munford, R. S., J. M. Andersen, and J. M. Dietschy. 1981. Sites of tissue binding and uptake *in vivo* of bacterial lipopolysaccharide-high density lipoprotein complexes. *J. Clin. Invest.* 68:1503.
- Munford, R. S., C. L. Hall, J. M. Lipton, and J. M. Dietschy. 1982. Biological activity, lipoprotein-binding behavior and *in vivo* disposition of extracted and native forms of *Salmonella typhimurium* lipopolysaccharides. *J. Clin. Invest.* 70:877.
- Rensen, P. C. N., M. van Oosten, E. van de Bilt, M. van Eck, J. Kuiper, and T. J. C. van Berkel. 1997. Human recombinant apolipoprotein E redirects lipopolysaccharide from Kupffer cells to liver parenchymal cells in rats *in vivo*. *J. Clin. Invest.* 99:2438.
- Harris, H. W., S. E. Brady, and J. H. Rapp. 2002. Hepatic endosomal trafficking of lipoprotein-bound endotoxin in rats. *J. Surg. Res.* 106:188.

7. Read, T. E., H. W. Harris, C. Grunfeld, K. R. Feingold, M. C. Calhoun, J. P. Kane, and J. H. Rapp. 1993. Chylomicrons enhance endotoxin excretion in Bile. *Infect. Immun.* 61:3496.
8. Cavaillon, J.-M., C. Fitting, N. Haeflner-Cavaillon, S. J. Kirsch, and H. S. Warren. 1990. Cytokine response by monocytes and macrophages to free and lipoprotein-bound lipopolysaccharide. *Infect. Immun.* 58:2375.
9. Kitchens, R. L., P. A. Thompson, S. Viriyakosol, G. E. O'Keefe, and R. S. Munford. 2002. Plasma CD14 decreases monocyte responses to LPS by transferring cell-bound LPS to plasma lipoproteins. *J. Clin. Invest.* 108:485.
10. Munford, R. S., C. L. Hall, and J. M. Dietschy. 1981. Binding of *Salmonella typhimurium* lipopolysaccharides to rat high-density lipoproteins. *Infect. Immun.* 34:835.
11. Wurfel, M. M., and S. D. Wright. 1994. Lipopolysaccharide-binding protein is carried on lipoproteins and acts as a shuttle in the neutralization of LPS by LPS-binding cofactor in the neutralization of LPS. *J. Exp. Med.* 180:1025.
12. Zweigner, J., H. J. Gramm, O. C. Singer, K. Wegscheider, and R. R. Schumann. 2001. High concentrations of lipopolysaccharide-binding protein in serum of patients with severe sepsis or septic shock inhibit the lipopolysaccharide response in human monocytes. *Blood* 15:3800.
13. Grunfeld, C., M. Marshall, J. K. Shigenaga, A. H. Moser, P. Tobias, and K. R. Feingold. 1999. Lipoproteins inhibit macrophage activation by lipoteichoic acid. *J. Lipid Res.* 40:245.
14. Ulevitch, R. J., A. R. Johnston, and D. B. Weinstein. 1979. New function for high density lipoproteins. *J. Clin. Invest.* 64:1516.
15. Van Lenten, B. J., A. M. Fogelman, M. E. Haberland, and P. A. Edwards. 1986. The role of lipoproteins and receptor-mediated endocytosis in the transport of bacterial lipopolysaccharide. *Proc. Natl. Acad. Sci. USA* 83:2704.
16. Victorov, A. V., N. V. Medvedeva, E. M. Gladkaya, A. D. Morozkin, E. A. Podrez, V. A. Kosykh, and V. A. Yurkiv. 1989. Composition and structure of lipopolysaccharide-human plasma low density lipoprotein complex. *Biochim. Biophys. Acta* 984:119.
17. Netea, M. G., P. N. M. Demacker, B. J. Kullberg, L. E. H. Jacobs, T. J. G. Verver-Jansen, O. C. Boerman, A. F. H. Stalenhoef, and J. W. M. van der Meer. 1998. Bacterial lipopolysaccharide binds and stimulates cytokine producing cells before neutralization by endogenous lipoproteins can occur. *Cytokine* 10:766.
18. Vreugdenhil, A. C. E., A. M. P. Snoek, C. van 't Veer, J.-W. M. Greve, and W. A. Buurman. 2001. Lipopolysaccharide binding protein circulates associated with apoB containing lipoproteins and enhances endotoxin-LDL/VLDL interaction. *J. Clin. Invest.* 107:225.
19. Gao, J. J., Q. Xue, E. G. Zuvanich, K. R. Haghi, and D. C. Morrison. 2001. Commercial preparations of lipoteichoic acid contain endotoxin that contributes to activation of mouse macrophages in vitro. *Infect. Immun.* 69:751.
20. Morath, S., A. Geyer, and T. Hartung. 2001. Structure-function relationship of cytokine induction by lipoteichoic acid from *Staphylococcus aureus*. *J. Exp. Med.* 193:393.
21. Terpstra, A. H. M. 1985. Isolation of serum chylomicrons prior to density gradient ultracentrifugation of other serum lipoprotein classes. *Anal. Biochem.* 150:221.
22. Terpstra, A. H. M., C. J. H. Woodward, and F. J. Sanchez-Muniz. 1981. Improved techniques for the separation of serum lipoproteins by density gradient ultracentrifugation: visualization by prestaining and rapid separation of serum lipoproteins from small volumes of serum. *Anal. Biochem.* 111:149.
23. Froom, A. H. M., M. A. Dentener, J. W. M. Greve, G. Ramsay, and W. A. Buurman. 1995. LPS toxicity regulating proteins in bacteremia. *J. Infect. Dis.* 171:1250.
24. Yu, B., and S. D. Wright. 1996. Catalytic properties of lipopolysaccharide (LPS) binding protein, transfer of CD14. *J. Biol. Chem.* 271:4100.
25. Engelberts, I., A. Moller, G. J. M. Schoen, C. J. van der Linden, and W. A. Buurman. 1991. Evaluation of measurement of human TNF in plasma by ELISA. *Lymphokine Cytokine Res.* 10:69.
26. Bouma, M. G., R. K. Stad, F. A. J. M van de Wildenberg, and W. A. Buurman. 1994. Differential regulatory effects of adenosine on cytokine release by activated human monocytes. *J. Immunol.* 153:4159.
27. Lamping, N., R. Dettmer, N. W. J. Schroder, D. Pfeil, W. Hallatschek, R. Burger, and R. R. Schumann. 1998. LPS-binding protein protects mice from septic shock caused by LPS or Gram-negative bacteria. *J. Clin. Invest.* 101:2065.
28. Bone, R. C. 1994. Gram-positive organisms and sepsis. *Arch. Intern. Med.* 154:26.
29. Nogare, A. R., 1991. Southwestern Internal Medicine Conference: septic shock. *Am. J. Med. Sci.* 302:50.
30. Bhakdi, S., T. Klonisch, P. Nuber, and W. Fischer. 1991. Stimulation of monokine production by lipoteichoic acids. *Infect. Immun.* 59:4614.
31. Eichbaum, E. B., H. W. Harris, J. P. Kane, and J. H. Rapp. 1991. Chylomicrons can inhibit endotoxin activity in vitro. *J. Surg. Res.* 51:413.
32. Hailman, E., J. J. Albers, G. Wolfbauer, A. Y. Tu, and S. D. Wright. 1996. Neutralization and transfer of lipopolysaccharide by phospholipid transfer protein. *J. Biol. Chem.* 24:12172.
33. Wurfel, M. M., and S. D. Wright. 1997. Lipopolysaccharide-binding protein and soluble CD14 transfer lipopolysaccharide to phospholipid bilayers. *J. Immunol.* 158:3925.
34. Read, T. E., C. Grunfeld, Z. Kumwenda, M. C. Calhoun, J. P. Kane, K. R. Feingold, and J. H. T. Rapp. 1995. Triglyceride-rich lipoproteins improve survival when given after endotoxin in rats. *Surgery* 117:62.
35. Vreugdenhil, A. C. E., M. A. Dentener, A. M. P. Snoek, J.-W. Greve, and W. A. Buurman. 1999. Lipopolysaccharide binding protein and serum amyloid A secretion by human intestinal epithelial cells during the acute phase response. *J. Immunol.* 163:2792.
36. O'Boyle, C. J., J. MacFie, C. J. Mitchell, D. Johnstone, P. M. Sagar, and P. C. Sedman. 1998. Microbiology of bacterial translocation in humans. *Gut* 42:29.
37. Fan, X., F. Stelter, R. Menzel, R. Jack, I. Spreitzer, T. Hartung, and C. Schütt. 1999. Structures in *Bacillus subtilis* are recognized by CD14 in a lipopolysaccharide binding protein-dependent reaction. *Infect. Immun.* 67:2964.
38. Braunschweig, C. L., P. Levy, P. M. Sheehan, and X. Wang. 2001. Enteral compared with parenteral nutrition: a meta-analysis. *Am. J. Clin. Nutr.* 74:534.
39. Moore, F. A., E. E. Moore, T. N. Jones, B. L. McCroskey, and V. M. Peterson. 1989. TEN versus TPN following major abdominal trauma reduced septic morbidity. *J. Trauma* 29:916.
40. Schilder, J. M., J. A. Hurteau, K. Y. Look, D. H. Moore, G. Raff, F. B. Stehman, and G. P. Sutton. 1997. A prospective controlled trial of early postoperative oral intake following major abdominal gynecologic surgery. *Gynecol. Oncol.* 67:235.
41. Hugh, P., and R. Barett. 1998. Kinetics of triglyceride rich lipoproteins: chylomicrons and very low density lipoproteins. *Atherosclerosis* 141(Suppl. 1):S35.