

BULK ANTIBODIES  
for *in vivo*  
RESEARCH

$\alpha$ -PD-1

$\alpha$ -PD-L1

$\alpha$ -4-1BB

$\alpha$ -CTLA4

$\alpha$ -LAG3

Discover More

BioCell



The Journal of  
Immunology

## CORRECTIONS

*J Immunol* 2004; 172:4647; ;  
doi: 10.4049/jimmunol.172.7.4647  
<http://www.jimmunol.org/content/172/7/4647>

This information is current as  
of October 17, 2019.

Why *The JI*? [Submit online.](#)

- **Rapid Reviews!** 30 days\* from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

\*average

**Subscription** Information about subscribing to *The Journal of Immunology* is online at:  
<http://jimmunol.org/subscription>

**Permissions** Submit copyright permission requests at:  
<http://www.aai.org/About/Publications/JI/copyright.html>

**Email Alerts** Receive free email-alerts when new articles cite this article. Sign up at:  
<http://jimmunol.org/alerts>

*The Journal of Immunology* is published twice each month by  
The American Association of Immunologists, Inc.,  
1451 Rockville Pike, Suite 650, Rockville, MD 20852  
Copyright © 2004 by The American Association of  
Immunologists All rights reserved.  
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



# CORRECTIONS

Bernhard Banas, Markus Wörnle, Thorsten Berger, Peter J. Nelson, Clemens D. Cohen, Matthias Kretzler, Jochen Pfirstinger, Matthias Mack, Martin Lipp, Hermann-Josef Gröne, and Detlef Schlöndorff. Roles of SLC/CCL21 and CCR7 in Human Kidney for Mesangial Proliferation, Migration, Apoptosis, and Tissue Homeostasis. *The Journal of Immunology* 2002;168:4301–4307.

In *Materials and Methods*, under the heading *RT-PCR analysis*, the primer sequences for CCR7 are incorrect. The other data on the probes (accession number and positions of the primers) are correct as published. The correct primer sequences are shown below.

sense: 5'-GCTCCAGGCACGCAACTTT-3'  
 antisense: 5'-ACCACGACCACAGCGATGA-3'  
 probe (FAM): 5'-AGCGCAACAAGGCCATCAAGGTG-3'

Leonid Gorelik, Anne H. Cutler, Greg Thill, Steven D. Miklasz, Dianna E. Shea, Christine Ambrose, Sarah A. Bixler, Lihe Su, Martin L. Scott, and Susan L. Kalled. Cutting Edge: BAFF Regulates CD21/35 and CD23 Expression Independent of Its B Cell Survival Function. *The Journal of Immunology* 2004;172:762–766.

In *Results*, there were errors in Table II. The legend and conclusion are correct as published. The revised table is shown below.

Table II. *BAFF/APRIL blockade in bcl-2 Tg mice*

Treatment	B Cells <sup>a</sup> (×10 <sup>7</sup> )	CD21/35 <sup>b</sup> (MFI)	CD23 <sup>b</sup> (MFI)
BCMA-Fc	15.8	13.7	145
BCMA-Fc	16.0	13.0	136
BCMA-Fc	24.3	12.1	174
Mean ± SD	18.7 ± 4.9	12.9 ± 0.8	151.7 ± 19.9
hIgG	21.8	16.9	256
hIgG	20.7	19.2	279
Mean	21.3	18.1	268

<sup>a</sup> Assessed on day 8 after 2 doses of BCMA-Fc.

<sup>b</sup> Assessed 2 days after a single dose of BCMA-Fc.

Sergei Kusmartsev, Yulia Nefedova, Daniel Yoder, and Dmitry I. Gabrilovich. Antigen-Specific Inhibition of CD8<sup>+</sup> T Cell Response by Immature Myeloid Cells in Cancer Is Mediated by Reactive Oxygen Species. *The Journal of Immunology* 2004;172:989–999.

In *Results*, the flow cytometry histogram labels MFI and M1 are incorrect in three of the four panels in Figure 6A. The revised figure is shown below.

