



Vaccine Adjuvants

Take your vaccine to the next level

InvivoGen



The Journal of
Immunology

Cutting Edge: Contact-Mediated Suppression by CD4⁺CD25⁺ Regulatory Cells Involves a Granzyme B-Dependent, Perforin-Independent Mechanism

This information is current as of October 21, 2021.

David C. Gondek, Li-Fan Lu, Sergio A. Quezada, Shimon Sakaguchi and Randolph J. Noelle

J Immunol 2005; 174:1783-1786; ;

doi: 10.4049/jimmunol.174.4.1783

<http://www.jimmunol.org/content/174/4/1783>

References This article **cites 27 articles**, 14 of which you can access for free at: <http://www.jimmunol.org/content/174/4/1783.full#ref-list-1>

Why *The JI*? Submit online.

- **Rapid Reviews! 30 days*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

**average*

Subscription Information about subscribing to *The Journal of Immunology* is online at: <http://jimmunol.org/subscription>

Permissions Submit copyright permission requests at: <http://www.aai.org/About/Publications/JI/copyright.html>

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at: <http://jimmunol.org/alerts>

The Journal of Immunology is published twice each month by
The American Association of Immunologists, Inc.,
1451 Rockville Pike, Suite 650, Rockville, MD 20852
Copyright © 2005 by The American Association of
Immunologists All rights reserved.
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



CUTTING EDGE

Cutting Edge: Contact-Mediated Suppression by CD4⁺CD25⁺ Regulatory Cells Involves a Granzyme B-Dependent, Perforin-Independent Mechanism¹

David C. Gondek,^{2*} Li-Fan Lu,^{2*} Sergio A. Quezada,^{*} Shimon Sakaguchi,[†] and Randolph J. Noelle^{3*}

CD4⁺CD25⁺ regulatory T cells (T_{reg}) are potent immunosuppressive cells that are pivotal in the regulation of peripheral tolerance. In this report, we identify granzyme B (GZ-B) as one of the key components of T_{reg}-mediated suppression. Induction of regulatory activity is correlated with the up-regulation of GZ-B expression. Proof of a functional involvement of GZ-B in contact-mediated suppression by T_{reg} is shown by the reduced ability of T_{reg} from GZ-B^{-/-} mice to suppress as efficiently as T_{reg} from WT mice. GZ-B-mediated suppression is perforin independent, because suppression by T_{reg} from perforin^{-/-} and WT is indistinguishable. Additionally, suppression mediated by T_{reg} appears to be mediated, in part, by the induction of apoptosis in the CD4⁺CD25⁻ effector cell. In summary, GZ-B is one of the key mechanisms through which CD4⁺CD25⁺ T_{reg} induce cell contact-mediated suppression. *The Journal of Immunology*, 2005, 174: 1783–1786.

Recent studies have underscored the importance of regulatory T cells (T_{reg})⁴ in preventing the emergence of autoimmune disease, dampening the intensity of immune responses to pathogens and mediating peripheral transplantation tolerance. Multiple subsets of T_{reg} have been implicated in these processes and include the thymically derived CD4⁺CD25⁺ T_{reg} (1), as well as an inducible regulatory T cell (Tr1) subset (2). A major focus of study has been to molecularly characterize the mechanisms that mediate T_{reg} suppression of immunity. It has been shown that Tr1 suppress predominantly by a cytokine-dependent mechanism characterized by IL-10 and TGF-β secretion (3). Similarly, a TGF-β-dependent mechanism has also been implicated in suppression by CD4⁺CD25⁺ T_{reg} (4–6). In addition to suppression via soluble factors, the CD4⁺CD25⁺ T_{reg} have been shown to mediate suppression via a contact-dependent mechanism (7). The mo-

lecular basis for contact-dependent suppression by CD4⁺CD25⁺ T_{reg} is not known.

Glucocorticoid-induced TNF-like receptor (GITR or TNFSF18) is a member of the TNFR family that is constitutively expressed on T_{reg} and inducibly expressed on CD4⁺CD25⁻ effector T cells (T_{eff}) (8, 9). Triggering of GITR has been shown to extinguish their contact-dependent suppressive activity (8, 10). Based on this overt change in biological function, transcriptional profiling of resting, activated T_{reg}, and anti-GITR-treated activated T_{reg} has led to the identification of a number of candidate molecules that may be involved in contact-dependent suppression. One such molecule that was identified as up-regulated in activated T_{reg} and whose expression was reduced via GITR-triggering is granzyme B (GZ-B).

GZ-B is a serine protease, secreted mainly by NK cells and CTLs (11), and is largely responsible for the induction of apoptosis in the target cell. However, recent reports have shown that human CD4⁺ T cells are also able to synthesize GZ-B and perforin (12, 13). Furthermore, studies by Ley and coworkers as well as others (14, 15) have shown that GZ-B is highly up-regulated in activated human T cells bearing a Tr1 phenotype. Moreover, Ley and coworkers (16) have shown CD4⁺CD25⁺ T_{reg} in the human system mediate suppression with requirement for granzyme A (GZ-A). These results suggest a possible role for granzyme in mediating T cell suppression. The data presented in this study implicate that GZ-B plays a pivotal role in the suppressive capacity of murine CD4⁺CD25⁺ T_{reg}.

Materials and Methods

Mice and materials

Congenic strains CD45.1 or CD45.2 C57BL/6 and perforin^{-/-} mice, 8–10 wk old, were purchased from The Jackson Laboratory. C57BL/6 GZ-B^{-/-} mice (15) were bred and maintained in our facility at Dartmouth Medical School.

*Department of Microbiology and Immunology, Dartmouth Medical School and Norris Cotton Cancer Center, Lebanon, NH 03756; and [†]Department of Experimental Pathology, Institute for Frontier Medical Sciences, and Department of Transplantation and Immunology, Faculty of Medicine, Kyoto University, Kyoto, Japan

Received for publication August 18, 2004. Accepted for publication December 7, 2004.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This work was supported by National Institutes of Health Grants CA91436-01 and AI48667.

² D.C.G. and L.-F.L. contributed equally to this manuscript.

³ Address correspondence and reprint requests to Dr. Randolph J. Noelle, Department of Microbiology and Immunology, Norris Cotton Cancer Center, Dartmouth Hitchcock Medical Center, Dartmouth Medical School, 1 Medical Center Drive, Lebanon, NH 03756. E-mail address: rjn@dartmouth.edu

⁴ Abbreviations used in this paper: T_{reg}, regulatory T cell; T_{eff}, effector T cell; GITR, glucocorticoid-induced TNF-like receptor; GZ-B, granzyme B; GZ-A, granzyme A; PI, propidium iodide.

Cell isolation, gene array, and real time

Single-cell suspensions were prepared from 8- to 10-wk-old mice and applied to CD4 enrichment. CD4⁺CD25⁻ and CD4⁺CD25⁺ T cells were further purified by magnetic separation with MACS (Miltenyi Biotec) according to the manufacturer's instructions. Enriched cell populations and purified cells were phenotypically analyzed by FACS. The purities of CD4⁺CD25⁻ and CD4⁺CD25⁺ T cells were 90–95%, respectively. Freshly isolated cells have been inoculated (10⁶/ml; complete RPMI 1640/10% FBS supplemented with 100 U of IL2) into a 24-well plate precoated with 10 μg/ml anti-CD3 (clone 2C11) with or without 10 μg/ml anti-GITR (clone DTA-1; Ref. 8) cultured at 37°C for 0, 12, and 48 h. Purified RNA were then analyzed by using Affymetrix mouse genome A430 oligonucleotide arrays or by real-time PCR analysis.

Cell culture and T cell suppression assay

GZ-B expression was assessed in freshly isolated CD4⁺CD25⁺ T cells or in cells cultured in vitro 24–72 h in the presence of plate-bound CD3 (1 μg/ml) with 100 U/ml IL-2.

Spleens and lymph nodes from wild-type, perforin^{-/-}, or GZ-B^{-/-} mice were magnetic bead sorted as stated above. Further purification of the T_{eff} subset was accomplished with a CD4⁺ T cell Isolation kit (Miltenyi Biotec). Effector cells were >95% pure at the end of this isolation. In a polyclonal T_{reg} suppressor assay, CD4⁺CD25⁻ T_{eff} cells (5 × 10⁴) were cocultured with irradiated T-depleted splenocytes (1 × 10⁵), 5 μg/ml anti-CD3, and indicated numbers of CD4⁺CD25⁺ cells for 3 days. In some experiments, 5 μg/ml anti-GITR was also added to the wells. Proliferation was assessed by incorporation of [³H]thymidine (1 μCi/well), which was added for the last 8 h of culture.

Cell surface, intracellular staining, and flow cytometry

Approximately 2 × 10⁵ cells from each of triplicate wells were collected and pooled. Cells were labeled with anti-CD45.1-allophycocyanin (clone A20; eBioscience). Samples were then resuspended in 1 × annexin staining buffer and treated with Annexin V^{FITC} (BD Pharmingen) and propidium iodide (PI; Sigma-Aldrich). For GZ-B expression assay, following isolation for fresh T_{reg} or 24–72 h for cultured T_{reg} cells were stained with anti-CD4-FITC (clone RM 4-5) and anti-CD25-PE (clone PC-61). Samples were then fixed and permeabilized (Cytofix/Cytoperm; BD Pharmingen) and stained with anti-human GZ-B-allophycocyanin (clone GB12; Caltag) diluted 1/200 in staining buffer. Throughout all steps, normal rat serum (5% v/v; Invitrogen Life Technologies) was used to block nonspecific binding. Samples were analyzed on FACScan (BD Biosciences). Anti-human GZ-B cross-reactivity with mouse GZ-B has been previously reported (15). For CFSE experiments, CD45.1⁺ cells were labeled with 5 μM CFSE and added to suppressor assay as described above.

Statistical analysis

Analysis of proliferation assays and real-time expression between the various treatment groups were analyzed by two-tailed, paired Student's *t* test. Values of *p* < 0.05 were considered significant.

Results and Discussion

Anti-GITR causes down-regulation of GZ-B in in vitro-cultured T_{reg}

CD4⁺CD25⁺ T_{reg} are suppressive to naive CD4⁺ T_{eff} in vitro following polyclonal and Ag-specific activation. Furthermore, the in vitro suppressive capacity has been shown to be contact dependent and ablated following treatment with anti-GITR (8). Global gene analysis of activated T_{reg} treated or untreated with anti-GITR was used to identify candidate genes involved in suppression. We examined naive and activated T_{eff} (purified CD4⁺CD25⁻ T cells) and T_{reg} (purified CD4⁺CD25⁺ T cells) in the presence of anti-CD3 with or without anti-GITR for 12 or 48 h. Of the ~22,700 genes examined, 259 were up-regulated >1.5-fold and 99 were down-regulated >1.5-fold in T_{reg} following treatment with anti-GITR and anti-CD3 relative to treatment with anti-CD3 alone. GZ-B, as has been shown previously, is up-regulated with T_{reg} activation via anti-CD3 alone (9, 17). Studies presented herein show that that GZ-B is down-regulated 2-fold with anti-CD3 in combination with anti-GITR (Fig. 1, *A* and *B*). The microarray data was confirmed by RT-PCR (Fig. 1*C*). At both the 12- and 48-h time point, the levels GZ-B expression are 2-fold greater with anti-CD3 alone treatment vs combining with anti-GITR stimula-




FIGURE 1. Activation-induced up-regulation of GZ-B in T_{reg} and its regulation by GITR. *A*, Cells were purified via MACS columns to >90% purity and cultured in vitro with anti-CD3 and IL-2 with or without anti-GITR for 12 h. RNA was prepared and hybridized to the Affymetrix A430 array. Relative expression indicates the mean log₂ ratio of changes in T_{reg} expression between anti-CD3-alone treatment and anti-CD3 with anti-GITR. *B*, Gene chip signal intensity comparison of GZ-A, GZ-B, and perforin following treatments for 12 or 48 h. *C*, Real-time RT-PCR analysis of GZ-B expression in T_{reg} treated as described above. Figure is representative of two independent experiments. *D*, GZ-B expression in freshly isolated T_{reg} or in vitro cultured for 24 or 72 h with plate-bound anti-CD3 and 100 U of IL-2.

tion. Moreover, protein expression of GZ-B recapitulates the results found via RT-PCR by increasing the abundance of GZ-B from 24 to 72 h (Fig. 1D). Additionally, after 12 h in culture, T_{reg} GZ-B mRNA expression is 20-fold greater than T_{eff} with CD3 stimulation alone (data not shown). We also examined expression levels of GZ-A and perforin at all time points. For both molecules, we see similar regulation to that of GZ-B with anti-GITR treatment; however, expression is at a much lower intensity at all time points (Fig. 1, A and B). These data were also confirmed by RT-PCR (data not shown). These data indicate that, immediately following activation, T_{reg} rapidly up-regulate GZ-B; however, GZ-A and perforin remain low in abundance relative to GZ-B expression.

GZ-B^{-/-} T_{reg} have reduced suppressive capacity in vitro

Coculture of T_{reg} with wild-type T_{eff} leads to suppression of proliferation in a dose-dependent manner. To functionally evaluate the role of GZ-B in the contact-mediated suppression by T_{reg} , the suppressive activity of T_{reg} from WT and GZ-B^{-/-} mice was compared (Fig. 2A). Data presented show that T_{reg} from WT mice at a 1:1 ratio suppress the proliferation of T_{eff} >90%, whereas T_{reg} from GZ-B^{-/-} mice suppress T_{eff} proliferation <50%. The reduced suppressive activity of T_{reg} from GZ-B^{-/-} mice is observed across a spectrum of T_{reg} : T_{eff} ratios, suggesting a functional role of GZ-B in contact-mediated suppression. A comparison of FoxP3 levels of GZ-B^{-/-} T_{reg} revealed no significant difference from those of WT T_{reg} (data not shown). Because loss of GZ-B does not completely extinguish T_{reg} suppression, additional contact-dependent mechanisms must be important in this system.




FIGURE 2. GZ-B mediates T_{reg} suppression via a perforin-independent mechanism. *A*, T_{reg} isolated from wild-type or GZ-B^{-/-} mice were cocultured with CD4⁺CD25⁻ T_{eff} and irradiated T-depleted APC with anti-CD3 for 72 h. Wells were pulsed with 1 μ Ci/well ³H for the last 8 h of culture, and analyzed as described in *Materials and Methods*. *B*, T_{reg} isolated from wild-type or perforin^{-/-} mice were cultured as above.

Typically, GZ-B requires cytosolic entry via perforin or a perforin-like molecule to induce cell death (18, 19). To determine whether suppression is mediated by the canonical GZ-B-perforin pathway, the suppressor activity of T_{reg} from perforin^{-/-} vs WT mice was determined (Fig. 2B). The suppressive activity of T_{reg} from perforin^{-/-} and WT mice was indistinguishable, suggesting that GZ-B-mediated suppression is via a novel, perforin-independent mechanism. Moreover, studies of perforin^{-/-} mice on the BALB/c background showed similar results (data not shown). Several reports indicate that, at high concentrations of GZ-B, the necessity of perforin to release GZ-B from the endosomal compartment can be circumvented (20, 21). Additionally, Choy et al. (22) have shown that GZ-B is able to induce cell death in a perforin-independent manner mediated by a combination of intracellular and extracellular events.

A recent report by Grossman et al. (16) indicates that human CD4⁺CD25⁺ T_{reg} mediate their suppressive effects via death induced by a GZ-A perforin-dependent mechanism. The differences between the use of GZ-A in humans and GZ-B in mice could be due to species differences, or subtle differences in the subsets and/or activation of T cells that were used. With regard to perforin dependency, the study by Ley (16) implicates perforin because of the fact that a calcium chelator relieves suppression. Although this is a reasonable assertion, they also show that CD18 is required, and it is known that this molecule requires calcium to form the tight synapse required for granzyme-mediated toxicity (23–25). In our studies using perforin knockout mice, suppression was indistinguishable from WT mice.

Induction of T_{eff} apoptosis is a component of contact-mediated suppression

Recent reports have re-examined apoptosis by T_{reg} of T_{eff} as a mechanism for suppression (15, 26). The molecule(s) that mediate the induction of T_{eff} apoptosis have not been resolved, and it is unlikely that FasL plays a central role (27). Based on the finding that GZ-B plays a functionally significant role in T_{reg} suppression, the ability of T_{reg} to induce T_{eff} apoptosis and cell death was re-examined. The induction of T_{eff} apoptosis by T_{reg} was determined following the in vitro coculture of activated T_{eff} and T_{reg} . Briefly, CD45.1⁺ (Ly5.2⁺) T_{eff} were cocultured with increasing numbers of CD45.2⁺ (Ly5.1⁺) T_{reg} , in the presence of anti-CD3. After 72 h of culture, apoptosis of the CD4⁺ T_{eff} was determined by multiparameter flow cytometry. The data show that there is a dose-dependent increase in cell death of the T_{eff} cells when cocultured with T_{reg} , such that ~50% more T_{eff} are dead at a 1:1 ratio than at a 1:16 ratio of T_{reg} to T_{eff} (Fig. 3A). Moreover, addition of anti-GITR relieves the suppression and apoptosis as evidenced by enhanced proliferation and cell survival (data not shown). In parallel experiments, we examined thymidine incorporation in a standard suppressor assay with T_{reg} from wild-type mice treated with anti-CD3 to determine levels of suppressive activity concurrent with PI/annexin staining (Fig. 3B). To distinguish between the antiproliferative and antiapoptotic effect of T_{reg} , we examined suppression and death with CFSE-labeled T_{eff} counterstained with PI. In Fig. 3C, we demonstrate that the T_{eff} have a greater percentage of PI⁺ cells when cocultured with T_{reg} . Interestingly, in addition to the induction of cell death, the proliferation of PI⁻ T_{eff} was also inhibited, which indicates multiple mechanisms are involved in T_{reg} -mediated suppression.




FIGURE 3. T_{reg} suppression is mediated by enhanced death in T_{eff}. *A*, Ly5.2⁺ T_{eff} and Ly5.1⁺ T_{reg} and APC were cocultured for 72 h in a standard suppressor assay. Wells were harvested and stained with anti-Ly5.2 to identify the CD4⁺CD25⁻ T_{eff}. Cells were then stained with PI and Annexin V^{FITC}. *B*, Proliferation of cells in wells run simultaneously with stained wells. Cells were pulsed with 1μCi/well ³H for the last 8 h of culture. *C*, CFSE-labeled cells were cultured for 72 h in standard suppressor assay, and then counterstained with PI.

The data presented above suggest that GZ-B is pivotal for complete suppression by T_{reg}. The fact that the expression of GZ-B is up-regulated by T_{reg} activation, and impaired by anti-GITR triggering is consistent with biological activity manifested by T_{reg} under these conditions. Functionally, the role of GZ-B in T_{reg} activities was strongly suggested by the reduced ability of T_{reg} from GZ-B^{-/-} mice to suppress T_{eff} proliferation. Surprisingly, it appears that the GZ-B-dependent suppression of T_{eff} activities is not dependent on perforin, because the T_{reg} from perforin^{-/-} mice suppressed T_{eff} proliferation equivalent to that observed from wild-type mice. Hence, the precise mechanism of GZ-B-mediated suppression is enigmatic. In this context, GZ-B-mediated apoptosis in the absence of perforin has been reported in other systems. Finally, death as a consequence of T_{reg} action was demonstrated by the fact that CD4⁺CD25⁺ T_{reg} in a dose-dependent manner, can induce apoptosis in T_{eff}. The current studies raise many questions as to how GZ-B mediates T_{eff} death, and the role of GZ-B in the in vivo function of T_{reg}.

References

- Jordan, M. S., A. Boesteanu, A. J. Reed, A. L. Petrone, A. E. Holenbeck, M. A. Lerman, A. Najj, and A. J. Caton. 2001. Thymic selection of CD4⁺CD25⁺ regulatory T cells induced by an agonist self-peptide. *Nat. Immunol.* 2:301.
- Levings, M. K., and M. G. Roncarolo. 2000. T-regulatory 1 cells: a novel subset of CD4 T cells with immunoregulatory properties. *J. Allergy Clin. Immunol.* 106:S109.
- Roncarolo, M. G., R. Bacchetta, C. Bordignon, S. Narula, and M. K. Levings. 2001. Type 1 T regulatory cells. *Immunol. Rev.* 182:68.
- Nakamura, K., A. Kitani, and W. Strober. 2001. Cell contact-dependent immunosuppression by CD4⁺CD25⁺ regulatory T cells is mediated by cell surface-bound transforming growth factor β. *J. Exp. Med.* 194:629.
- Piccirillo, C. A., J. J. Letterio, A. M. Thornton, R. S. McHugh, M. Mamura, H. Mizuhara, and E. M. Shevach. 2002. CD4⁺CD25⁺ regulatory T cells can mediate suppressor function in the absence of transforming growth factor β1 production and responsiveness. *J. Exp. Med.* 196:237.
- Nakamura, K., A. Kitani, I. Fuss, A. Pedersen, N. Harada, H. Nawata, and W. Strober. 2004. TGF-β1 plays an important role in the mechanism of CD4⁺CD25⁺ regulatory T cell activity in both humans and mice. *J. Immunol.* 172:834.
- Shevach, E. M. 2002. CD4⁺CD25⁺ suppressor T cells: more questions than answers. *Nat. Rev. Immunol.* 2:389.
- Shimizu, J., S. Yamazaki, T. Takahashi, Y. Ishida, and S. Sakaguchi. 2002. Stimulation of CD25⁺CD4⁺ regulatory T cells through GITR breaks immunological self-tolerance. *Nat. Immunol.* 3:135.
- McHugh, R. S., M. J. Whitters, C. A. Piccirillo, D. A. Young, E. M. Shevach, M. Collins, and M. C. Byrne. 2002. CD4⁺CD25⁺ immunoregulatory T cells: gene expression analysis reveals a functional role for the glucocorticoid-induced TNF receptor. *Immunity* 16:311.
- Ji, H. B., G. Liao, W. A. Faubion, A. C. Abadia-Molina, C. Cozzo, F. S. Laroux, A. Caton, and C. Terhorst. 2004. Cutting edge: the natural ligand for glucocorticoid-induced TNF receptor-related protein abrogates regulatory T cell suppression. *J. Immunol.* 172:5823.
- Lord, S. J., R. V. Rajotte, G. S. Korbutt, and R. C. Bleackley. 2003. Granzyme B: a natural born killer. *Immunol. Rev.* 193:31.
- Appay, V., J. J. Zaunders, L. Papagno, J. Sutton, A. Jaramillo, A. Waters, P. Easterbrook, P. Grey, D. Smith, A. J. McMichael, et al. 2002. Characterization of CD4⁺ CTLs ex vivo. *J. Immunol.* 168:5954.
- van Leeuwen, E. M., E. B. Remmerswaal, M. T. Vossen, A. T. Rowshani, P. M. Wertheim-van Dillen, R. A. van Lier, and I. J. ten Berge. 2004. Emergence of a CD4⁺CD28⁻ granzyme B⁺, cytomegalovirus-specific T cell subset after recovery of primary cytomegalovirus infection. *J. Immunol.* 173:1834.
- Kemper, C., A. C. Chan, J. M. Green, K. A. Brett, K. M. Murphy, and J. P. Atkinson. 2003. Activation of human CD4⁺ cells with CD3 and CD46 induces a T-regulatory cell 1 phenotype. *Nature* 421:388.
- Grossman, W. J., J. W. Verbsky, B. L. Tollefsen, C. Kemper, J. P. Atkinson, and T. J. Ley. 2004. Differential expression of granzymes A and B in human cytotoxic lymphocyte subsets and T regulatory cells. *Blood* 104:2840.
- Grossman, W. J., J. W. Verbsky, W. Barchet, M. Colonna, J. P. Atkinson, and T. J. Ley. 2004. Human T regulatory cells can use the perforin pathway to cause autologous target cell death. *Immunity* 21:589.
- Herman, A. E., G. J. Freeman, D. Mathis, and C. Benoist. 2004. CD4⁺CD25⁺ T regulatory cells dependent on ICOS promote regulation of effector cells in the prediabetic lesion. *J. Exp. Med.* 199:1479.
- Froelich, C. J., K. Orth, J. Turbov, P. Seth, R. Gottlieb, B. Babiorek, G. M. Shah, R. C. Bleackley, V. M. Dixit, and W. Hanna. 1996. New paradigm for lymphocyte granule-mediated cytotoxicity: target cells bind and internalize granzyme B, but an endosomal agent is necessary for cytosolic delivery and subsequent apoptosis. *J. Biol. Chem.* 271:29073.
- Browne, K. A., E. Blink, V. R. Sutton, C. J. Froelich, D. A. Jans, and J. A. Trapani. 1999. Cytosolic delivery of granzyme B by bacterial toxins: evidence that endosomal disruption, in addition to transmembrane pore formation, is an important function of perforin. *Mol. Cell Biol.* 19:8604.
- Trapani, J. A., V. R. Sutton, K. Y. Thia, Y. Q. Li, C. J. Froelich, D. A. Jans, M. S. Sandrin, and K. A. Browne. 2003. A clathrin/dynamin- and mannose-6-phosphate receptor-independent pathway for granzyme B-induced cell death. *J. Cell Biol.* 160:223.
- Kurschus, F. C., M. Kleinschmidt, E. Fellows, K. Dornmair, R. Rudolph, H. Lilie, and D. E. Jenne. 2004. Killing of target cells by redirected granzyme B in the absence of perforin. *FEBS Lett.* 562:87.
- Choy, J. C., V. H. Hung, A. L. Hunter, P. K. Cheung, B. Motyka, I. S. Goping, T. Sawchuk, R. C. Bleackley, T. J. Podor, B. M. McManus, and D. J. Granville. 2004. Granzyme B induces smooth muscle cell apoptosis in the absence of perforin: involvement of extracellular matrix degradation. *Arterioscler. Thromb. Vasc. Biol.* 24:2245.
- Somersalo, K., N. Anikeeva, T. N. Sims, V. K. Thomas, R. K. Strong, T. Spies, T. Lebedeva, Y. Sykulev, and M. L. Dustin. 2004. Cytotoxic T lymphocytes form an antigen-independent ring junction. *J. Clin. Invest.* 113:49.
- van Kooyk, Y., P. Weder, F. Hogervorst, A. J. Verhoeven, G. van Seventer, A. A. te Velde, J. Borst, G. D. Keizer, and C. G. Figdor. 1991. Activation of LFA-1 through a Ca²⁺-dependent epitope stimulates lymphocyte adhesion. *J. Cell Biol.* 112:345.
- Labadia, M. E., D. D. Jeanfavre, G. O. Caviness, and M. M. Morelock. 1998. Molecular regulation of the interaction between leukocyte function-associated antigen-1 and soluble ICAM-1 by divalent metal cations. *J. Immunol.* 161:836.
- Dai, Z., Q. Li, Y. Wang, G. Gao, L. S. Diggs, G. Tellides, and F. G. Lakkis. 2004. CD4⁺CD25⁺ regulatory T cells suppress allograft rejection mediated by memory CD8⁺ T cells via a CD30-dependent mechanism. *J. Clin. Invest.* 113:310.
- Takahashi, T., Y. Kuniyasu, M. Toda, N. Sakaguchi, M. Itoh, M. Iwata, J. Shimizu, and S. Sakaguchi. 1998. Immunologic self-tolerance maintained by CD25⁺CD4⁺ naturally anergic and suppressive T cells: induction of autoimmune disease by breaking their anergic/suppressive state. *Int. Immunol.* 10:1969.