



Explore what's possible with innovative
research tools

Discover the difference >



In Vivo Generation of Pathogen-Specific Th1 Cells in the Absence of the IFN- γ Receptor

Jodie S. Haring, Vladimir P. Badovinac, Matthew R. Olson,
Steven M. Varga and John T. Harty

This information is current as
of September 18, 2021.

J Immunol 2005; 175:3117-3122; ;
doi: 10.4049/jimmunol.175.5.3117
<http://www.jimmunol.org/content/175/5/3117>

References This article **cites 31 articles**, 16 of which you can access for free at:
<http://www.jimmunol.org/content/175/5/3117.full#ref-list-1>

Why *The JI*? Submit online.

- **Rapid Reviews! 30 days*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

**average*

Subscription Information about subscribing to *The Journal of Immunology* is online at:
<http://jimmunol.org/subscription>

Permissions Submit copyright permission requests at:
<http://www.aai.org/About/Publications/JI/copyright.html>

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://jimmunol.org/alerts>

The Journal of Immunology is published twice each month by
The American Association of Immunologists, Inc.,
1451 Rockville Pike, Suite 650, Rockville, MD 20852
Copyright © 2005 by The American Association of
Immunologists All rights reserved.
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



In Vivo Generation of Pathogen-Specific Th1 Cells in the Absence of the IFN- γ Receptor¹

Jodie S. Haring,* Vladimir P. Badovinac,* Matthew R. Olson,* Steven M. Varga,* and John T. Harty^{2*†}

The precise mechanisms that govern the commitment of CD4 T cells to become Th1 or Th2 cells in vivo are incompletely understood. Recent experiments demonstrate colocalization of the IFN- γ R chains with the TCR during activation of naive CD4 T cells, suggesting that association of these molecules may be involved in determining lineage commitment. To test the role of IFN- γ and its receptor in the generation of Th1 Ag-specific CD4 T cells, we analyzed mice after infection with *Listeria monocytogenes* or lymphocytic choriomeningitis virus. In the absence of IFN- γ , Ag-specific CD4 T cells were generated in response to both these infections. In addition, IFN- γ -producing (Th1) Ag-specific CD4 T cells were generated in mice lacking the ligand-binding chain of the IFN- γ R (IFN- γ R1^{-/-}) or the signaling chain (IFN- γ R2^{-/-}). There was no increase in the number of IL-4-producing Ag-specific CD4 T cells, nor was there a decrease in the expression of T-bet in the absence of functional IFN- γ signaling, indicating that the cells were committed Th1 cells. Thus, both chains of the IFN- γ R are dispensable for the generation of Th1 Ag-specific CD4 T cells after infection in vivo. *The Journal of Immunology*, 2005, 175: 3117–3122.

Early during an immune response naive CD4 T cells commit themselves to either a Th1 or a Th2 developmental pathway. This decision is strongly influenced by the cytokines present in the localized environment during activation of the CD4 T cells. In particular, IL-12 and IFN- γ have been identified as key regulators of Th1 development, and IL-4 is considered the key regulator of Th2 development (1). The lack of proper lineage development can adversely affect the outcome of infection, which is most clearly observed in C57BL/6 (B6) and BALB/c mice infected with the parasite *Leishmania major*. B6 mice develop a protective Th1 response and clear the infection, whereas BALB/c mice mount a nonprotective Th2 response, fail to control the infection, and develop progressive disease (2).

Most of the data addressing the mechanisms underlying Th1-Th2 lineage commitment by CD4 T cells have been acquired using in vitro experiments that involve culturing of cells in specific cytokine environments. It has been recently demonstrated that CD4 T cells cultured under Th1 conditions express the transcription factor T-bet (3, 4), whereas CD4 T cells cultured under Th2 conditions express the transcription factor GATA-3 (5, 6). The expression of these transcription factors is extremely specific for each lineage. Forced expression of T-bet can make Th2 CD4 T cells produce IFN- γ , a Th1-specific cytokine (3).

It has long been observed that the cytokines that foster the development of Th1 cells inhibit the development of Th2 cells and vice versa (7, 8); however, the exact mechanism(s) by which this is accomplished is currently unknown. A recent report (9) dem-

onstrated that both chains of the IFN- γ R colocalized with the TCR during activation of naive CD4 T cells in vitro with either Ag-expressing APCs or TCR cross-linking with Ab. In addition, the authors demonstrated that the TCR-IFN- γ R colocalization did not occur in the presence of IL-4. These data suggest the intriguing hypotheses that TCR-IFN- γ R colocalization could be important for the generation of Th1 CD4 T cells and that inhibition of this process could be the mechanism by which the Th2-inducing cytokine IL-4 inhibited the development of Th1 CD4 T cells.

To address the potential requirement for TCR-IFN- γ R colocalization in Th1 CD4 T cell generation in vivo, we infected wild-type (wt),³ IFN- γ ^{-/-}, IFN- γ R1^{-/-}, and IFN- γ R2^{-/-} mice with *Listeria monocytogenes* (LM) or lymphocytic choriomeningitis virus (LCMV), both of which induce vigorous CD4 T cell responses to well-characterized Ags (10–12). Our data demonstrate that Th1 Ag-specific CD4 T cells can be generated in the absence of either chain of the IFN- γ R, indicating that colocalization of the TCR with an intact IFN- γ R is not required for the generation of Th1-committed Ag-specific CD4 T cells.

Materials and Methods

Mice

C57BL/6 mice (National Cancer Institute), 129SVE mice (Taconic Farms), and IFN- γ ^{-/-} and IFN- γ R1^{-/-} mice (B6 background; The Jackson Laboratory) were purchased for use in these experiments. IFN- γ R2^{-/-} mice (129SVE background) (13) were generously provided by Dr. P. Rothman (University of Iowa, Iowa City, IA). All animal experiments followed approved institutional animal care and use committee protocols.

Bacteria and viruses

LM expressing OVA was a gift from Dr. L. Lefrancois (University of Connecticut, Farmington, CT) (14). An attenuated version of this strain was created by introducing an in-frame deletion in the *actA* gene as previously described (15). Bacteria were grown and quantified as previously described (16, 17). All bacterial infections were via i.v. injection. The Armstrong strain of LCMV was propagated and titrated as previously described (18). Approximately 2×10^5 PFU was given to each mouse i.p.

*Department of Microbiology and [†]Interdisciplinary Program in Immunology, University of Iowa, Iowa City, IA 52242

Received for publication February 15, 2005. Accepted for publication June 29, 2005.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This work was supported by National Institutes of Health Grants AI42767, AI46653, and AI50073 (to J.T.H.); T32AI0726-19 (to J.S.H.); and 2T32AI0075-33 (to M.R.O.).

² Address correspondence and reprint requests to Dr. John T. Harty, Department of Microbiology, 3-512 BSB, 51 Newton Road, Iowa City, IA 52242. E-mail address: john-harty@uiowa.edu

³ Abbreviations used in this paper: wt, wild type; ICS, intracellular cytokine staining; LCMV, lymphocytic choriomeningitis virus; LLO, listeriolysin O; LM, *Listeria monocytogenes*; NP, nuclear protein; p.i., postinfection.

Intracellular cytokine staining (ICS) and quantification of Ag-specific CD4 T cells

Surface staining for CD4 (L3T4; BD Pharmingen) and Thy1.2 (53-2.1; BD Pharmingen) and intracellular staining for IFN- γ (XMG1.2; BD Pharmingen), TNF (MP6-XT22; e-Bioscience), IL-4 (11B11; BD Pharmingen), or IL-5 (TRFK5; BD Pharmingen) were performed as previously described after incubation with listeriolysin O (LLO)_{190–201}, gp61–80, nuclear protein (NP)_{309–328}, or no peptide (19). The total number of Ag-specific CD4 T cells per spleen was calculated by multiplying the frequency of CD4⁺/Thy1.2⁺/IFN- γ ⁺ or TNF⁺ cells after stimulation with specific peptide by the total number of splenocytes.

ELISPOT

The ELISPOT assay was performed as previously described (20). Briefly, microtiter plates were coated with anti-IL-4 (clone 11B11; eBioscience). Whole splenocytes (5×10^6 to 2.5×10^5) from wt 129SVE and IFN- γ R2^{-/-} mice on day 7 postinfection (p.i.) were plated in triplicate together with 1×10^6 irradiated splenocytes pulsed with 1 mM LLO_{190–201} or nonpulsed splenocytes. Cells were cultured for 41 h before the biotinylated anti-IL-4 capture Ab (clone BVD6-24G2; eBioscience) was added. The assay was developed using the substrate 3-amino-9-ethylcarbazole. The number of spots per 1×10^6 CD4 T cells was calculated using the frequency of CD4⁺ cells determined by FACS. The plates were analyzed using an immunospot analyzer (Cellular Technology) according to the manufacturer's instructions.

CD4 T cell purification and RT-PCR

Splenocytes were stained with PE-conjugated CD4 (clone L3T4), then labeled with anti-PE-coated magnetic beads according to manufacturer's instructions (Miltenyi Biotec). Labeled CD4 T cells were then recovered by AutoMACS separation (Posseld program). Purity was assessed by FACS analysis before RNA isolation. All cell samples were purified to >90% CD4⁺. RNA was isolated from purified T cells using the RNeasy Mini Kit with additional on-column DNase treatment according to the manufacturer's instructions (Qiagen). cDNA was synthesized using a reaction mix including random hexamers and Moloney murine leukemia virus reverse transcriptase.

Amplification of cDNA for T-bet and GATA-3 from purified CD4 T cells was performed using RT-PCR as previously described (21). GAPDH was amplified together with each target gene and was used to normalize expression levels. Appropriate bands on ethidium bromide-stained gels were quantitated using ImageQuant 3.3. Oligos used in these experiments were (5'-3'): T-bet forward, GCGCCAGGAAGTTTCATTTGGGAA; T-bet reverse, ACAGCTCGGAAGTCCGCTTCATAA; GATA-3 forward, AGAAAGAAGGCATCCAGACCCGAA; and GATA-3 reverse, AGGACCTCTTCGCACACTTGGAGA. GAPDH primers were part of the TaqMan Rodent GAPDH Control Reagents kit (Applied Biosystems).

Results

IFN- γ can up-regulate the autocrine production of IFN- γ by DCs (22) and is considered a major effector cytokine of Th1 CD4 T cells (23). To determine the requirement for IFN- γ in the generation of Ag-specific CD4 T cells, wt B6 and IFN- γ R1^{-/-} B6 mice were infected with ~ 0.1 LD₅₀ of an attenuated *actA*-deficient strain of LM that is cleared from both strains by day 5 p.i. (data not shown) (16). CD4 T cells specific for LLO_{190–201}, an I-A^b-restricted CD4 T cell epitope, were detected by peptide stimulation and ICS for TNF. As shown in Fig. 1, TNF-producing LLO_{190–201}-specific CD4 T cells were detected in both wt B6 mice (Fig. 1A) and IFN- γ R1^{-/-} B6 mice (Fig. 1B) on day 7 p.i. When total numbers of LLO_{190–201}-specific CD4 T cells per spleen were calculated (Fig. 1C), it was determined that the IFN- γ R1^{-/-} B6 mice were capable of generating a slightly larger Ag-specific CD4 T cell response than the wt B6 mice. Total numbers of splenocytes were not different between the two groups of mice (wt B6, $1.45 \times 10^8 \pm 2.85 \times 10^7$; IFN- γ R1^{-/-}, $1.42 \times 10^8 \pm 3.40 \times 10^7$; $p = 0.86$). These data demonstrate that IFN- γ is not required for the generation of Ag-specific CD4 T cells after infection with LM.

The IFN- γ R is composed of two unique chains, IFN- γ R1 and IFN- γ R2 (24, 25). Both chains are required for responsiveness to




FIGURE 1. Ag-specific CD4 T cell generation in the absence of IFN- γ . IFN- γ R1^{-/-} (B6 background) and wt B6 mice were infected with LM. A and B, ICS for TNF was performed on splenocytes harvested from wt B6 (A) and IFN- γ R1^{-/-} (B) mice on day 7 p.i. after a 6-h in vitro stimulation with LLO_{190–201} (top panels) or no peptide (bottom panels). FACS plots were first gated on CD4⁺ cells and are representative of three experiments. Numbers indicate the frequency of Thy1.2⁺/TNF⁺ cells within the CD4⁺ population. C, Total number of Ag-specific CD4 T cells per spleen. Statistical analysis was performed using Student's *t* test. *, $p < 0.05$.

IFN- γ (26). IFN- γ R1 is the ligand-binding portion of the receptor and is constitutively expressed on most cells (27), including naive T cells (28). A recent report demonstrated that IFN- γ R1 colocalized with the TCR during, but not before, naive CD4 T cell activation in vitro (9). Colocalization of IFN- γ R1 with the TCR was abrogated in the presence of IL-4 via a process dependent on STAT6, which is a major signaling component directly downstream of the IL-4R. Thus, IFN- γ R recruitment to the TCR could potentially serve an important function in Th1 lineage commitment by CD4 T cells (9). To test the requirement for IFN- γ R1 colocalization with the TCR for the generation of Th1 Ag-specific CD4 T cells in vivo, we infected wt and IFN- γ R1^{-/-} mice with *actA*⁻ LM and analyzed splenocytes from these mice on day 7 p.i. for the presence of IFN- γ -producing (Th1) LLO_{190–201}-specific CD4 T cells. IFN- γ R1^{-/-} mice cleared the *actA*⁻ LM infection by day 5 p.i. (data not shown). As shown in Fig. 2, substantial expansion of IFN- γ -producing LLO_{190–201}-specific CD4 T cells was detected on day 7 p.i. in both wt B6 and IFN- γ R1^{-/-} mice, which resulted in similar frequencies (Fig. 2, A and B) and a <2-fold difference in total numbers of Ag-specific CD4 T cells (Fig. 2C). Total numbers of splenocytes were not different between the two groups of mice




FIGURE 2. IFN- γ R1 is not required for the generation of Th1 Ag-specific CD4 T cells after infection. Wt B6 and IFN- γ R1 $^{-/-}$ mice were infected with LM. *A* and *B*, ICS for IFN- γ was performed on splenocytes harvested from infected wt (*A*) and IFN- γ R1 $^{-/-}$ mice (*B*) on day 7 p.i. after incubation with LLO_{190–201} (*top panels*) or no peptide (*bottom panels*). The FACS plots presented were first gated on CD4 $^{+}$ cells and are representative of three mice per group. Numbers indicate the frequency of Thy1.2 $^{+}$ /IFN- γ $^{+}$ cells within the CD4 $^{+}$ population. *C*, Total number of Th1 Ag-specific CD4 T cells per spleen. Statistical analysis was performed using Student's *t* test. *, $p < 0.05$.

(wt B6, $1.45 \times 10^8 \pm 2.85 \times 10^7$; IFN- γ R1 $^{-/-}$ mice, $1.40 \times 10^8 \pm 2.56 \times 10^7$; $p = 0.72$). These data indicate that in the absence of IFN- γ R1 $^{-/-}$, and therefore colocalization of this cytokine receptor component with the TCR during T cell activation, Ag-specific CD4 T cells capable of producing IFN- γ (probably Th1 cells) are generated after infection with LM.

IFN- γ R2 has a very short extracellular domain and does not bind IFN- γ directly. It is not normally preassociated with IFN- γ R1, but instead complexes with IFN- γ R1 upon ligand binding and is required for subsequent signaling (24, 25). IFN- γ R2 was also shown to colocalize with the TCR during naive CD4 T cell activation (9). To directly test a role for IFN- γ R2 in the generation of Th1 Ag-specific CD4 T cells in vivo, we infected wt 129SVE and IFN- γ R2 $^{-/-}$ mice (129SVE background) (13) with LM and measured the LLO_{190–201}-specific CD4 T cell response on day 7 p.i. by ICS for IFN- γ . IFN- γ R2 $^{-/-}$ and wt 129 SVE mice cleared the *actA* $^{-}$ LM infection by day 5 p.i. (data not shown). The ICS data indicate that IFN- γ -producing (Th1) LLO_{190–201}-specific

CD4 T cells were generated with similar frequencies in wt and IFN- γ R2 $^{-/-}$ mice (Fig. 3, *A* and *B*). When total numbers of LLO_{190–201}-specific CD4 T cells per spleen were calculated, the data showed that IFN- γ R2 $^{-/-}$ mice had approximately twice as many Th1 Ag-specific CD4 T cells on day 7 p.i. compared with wt 129 mice. The total numbers of splenocytes were not different between the two groups of mice (wt 129 mice, $1.15 \times 10^8 \pm 1.84 \times 10^7$; IFN- γ R2 $^{-/-}$ mice, $1.30 \times 10^8 \pm 2.66 \times 10^7$; $p = 0.17$). In combination, the data presented above indicate that neither IFN- γ R1 nor IFN- γ R2 is required for generation of Ag-specific CD4 T cells capable of making IFN- γ , and, therefore, probably Th1 cells, after infection with LM.

To ensure that these results were not peculiar to LM infection or the LLO_{190–201} epitope, we infected wt B6, IFN- γ R1 $^{-/-}$, and IFN- γ R2 $^{-/-}$ mice with the arenavirus LCMV Armstrong strain and measured IFN- γ -producing Th1 CD4 T cells specific for gp61–80 and NP_{309–328} on day 8 p.i. (Fig. 4*A*). IFN- γ -producing CD4 T cells specific for each Ag were generated with similar frequencies (Fig. 4*A*) and total numbers (Fig. 4, *B* and *C*) in the




FIGURE 3. IFN- γ R2 is not required for the generation of Th1 Ag-specific CD4 T cells after infection. Wt 129SVE and IFN- γ R2 $^{-/-}$ mice were infected with LM. *A* and *B*, ICS for IFN- γ was performed on splenocytes harvested from infected wt (*A*) and IFN- γ R2 $^{-/-}$ (*B*) mice on day 7 p.i. after incubation with LLO_{190–201} (*top panels*) or no peptide (*bottom panels*). FACS plots presented were all first gated on CD4 $^{+}$ cells and are representative of at least three mice per group. Numbers indicate the frequency of Thy1.2 $^{+}$ /IFN- γ $^{+}$ cells within the CD4 $^{+}$ population. *C*, Total number of Th1 Ag-specific CD4 T cells per spleen. Statistical analysis was performed using Student's *t* test. *, $p < 0.05$.




FIGURE 4. Th1 Ag-specific CD4 T cells are generated in the absence of IFN- γ R1 or IFN- γ R2 after viral infection. Wt B6, IFN- γ R1^{-/-}, and IFN- γ R2^{-/-} mice were infected with LCMV. *A*, ICS for IFN- γ was performed on splenocytes harvested on day 8 p.i. after incubation with gp61–80 (*top panels*), NP_{309–328} (*middle panels*), or no peptide (*bottom panels*). The FACS plots presented were first gated on CD4⁺ cells and are representative of at least three mice per group. Numbers indicate frequencies of Thy1.2⁺/IFN- γ ⁺ within the CD4⁺ population. *B*, Total number of Th1 gp61–80-specific CD4 T cells per spleen. *C*, Total number of Th1 NP_{309–328}-specific CD4 T cells per spleen. Statistical analysis was performed using Student's *t* test. *, *p* < 0.05.

different groups of mice. It has been previously reported that wt 129 mice clear LCMV Armstrong infection on approximately day 8 p.i. In contrast, there was prolonged infection in IFN- γ R2^{-/-} mice, with clearance reported to be between days 10 and 15 p.i., which resulted in increased total numbers of CD8 T cells at these time points (29). Consistent with these results and as was observed after LM infection, IFN- γ R2^{-/-} mice consistently exhibited higher numbers of Th1 Ag-specific CD4 T cells compared with wt 129 mice (Figs. 3, *B* and *C*, and 4). We also infected IFN- γ ^{-/-} mice with LCMV and used ICS for TNF to detect gp61–80- and NP_{309–328}-specific CD4 T cells. We were able to detect CD4 T

cells specific for each of these epitopes on day 8 p.i. (data not shown), indicating that IFN- γ is also not required for the generation of Ag-specific CD4 T cells after infection with LCMV.

Although the ability to produce IFN- γ is a hallmark characteristic of Th1 CD4 T cells, it remained possible that Ag-specific CD4 T cells generated after LM or LCMV infection in the absence of the IFN- γ R were uncommitted to either a Th1 or Th2 lineage. To formally show that these cells were Th1 Ag-specific CD4 T cells after infection and did not also possess characteristics of Th2 cells, we stimulated splenocytes from IFN- γ R2^{-/-} and wt 129SVE mice on day 7 p.i. with LLO_{190–201} and costained for IFN- γ and either IL-4 or IL-5. As shown in Fig. 5, *A* and *B*, we were unable to detect any IFN- γ -producing LLO_{190–201}-specific CD4 T cells that also produced IL-4 or IL-5 in either wt 129SVE or IFN- γ R2^{-/-} mice. In addition, using a more sensitive ELISPOT assay, we determined that the number of IL-4-secreting LLO_{190–201}-specific CD4 T cells was not statistically different between wt 129SVE mice and IFN- γ R2^{-/-} mice (Fig. 5*C*). In summary, we did not detect uncommitted Ag-specific CD4 T cells in IFN- γ R2^{-/-} mice, nor did we document any increase in the number of Ag-specific Th2 CD4 T cells.

To further investigate Th1 and Th2 characteristics of CD4 T cells after LM infection, we purified CD4 T cells from wt 129SVE and IFN- γ R2^{-/-} mice on day 7 p.i. and performed RT-PCR for the transcription factors T-bet and GATA-3 (Fig. 5, *D* and *E*). Compared with CD4 T cells isolated from wt mice, IFN- γ R2^{-/-} CD4 T cells did not express statistically less T-bet or more GATA-3, indicating that they were not more Th2-like than wt CD4 T cells. Taken together, these data strongly support the conclusion that Th1-committed Ag-specific CD4 T cells are generated in the absence of the IFN- γ R.

Discussion

Using two different infectious model systems, we have shown that the development of Th1 Ag-specific CD4 T cells does not require either of the IFN- γ R chains. It is a strong likelihood that during the process of Th1-Th2 lineage commitment, when multiple signals are being received by naive CD4 T cells in a short period of time, ligation of one receptor may influence how signals received through other receptors are translated into cellular responses. These interactions may be amplified under *in vitro* stimulation conditions, when the full spectrum of inflammatory signals may not be present.

There are statistical differences in the magnitudes of the Th1 Ag-specific CD4 T cell responses on day 7 (LM) and day 8 (LCMV) p.i. between wt and the various knockout mice. However, without thorough kinetic experiments and determination of the peak of the CD4 T cell response in IFN- γ and receptor-deficient mice, statistical differences in the numbers of cells at this one time point may be misleading. We sought to determine whether Th1 Ag-specific CD4 T cells could be primed in the absence of IFN- γ or its receptor. Our results clearly show this to be true, although the number of IFN- γ -producing CD4 T cells was not always equivalent. As we have previously documented, CD8 T cell responses in the absence of IFN- γ exhibit different kinetics than those in wt mice (18). We have generated preliminary data to suggest that Ag-specific CD4 T cell response kinetics are also aberrant in IFN- γ -deficient mice. Studies are currently underway to carefully explore interesting differences in the magnitude and kinetics of Ag-specific CD4 T cell responses in the absence of IFN- γ or its receptor.

It remains a possibility that the IFN- γ R participates together with TCR signaling in Th1 lineage commitment of CD4 T cells




FIGURE 5. Ag-specific CD4 T cells generated after LM infection are committed Th1 cells. CD4 cells from wt 129SVE and $\text{IFN-}\gamma\text{R2}^{-/-}$ mice on day 7 p.i. with LM were analyzed for their ability to produce Th2 cytokines and expression of Th1- or Th2-specific transcription factors. ICS for $\text{IFN-}\gamma$ and IL-4 (A) or IL-5 (B) was performed after incubation with $\text{LLO}_{190-201}$ (top panels) or no peptide (bottom panels). Numbers indicate the frequency of $\text{IFN-}\gamma^{+}/\text{IL-4}^{-}$ or IL-5^{+} or $\text{IFN-}\gamma^{+}/\text{IL-4}^{+}$ or IL-5^{-} cells within the $\text{Thy1.2}^{+}/\text{CD4}^{+}$ population. Data are representative of three mice per group. C, Results of an IL-4 ELISPOT assay reported as the number of IL-4-producing spots per 1×10^6 CD4 cells. Cells were either stimulated for 41 h with $\text{LLO}_{190-201}$ or left unstimulated. The number of spots detected in the unstimulated wells was subtracted from the number detected in the peptide-stimulated wells. Data are the mean \pm SD of three mice per group. Statistical analysis was performed using Student's *t* test. D and E, Total CD4 T cells were purified from the indicated mice on day 7 p.i. RT-PCR was performed for T-bet (D) and GATA-3 (E). Data are reported as a ratio of T-bet or GATA-3 expression to GAPDH expression. Each symbol represents one mouse. Lines indicate averages per group.

under some conditions. Perhaps during infections where IL-4 is produced in much higher quantities than after LM or LCMV infections, the inhibition of $\text{IFN-}\gamma\text{R}$ colocalization with the TCR may negatively influence Th1 polarization of CD4 T cells, resulting in the development of Th2 Ag-specific CD4 T cells as is seen in *L. major* infection of BALB/c mice (30, 31). However, our experiments clearly demonstrate that after infection with LM or LCMV, signaling through the $\text{IFN-}\gamma\text{R}$ is not a requirement for the development of Th1 CD4 T cells in vivo.

Acknowledgments

We thank Rebecca Podyminogin for excellent laboratory assistance.

Disclosures

The authors have no financial conflict of interest.

References

- Ho, I. C., and L. H. Glimcher. 2002. Transcription: tantalizing times for T cells. *Cell* 109(Suppl.): S109–S120.
- Sacks, D., and N. Noben-Trauth. 2002. The immunology of susceptibility and resistance to *Leishmania major* in mice. *Nat. Rev. Immunol.* 2: 845–858.
- Szabo, S. J., S. T. Kim, G. L. Costa, X. Zhang, C. G. Fathman, and L. H. Glimcher. 2000. A novel transcription factor, T-bet, directs Th1 lineage commitment. *Cell* 100: 655–669.
- Szabo, S. J., B. M. Sullivan, C. Stemmann, A. R. Satoskar, B. P. Sleckman, and L. H. Glimcher. 2002. Distinct effects of T-bet in Th1 lineage commitment and $\text{IFN-}\gamma$ production in CD4 and CD8 T cells. *Science* 295: 338–342.
- Zhang, D. H., L. Cohn, P. Ray, K. Bottomly, and A. Ray. 1997. Transcription factor GATA-3 is differentially expressed in murine Th1 and Th2 cells and controls Th2-specific expression of the interleukin-5 gene. *J. Biol. Chem.* 272: 21597–21603.
- Zheng, W., and R. A. Flavell. 1997. The transcription factor GATA-3 is necessary and sufficient for Th2 cytokine gene expression in CD4 T cells. *Cell* 89: 587–596.
- O'Garra, A., and K. Murphy. 1996. Role of cytokines in development of Th1 and Th2 cells. *Chem. Immunol.* 63: 1–13.
- O'Garra, A., and K. Murphy. 1994. Role of cytokines in determining T-lymphocyte function. *Curr. Opin. Immunol.* 6: 458–466.
- Maldonado, R. A., D. J. Irvine, R. Schreiber, and L. H. Glimcher. 2004. A role for the immunological synapse in lineage commitment of CD4 lymphocytes. *Nature* 431: 527–532.
- Corbin, G. A., and J. T. Harty. 2004. Duration of infection and antigen display have minimal influence on the kinetics of the CD4^{+} T cell response to *Listeria monocytogenes* infection. *J. Immunol.* 173: 5679–5687.
- Geginat, G., S. Schenk, M. Skoberne, W. Goebel, and H. Hof. 2001. A novel approach of direct ex vivo epitope mapping identifies dominant and subdominant CD4 and CD8 T cell epitopes from *Listeria monocytogenes*. *J. Immunol.* 166: 1877–1884.
- Varga, S. M., and R. M. Welsh. 2000. High frequency of virus-specific interleukin-2-producing CD4^{+} T cells and Th1 dominance during lymphocytic choriomeningitis virus infection. *J. Virol.* 74: 4429–4432.
- Lu, B., C. Ebersperger, Z. Dembic, Y. Wang, M. Kvatyuk, T. Lu, R. L. Coffman, S. Pestka, and P. B. Rothman. 1998. Targeted disruption of the interferon- γ receptor 2 gene results in severe immune defects in mice. *Proc. Natl. Acad. Sci. USA* 95: 8233–8238.
- Pope, C., S. K. Kim, A. Marzo, D. Masopust, K. Williams, J. Jiang, H. Shen, and L. Lefrancois. 2001. Organ-specific regulation of the CD8 T cell response to *Listeria monocytogenes* infection. [Published erratum appears in *J. Immunol.* 2001 166: 5840.] *J. Immunol.* 166: 3402–3409.
- Tvinnereim, A. R., S. E. Hamilton, and J. T. Harty. 2002. CD8^{+} -T-cell response to secreted and nonsecreted antigens delivered by recombinant *Listeria monocytogenes* during secondary infection. *Infect. Immun.* 70: 153–162.
- Harty, J. T., and M. J. Bevan. 1995. Specific immunity to *Listeria monocytogenes* in the absence of $\text{IFN-}\gamma$. *Immunity* 3: 109–117.
- Harty, J. T., and M. J. Bevan. 1992. CD8^{+} T cells specific for a single nonamer epitope of *Listeria monocytogenes* are protective in vivo. *J. Exp. Med.* 175: 1531–1538.
- Badovinac, V. P., A. R. Tvinnereim, and J. T. Harty. 2000. Regulation of antigen-specific CD8^{+} T cell homeostasis by perforin and $\text{IFN-}\gamma$. *Science* 290: 1354–1358.
- Badovinac, V. P., and J. T. Harty. 2000. Intracellular staining for TNF and $\text{IFN-}\gamma$ detects different frequencies of antigen-specific CD8^{+} T cells. *J. Immunol. Methods* 238: 107–117.
- Varga, S. M., E. L. Wissinger, and T. J. Braciale. 2000. The attachment (G) glycoprotein of respiratory syncytial virus contains a single immunodominant epitope that elicits both Th1 and Th2 CD4^{+} T cell responses. *J. Immunol.* 165: 6487–6495.

21. Haring, J. S., G. A. Corbin, and J. T. Harty. 2005. Dynamic regulation of IFN- γ signaling in antigen-specific CD8⁺ T cells responding to infection. *J. Immunol.* 174: 6791–6802.
22. Lugo-Villarino, G., R. Maldonado-Lopez, R. Possemato, C. Penaranda, and L. H. Glimcher. 2003. T-bet is required for optimal production of IFN- γ and antigen-specific T cell activation by dendritic cells. *Proc. Natl. Acad. Sci. USA* 100: 7749–7754.
23. Murphy, K. M., and S. L. Reiner. 2002. The lineage decisions of helper T cells. *Nat. Rev. Immunol.* 2: 933–944.
24. Boehm, U., T. Klamp, M. Groot, and J. C. Howard. 1997. Cellular responses to interferon- γ . *Annu. Rev. Immunol.* 15: 749–795.
25. Bach, E. A., M. Aguet, and R. D. Schreiber. 1997. The IFN γ receptor: a paradigm for cytokine receptor signaling. *Annu. Rev. Immunol.* 15: 563–591.
26. Pernis, A., S. Gupta, K. J. Gollob, E. Garfein, R. L. Coffman, C. Schindler, and P. Rothman. 1995. Lack of interferon γ receptor β chain and the prevention of interferon γ signaling in TH1 cells. *Science* 269: 245–247.
27. Farrar, M. A., and R. D. Schreiber. 1993. The molecular cell biology of interferon- γ and its receptor. *Annu. Rev. Immunol.* 11: 571–611.
28. Skrenta, H., Y. Yang, S. Pestka, and C. G. Fathman. 2000. Ligand-independent down-regulation of IFN- γ receptor 1 following TCR engagement. *J. Immunol.* 164: 3506–3511.
29. Lohman, B. L., and R. M. Welsh. 1998. Apoptotic regulation of T cells and absence of immune deficiency in virus-infected γ interferon receptor knockout mice. *J. Virol.* 72: 7815–7821.
30. Stetson, D. B., M. Mohrs, V. Mallet-Designe, L. Teyton, and R. M. Locksley. 2002. Rapid expansion and IL-4 expression by *Leishmania*-specific naive helper T cells in vivo. *Immunity* 17: 191–200.
31. Launois, P., I. Maillard, S. Pingel, K. G. Swihart, I. Xenarios, H. Acha-Orbea, H. Diggelmann, R. M. Locksley, H. R. MacDonald, and J. A. Louis. 1997. IL-4 rapidly produced by Vb4/Va8 CD4⁺ T cells instructs Th2 development and susceptibility to *Leishmania major* in BALB/c mice. *Immunity* 6: 541–549.