



## Two data sets, one step, zero doubt

New Attune™ CytPix™ Flow Cytometer

[See it now >](#)

**invitrogen**  
by Thermo Fisher Scientific



## Cutting Edge: Resistance to HIV-1 Infection among African Female Sex Workers Is Associated with Inhibitory KIR in the Absence of Their HLA Ligands

This information is current as of January 22, 2022.

Wim Jennes, Sonja Verheyden, Christian Demanet, Christiane A. Adjé-Touré, Bea Vuylsteke, John N. Nkengasong and Luc Kestens

*J Immunol* 2006; 177:6588-6592; ;

doi: 10.4049/jimmunol.177.10.6588

<http://www.jimmunol.org/content/177/10/6588>

**References** This article **cites 26 articles**, 8 of which you can access for free at: <http://www.jimmunol.org/content/177/10/6588.full#ref-list-1>

**Why *The JI*? Submit online.**

- **Rapid Reviews! 30 days\*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

*\*average*

**Subscription** Information about subscribing to *The Journal of Immunology* is online at: <http://jimmunol.org/subscription>

**Permissions** Submit copyright permission requests at: <http://www.aai.org/About/Publications/JI/copyright.html>

**Email Alerts** Receive free email-alerts when new articles cite this article. Sign up at: <http://jimmunol.org/alerts>

*The Journal of Immunology* is published twice each month by  
The American Association of Immunologists, Inc.,  
1451 Rockville Pike, Suite 650, Rockville, MD 20852  
Copyright © 2006 by The American Association of  
Immunologists All rights reserved.  
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



## CUTTING EDGE

## Cutting Edge: Resistance to HIV-1 Infection among African Female Sex Workers Is Associated with Inhibitory KIR in the Absence of Their HLA Ligands<sup>1</sup>

Wim Jennes,<sup>2,3\*</sup> Sonja Verheyden,<sup>2†</sup> Christian Demanet,<sup>†</sup> Christiane A. Adjé-Touré,<sup>‡</sup> Bea Vuylsteke,<sup>\*‡</sup> John N. Nkengasong,<sup>‡§</sup> and Luc Kestens<sup>\*</sup>

*NK cells are regulated in part by killer Ig-like receptors (KIR) that interact with HLA molecules on potential target cells. KIR and HLA loci are highly polymorphic and certain KIR/HLA combinations were found to protect against HIV disease progression. We show in this study that KIR/HLA interactions also influence resistance to HIV transmission. HIV-exposed but seronegative female sex workers in Abidjan, Côte d'Ivoire, frequently possessed inhibitory KIR genes in the absence of their cognate HLA genes: KIR2DL2/KIR2DL3 heterozygosity in the absence of HLA-C1 and KIR3DL1 homozygosity in the absence of HLA-Bw4. HIV-seropositive female sex workers were characterized by corresponding inhibitory KIR/HLA pairings: KIR2DL3 homozygosity together with HLA-C1 and a trend toward KIR3DL1/HLA-Bw4 homozygosity. Absence of ligands for inhibitory KIR could lower the threshold for NK cell activation. In addition, exposed seronegatives more frequently possessed AB KIR genotypes, which contain more activating KIR. The data support an important role for NK cells and KIR/HLA interactions in antiviral immunity. The Journal of Immunology, 2006, 177: 6588–6592.*

The NK cells play an important role in the innate immune system by providing the first line of defense against viral infections and tumors (1). NK cell activity is partially controlled by distinct inhibitory and activating killer Ig-like receptors (KIR)<sup>4</sup> that recognize specific ligands on potential target cells (2). NK effector functions occur only when inhibitory signals are overcome by activating signals. KIR are encoded by 15 polymorphic genes located on chromosome 19 and contain two or three external Ig-like domains (2D, 3D)

with either long (2DL, 3DL) or short (2DS, 3DS) cytoplasmic tails, corresponding to their function as inhibitory or activating receptors, respectively (3). Several inhibitory KIR have well-defined HLA class I ligands. KIR2DL2 and KIR2DL3 bind to HLA-Cw group 1 (HLA-C1), which have asparagine at position 80, whereas KIR2DL1 binds the mutually exclusive HLA-Cw group 2 (HLA-C2) with a lysine at this position (4). KIR3DL1 binds HLA-B molecules with the serologically defined Bw4 epitope, specified by five variable amino acids spanning positions 77–83 (5). The alternative serotype, Bw6, is not known to bind any KIR. Despite high sequence similarity with inhibitory receptors, activating 2DS and 3DS KIR show either weak or undetectable binding to HLA class I (6–8).

The extreme levels of population diversity and rapid evolution of both *KIR* and *HLA* genes suggest that they are under pathogen-mediated selection (9). *KIR* and *HLA* loci map to separate chromosomes resulting in variation in the number and kind of KIR/HLA ligand pairs, potentially influencing disease outcome at the individual level. Indeed, specific *KIR/HLA* genotypes favoring NK cell activation were found to protect against disease progression after HIV-1 or hepatitis C virus infection (10, 11) and were associated with increased susceptibility to autoimmune disorders like psoriatic arthritis (12) and type I diabetes (13).

Rare individuals remain HIV-seronegative despite frequent unprotected exposure to the virus and several mechanisms of resistance have been proposed (HIV-exposed seronegative (ESN), reviewed in ref. 14). *HLA* polymorphism has been shown to play a role in protection against HIV infection (15, 16), but its mode of action remains incompletely resolved. In this study, we analyzed the genes encoding KIR and their HLA ligands in ESN female sex workers (FSWs), HIV-1-seropositive (SP) FSWs, and HIV-seronegative female blood donors (FBDs) from Abidjan, Côte d'Ivoire.

\*Department of Microbiology, Institute of Tropical Medicine, Antwerp, Belgium; †HLA Laboratory, Academisch Ziekenhuis-Vrije Universiteit Brussel, Brussels, Belgium; ‡Projet RETRO-CI, Abidjan, Côte d'Ivoire; §Division of HIV/AIDS Prevention, National Center for HIV, STD, and TB Prevention, Centers for Disease Control and Prevention, Atlanta, GA.

Received for publication April 25, 2006. Accepted for publication September 11, 2006.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup> This work was supported by the Belgian Fund for Scientific Research (FWO-Vlaanderen), Grant G.0660.06, and the Centers for Disease Control and Prevention, Division of

HIV/AIDS Prevention, Atlanta, GA. S.V. was supported by a grant from the Scientific Fund W. Gepts AZ-VUB.

<sup>2</sup> W.J. and S.V. contributed equally to this work.

<sup>3</sup> Address correspondence and reprint requests to Dr. Wim Jennes, Department of Microbiology, Laboratory of Immunology, Institute of Tropical Medicine, Nationalestraat 155, 2000 Antwerpen, Belgium. E-mail address: wjennes@itg.be

<sup>4</sup> Abbreviations used in this paper: KIR, killer Ig-like receptor; ESN, HIV-exposed seronegative; FBD, female blood donor; FSW, female sex worker; SP, HIV-1-seropositive; CI, confidence interval.

## Materials and Methods

### Study subjects

Twenty-one ESN and 20 SP FSWs were enrolled at a confidential FSW clinic in Abidjan, Côte d'Ivoire, between June 1998 and May 2000. The women were followed as part of a clinical trial testing the efficacy of a nonoxynol-9 microbicide gel (17). Twenty-five HIV-seronegative FBDs were enrolled at the blood transfusion center in Abidjan, Côte d'Ivoire. The study was approved by ethical committees of the Ministry of Health, Côte d'Ivoire, and the Institute of Tropical Medicine, Antwerp, Belgium, and by the Institutional Review Board of the Centers for Disease Control and Prevention, Atlanta, GA. All subjects gave written informed consent before enrollment.

### Laboratory methods

Whole blood was drawn in EDTA tubes (BD Biosciences). Plasma was separated from whole blood by centrifugation and tested for HIV by ELISA and Western blot, and confirmed by HIV RT-PCR. PBMC were separated from whole blood by gradient centrifugation and stored in liquid nitrogen.

### KIR and HLA class I genotyping

Genomic DNA was extracted from PBMC using a QIAamp DNA blood mini kit (Qiagen). KIR typing was performed with the PCR sequence-specific primer technique as previously reported (18). In the assessment of KIR genotypes, group A haplotypes were defined by the presence of *KIR2DL1*, *KIR2DL3*, *KIR3DL1*, and *KIR2DS4*. Group B haplotypes were defined by lack of *KIR3DL1* and *KIR2DS4* in the presence of inhibitory *KIR2DL2* and *KIR2DL5* and one or more of the activating *KIR2DS1*, *KIR2DS2*, *KIR2DS3*, *KIR2DS5*, and *KIR3DS1* (19). *HLA-B* and *-C* typing at the allele group level was performed by PCR sequence-specific oligonucleotides methodology (Tepnel Lifecodes).


### Statistical methods

Differences in continuous variables were analyzed with Mann-Whitney *U* tests. Fisher's exact tests were used to calculate statistical significances and exact 95% confidence intervals (CIs) of odds ratios of allele frequency differences. Occurrence of trend was analyzed with the exact non-parametric Cochran-Armitage test. Level of statistical significance for all analyses was set at  $p < 0.05$ . Statistical analyses were performed with SAS version 9.1 (The SAS Institute).

## Results and Discussion

Twenty-one ESN FSWs, 20 SP FSWs, and 25 FBDs were included in the study (Table I). First, we analyzed the frequencies of individual KIR genes and KIR genotypes. ESN FSWs showed significantly higher frequencies of inhibitory *KIR2DL2* and *KIR2DL5* than SP FSWs (Fig. 1A). ESN FSWs also showed higher frequencies of activating *KIR2DS2* and *KIR2DS3*, but the differences were not statistically significant (Fig. 1B). All study subjects displayed *KIR3DL1* and *KIR2DS4* genes, therefore only AA and AB genotypes were present. ESN FSWs more frequently displayed AB genotypes, whereas SP FSWs more frequently displayed AA genotypes (Fig. 1C). In general, FBDs showed frequencies of individual KIR genes and KIR genotypes in between those from ESN and SP FSWs (Fig. 1).

Next, we analyzed the presence of inhibitory KIR in association with their known HLA ligand genes (Tables II and III). First, we analyzed gene frequencies of the inhibitory receptors *KIR2DL2* and *KIR2DL3*, which segregate as alleles of the same gene locus, together with their HLA-C1 ligand (Table II).



**FIGURE 1.** Frequencies of individual KIR genes and KIR genotypes among ESN FSWs, SP FSWs, and FBDs. A, Inhibitory KIR genes; B, activating KIR genes; C, KIR genotypes. Gene and genotype frequencies were compared between ESN and FBDs and between ESN and SP FSWs with non-parametric Fisher's exact tests, *p* values  $< 0.1$  are shown. SP FSWs,  $n = 20$ , white bars; FBDs,  $n = 25$ , hatched bars; ESN FSWs,  $n = 21$ , black bars.

There were significantly more *KIR2DL2/KIR2DL3* heterozygotes among ESN FSWs, whereas SP FSWs showed a markedly higher proportion of *KIR2DL3/KIR2DL3* homozygotes. There was a trend toward a higher frequency of *HLA-C1/C2* heterozygotes among SP FSWs, potentially resulting in more inhibitory NK signals via KIR2DL receptors. More specifically, we found that the increased frequency of homozygous *KIR2DL3* among SP FSWs was significant only if both *HLA-C1* and *-C2* were present. ESN FSWs retained an increased frequency of heterozygous *KIR2DL2/KIR2DL3* only in the absence of their HLA-C1 ligand gene (i.e., *C2/C2* homozygous). The trends for the KIR/HLA comparisons between ESN FSWs and FBDs were similar to those between ESN and SP FSWs, although they did not always reach statistical significance probably because of the low sample size.

Next, we analyzed frequencies of inhibitory *KIR3DL1* and activating *KIR3DS1*, also alleles of the same gene locus, together with the HLA-Bw4 ligand *KIR3DL1* (Table III).

Table I. Characteristics of female sex workers and female blood donors included in the study<sup>a</sup>

	ESN ( $n = 21$ )	FBD ( $n = 25$ )	SP ( $n = 20$ )	ESN vs FBD <i>p</i>	ESN vs SP <i>p</i>
Age (years)	30 (26–35)	22 (21–27)	33 (25–38)	0.008	0.592
Duration of commercial sex work (mo)	43 (25–76)	NA	33 (16–91)	NA	0.554
CD4 <sup>+</sup> T cells (cells/ $\mu$ l)	1221 (987–1444)	NA	437 (296–747)	NA	$< 0.001$

<sup>a</sup> Data are median values (interquartile range). NA, Not available. *p* values calculated with Mann-Whitney *U* tests.

Table II. Frequencies of inhibitory *KIR2DL2* and *KIR2DL3* alleles and their *HLA-C1* ligand alleles gene among female sex workers and female blood donors<sup>a</sup>

	ESN (n = 21)	FBD (n = 25)	SP (n = 20)	ESN vs FBD			ESN vs SP		
				OR	95% CI	p	OR	95% CI	p
<i>KIR</i> alleles									
<i>2DL2/2DL2</i>	24	20	15	1.25	0.24–6.48	1.000	1.77	0.28–13.1	0.697
<i>2DL2/2DL3</i>	62	44	25	2.07	0.55–7.98	0.253	4.88	1.08–23.5	0.028*
<i>2DL3/2DL3</i>	14	36	60	0.30	0.05–1.50	0.176	0.11	0.02–0.60	0.004*
<i>HLA-C</i> alleles									
<i>C1/C1</i>	29	28	15	1.03	0.23–4.49	1.000	2.27	0.39–16.2	0.454
<i>C1/C2</i>	43	56	70	0.59	0.16–2.20	0.554	0.32	0.07–1.38	0.118
<i>C2/C2</i>	29	16	15	2.10	0.41–11.8	0.475	2.27	0.39–16.2	0.454
<i>KIR-HLA</i> combinations									
<i>2DL2/2DL3 + C1/C1</i>	19	12	10	1.73	0.25–13.2	0.686	2.12	0.26–25.8	0.663
<i>2DL2/2DL3 + C1/C2</i>	24	32	15	0.66	0.14–2.92	0.744	1.77	0.28–13.1	0.697
<i>2DL2/2DL3 + C2/C2</i>	19	0	0	∞	1.16–∞	0.037*	∞	0.92–∞	0.107
<i>2DL3/2DL3 + C1/C1</i>	0	8	5	0.00	0.00–4.11	0.493	0.00	0.00–18.1	0.488
<i>2DL3/2DL3 + C1/C2</i>	10	20	45	0.42	0.04–3.02	0.428	0.13	0.01–0.82	0.015*
<i>2DL3/2DL3 + C2/C2</i>	5	8	10	0.58	0.01–12.0	1.000	0.45	0.01–9.51	0.606

<sup>a</sup> Data are percentages. Statistical significance and exact 95% CI of odds ratios (OR) were calculated with Fisher's exact tests.

\*,  $p < 0.05$ .

*KIR3DL1* and *KIR3DS1* frequencies were similar among ESN and SP FSWs. SP FSWs tended to be more frequently homozygous for *HLA-Bw4*, on its own and in combination with *KIR3DL1*, potentially resulting in more NK inhibition via *KIR3DL1*. A significantly higher proportion of ESN FSWs who were homozygous for *KIR3DL1* lacked *HLA-Bw4* (i.e., *Bw6* homozygous), resulting in the absence of NK inhibition via *KIR3DL1*. Here, the trends for the *KIR/HLA* comparisons between ESN FSWs and FBDs did not confirm those between ESN and SP FSWs: similar proportions of ESN FSWs and FBDs were *HLA-Bw4* and *KIR3DL1/HLA-Bw6* homozygous. None of the subjects included in the study were homozygous for *KIR3DS1* (Table III). Slow progression toward AIDS was previously associated with *KIR3DS1* in the presence of *HLA-Bw4* alleles with isoleucine at position 80 (*Bw4-80Ile*) (10). In our study, frequencies of *Bw4-80Ile* were similar among ESN, SP FSWs, and FBDs. Compared with ESN FSWs, SP FSWs displayed a higher proportion of the alternative *HLA-Bw4* allele with a threonine at position 80 ( $p = 0.028$ ). No statistically

significant interactions were observed between *KIR3DS1* and *HLA-B* alleles, possibly as the result of the low frequency of *KIR3DS1* in this population (12% compared with 42% among healthy Caucasians; see Ref. 18). No correlations were found between the observed *KIR2DL/HLA-C* and *KIR3DL1/HLA-B* interactions (data not shown).

At the time of sample collection, HIV-1 seroprevalence was 32% among FSWs (20), and 14% among their clients (21). HIV-1 seroincidence among FSWs participating in the non-oxynol-9 trial was 4% (17). Using self-reported data, ESN FSWs in Abidjan were estimated to have on average 52 unprotected exposures to HIV-1 per year (22). None of the FSWs tested in Abidjan showed the *CCR5* 32-bp deletion (23), which is in agreement with other populations in Africa (24). Together, this suggests that ESN FSWs in Abidjan are at high risk for acquiring HIV-1 infection. ESN FSWs enrolled in this study reported a median duration of commercial sex work of 3.5 years, ranging from as short as 2 mo to as long as 20 years (Table I). Putative HIV-1 resistance factors may be selected for in ESN

Table III. Frequencies of *KIR3DL1* and *KIR3DS1* alleles and the *HLA-Bw4* ligand alleles among female sex workers and female blood donors<sup>a</sup>

	ESN (n = 21)	FBD (n = 25)	SP (n = 20)	ESN vs FBD			ESN vs SP		
				OR	95% CI	p	OR	95% CI	p
<i>KIR</i> alleles									
<i>3DL1/3DL1</i>	95	84	85	3.81	0.33–197	0.357	3.53	0.25–194	0.343
<i>3DL1/3DS1</i>	5	16	15	0.26	0.01–3.03	0.357	0.28	0.01–4.02	0.343
<i>3DS1/3DS1</i>	0	0	0	NA	NA	NA	NA	NA	NA
<i>HLA-B</i> alleles <sup>b</sup>									
<i>Bw4/Bw4</i>	10	13	38	0.70	0.05–6.90	1.000	0.18	0.02–1.28	0.055
<i>Bw4/Bw6</i>	62	52	56	1.49	0.38–5.88	0.557	1.26	0.28–5.75	0.749
<i>Bw6/Bw6</i>	29	35	6	0.75	0.17–3.21	0.752	6.00	0.59–294	0.113
<i>KIR-HLA</i> combinations <sup>b</sup>									
<i>3DL1/3DL1 + Bw4/Bw4</i>	10	4	31	2.32	0.11–143	0.599	0.23	0.02–1.78	0.202
<i>3DL1/3DL1 + Bw4/Bw6</i>	57	48	50	1.45	0.38–5.63	0.563	1.33	0.30–5.96	0.746
<i>3DL1/3DL1 + Bw6/Bw6</i>	29	35	0	0.75	0.17–3.21	0.752	∞	1.37–∞	0.027*
<i>3DL1/3DS1 + Bw4/Bw4</i>	0	9	6	0.00	0.00–3.77	0.489	0.00	0.00–14.5	0.432
<i>3DL1/3DS1 + Bw4/Bw6</i>	5	4	6	1.10	0.01–90.2	1.000	0.75	0.01–62.8	1.000
<i>3DL1/3DS1 + Bw6/Bw6</i>	0	0	6	NA	NA	NA	0.00	0.00–14.5	0.432

<sup>a</sup> Data are percentages. NA, Not available. Statistical significance and exact 95% CI of odds ratios (OR) were calculated with Fisher's exact tests.

<sup>b</sup> Data available for 16 SP FSWs and 23 FBDs due to insufficient amounts of DNA.


\*,  $p < 0.05$ .



subjects with a longer history of high-risk behavior. To test this, we correlated the observed *KIR/HLA* associations with the duration of commercial sex work of the ESN FSWs. For the *KIR* and *KIR/HLA* combinations that reached statistical significance in Tables II and III, we observed a statistically significant trend over SP and ESN FSWs who had done more and <3 years of commercial sex work (Fig. 2).

In this study, the ESN status of FSWs was associated with the occurrence of certain inhibitory *KIR* genes in the absence of their cognate *HLA* genes: *KIR2DL2/KIR2DL3* heterozygosity in the absence of *HLA-C1*, and independent from this, *KIR3DL1* homozygosity in the absence of *HLA-Bw4*. Together, 38% of ESN FSWs showed at least one such *KIR/HLA* mismatch compared with 0% of SP FSWs ( $p = 0.006$ ). Absence of HLA ligands for inhibitory KIR may lower the threshold for NK cell activation via activating KIR, resulting in NK cytotoxic activity and early elimination of HIV-infected cells. In agreement with this, a higher frequency of ESN FSWs possessed *AB KIR* genotypes, which contain a higher number of activating *KIR*. In that respect, our data corroborate a recent study showing increased NK cell-mediated cytotoxicity and cytokine and  $\beta$ -chemokine secretion among ESN intravascular drug users (25). In contrast with this, SP FSWs were characterized by corresponding inhibitory *KIR/HLA* ligand pairs that can directly inhibit NK cell effector functions: *KIR2DL3* homozygosity in the presence of heterozygous *HLA-C1/C2*, and independent from this, a trend toward *KIR3DL1/HLA-Bw4* homozygosity. Together, 61% of SP FSWs showed at least one such inhibitory *KIR/HLA* pair compared with 19% of ESN FSWs ( $p = 0.009$ ). In addition, SP FSWs more frequently showed *AA KIR* genotypes that contain lower numbers of activating *KIR*.

Although SP FSWs showed a wide range of CD4 counts (median, 437 cells/ $\mu$ l; range, 144–1712 cells/ $\mu$ l; Table I), it is unclear to what extent rapid and slow progressor patients were



**FIGURE 2.** Test for trend of *KIR* and *HLA* frequencies with duration of commercial sex work of ESN FSWs. *A*, *KIR* alleles; *B*, *KIR/HLA* allele combinations. *KIR* alleles and *KIR/HLA* allele combinations that reached statistical significance in Tables II and III were tested. Tests for trend were performed with exact non-parametric Cochran-Armitage tests. SP FSWs,  $n = 20$  (except for *KIR3DL1/3DL1* + *HLA Bw6/Bw6*,  $n = 16$ ), white bars; ESN <3 years, HIV-exposed seronegative FSWs with <3 years of commercial sex work,  $n = 9$ , hatched bars; ESN  $\geq 3$  years, HIV-exposed seronegative FSWs with >3 years of commercial sex work,  $n = 12$ , black bars.

equally represented; this cannot be known in a cross-sectional study. Thus, it cannot be ruled out that *KIR/HLA* combinations related to differential disease progression in the SP group have influenced our results. However, although less pronounced, the *KIR* and *HLA* frequency differences among ESN and SP FSWs were generally confirmed by comparisons between ESN FSWs and FBDs. This supports the conclusion that the observed associations are related to HIV resistance rather than differential HIV disease progression.

Despite large gene frequency differences, the small sample size resulted in *KIR/HLA* interactions with relatively weak statistical significance. Because of the exploratory nature of the study, the testing of distinct hypotheses rather than a comprehensive screening for genes and associations, and restriction of *HLA-B* and *-C* allele analysis to four distinct *KIR* binding groups, correction for multiple testing was not applied. Exact testing for small sample sizes was performed for all analyses. Nevertheless, confirmation of our findings in larger populations of healthy vs HIV-infected subjects, and in functional studies, is needed.

Our data are the first to show that *KIR/HLA* interactions may influence susceptibility to virus transmission. Moreover, the *KIR/HLA* gene combinations favoring NK activation over inhibition by lack of HLA ligands for inhibitory *KIR* may be more straightforward than those previously described for protection against viral disease progression. Indeed, delayed onset of AIDS was associated with the activating *KIR3DS1* receptor in combination with its only presumptive *HLA-B Bw4-80Ile* ligand (10). Resolution of hepatitis C virus infection was found to be associated with corresponding *KIR2DL3/HLA-C1* gene combinations, only putatively resulting in the weakest inhibitory signals (11).

The non-classical *HLA* class I molecules *HLA-E* and *-G* are ligands for the inhibitory NK cell receptors *CD94/NKG2A* and *KIR2DL4*, respectively. A recent study found genetic variants of *HLA-E* and *HLA-G* with a potentially lower affinity for their inhibitory NK cell receptors to be associated with a decreased risk of HIV transmission (26). These findings further support a role for NK cells in protection against virus transmission and suggest that parallel inhibitory NK receptor/*HLA* ligand mechanisms may be at play.

## Acknowledgments

We thank the community of female sex workers in Abidjan for their cooperation, and Joris Menten and Odette Tossou for help with data analysis.

## Disclosures

The authors have no financial conflict of interest.

## References

- Hamerman, J. A., K. Ogasawara, and L. L. Lanier. 2005. NK cells in innate immunity. *Curr. Opin. Immunol.* 17: 29–35.
- Vilches, C., and P. Parham. 2002. KIR: diverse, rapidly evolving receptors of innate and adaptive immunity. *Annu. Rev. Immunol.* 20: 217–251.
- Martin, A. M., E. M. Freitas, C. S. Witt, and F. T. Christiansen. 2000. The genomic organization and evolution of the natural killer immunoglobulin-like receptor (*KIR*) gene cluster. *Immunogenetics* 51: 268–280.
- Moretta, A., M. Vitale, C. Bottino, A. M. Orengo, L. Morelli, R. Augugliaro, M. Barbaresi, E. Ciccone, and L. Moretta. 1993. P58 molecules as putative receptors for major histocompatibility complex (MHC) class I molecules in human natural killer (NK) cells. Anti-p58 antibodies reconstitute lysis of MHC class I-protected cells in NK clones displaying different specificities. *J. Exp. Med.* 178: 597–604.
- Gumperz, J. E., V. Litwin, J. H. Phillips, L. L. Lanier, and P. Parham. 1995. The *Bw4* public epitope of *HLA-B* molecules confers reactivity with natural killer cell clones that express *NKB1*, a putative *HLA* receptor. *J. Exp. Med.* 181: 1133–1144.

6. Biassoni, R., A. Pessino, A. Malaspina, C. Cantoni, C. Bottino, S. Sivori, L. Moretta, and A. Moretta. 1997. Role of amino acid position 70 in the binding affinity of p50.1 and p58.1 receptors for HLA-Cw4 molecules. *Eur. J. Immunol.* 27: 3095–3099.
7. Vales-Gomez, M., H. T. Reyburn, R. A. Erskine, and J. Strominger. 1998. Differential binding to HLA-C of p50-activating and p58-inhibitory natural killer cell receptors. *Proc. Natl. Acad. Sci. USA* 95: 14326–14331.
8. Winter, C. C., J. E. Gumperz, P. Parham, E. O. Long, and N. Wagtmann. 1998. Direct binding and functional transfer of NK cell inhibitory receptors reveal novel patterns of HLA-C allotype recognition. *J. Immunol.* 161: 571–577.
9. Parham, P. 2005. MHC class I molecules and KIRs in human history, health and survival. *Nat. Rev. Immunol.* 5: 201–214.
10. Martin, M. P., X. Gao, J. H. Lee, G. W. Nelson, R. Detels, J. J. Goedert, S. Buchbinder, K. Hoots, D. Vlahov, J. Trowsdale, et al. 2002. Epistatic interaction between KIR3DS1 and HLA-B delays the progression to AIDS. *Nat. Genet.* 31: 429–434.
11. Khakoo, S. I., C. L. Thio, M. P. Martin, C. R. Brooks, X. Gao, J. Astemborski, J. Cheng, J. J. Goedert, D. Vlahov, M. Hilgartner, et al. 2004. HLA and NK cell inhibitory receptor genes in resolving hepatitis C virus infection. *Science* 305: 872–874.
12. Nelson, G. W., M. P. Martin, D. Gladman, J. Wade, J. Trowsdale, and M. Carrington. 2004. Cutting edge: heterozygote advantage in autoimmune disease: hierarchy of protection/susceptibility conferred by HLA and killer Ig-like receptor combinations in psoriatic arthritis. *J. Immunol.* 173: 4273–4276.
13. van der Slik, A. R., B. P. Koeleman, W. Verduijn, G. J. Bruining, B. O. Roep, and M. J. Giphart. 2003. KIR in type 1 diabetes: disparate distribution of activating and inhibitory natural killer cell receptors in patients versus HLA-matched control subjects. *Diabetes* 52: 2639–2642.
14. Kulkarni, P. S., S. T. Butera, and A. C. Duerr. 2003. Resistance to HIV-1 infection: lessons learned from studies of highly exposed persistently seronegative (HEPS) individuals. *AIDS Rev.* 5: 87–103.
15. MacDonald, K. S., K. R. Fowke, J. Kimani, V. A. Dunand, N. J. Nagelkerke, T. B. Ball, J. Oyugi, E. Njagi, L. K. Gaur, R. C. Brunham, et al. 2000. Influence of HLA supertypes on susceptibility and resistance to human immunodeficiency virus type 1 infection. *J. Infect. Dis.* 181: 1581–1589.
16. Dorak, M. T., J. Tang, A. Penman-Aguilar, A. O. Westfall, I. Zulu, E. S. Lobashevsky, N. G. Kancheya, M. M. Schaen, S. A. Allen, and R. A. Kaslow. 2004. Transmission of HIV-1 and HLA-B allele-sharing within serodiscordant heterosexual Zambian couples. *Lancet* 363: 2137–2139.
17. Van Damme, L., G. Ramjee, M. Alary, B. Vuylsteke, V. Chandeying, H. Rees, P. Sirivongrangsorn, L. Mukenge-Tshibaka, V. Ertiegne-Traore, C. Uahowitchai, et al. 2002. Effectiveness of COL-1492, a nonoxynol-9 vaginal gel, on HIV-1 transmission in female sex workers: a randomised controlled trial. *Lancet* 360: 971–977.
18. Verheyden, S., M. Bernier, and C. Demanet. 2004. Identification of natural killer cell receptor phenotypes associated with leukemia. *Leukemia* 18: 2002–2007.
19. Marsh, S. G., P. Parham, B. Dupont, D. E. Geraghty, J. Trowsdale, D. Middleton, C. Vilches, M. Carrington, C. Witt, L. A. Guethlein, et al. 2003. Killer-cell immunoglobulin-like receptor (KIR) nomenclature report, 2002. *Hum. Immunol.* 64: 648–654.
20. Ghys, P. D., M. O. Diallo, V. Ertiegne-Traore, K. Kale, O. Tawil, M. Carael, M. Traore, G. Mah-Bi, K. M. De Cock, S. Z. Wiktor, et al. 2002. Increase in condom use and decline in HIV and sexually transmitted diseases among female sex workers in Abidjan, Cote d'Ivoire, 1991–1998. *AIDS* 16: 251–258.
21. Vuylsteke, B. L., P. D. Ghys, M. Traore, Y. Konan, G. Mah-Bi, C. Maurice, D. Soroh, J. N. Diarra, T. H. Roels, and M. Laga. 2003. HIV prevalence and risk behavior among clients of female sex workers in Abidjan, Cote d'Ivoire. *AIDS* 17: 1691–1694.
22. Jennes, W., B. Vuylsteke, M. Y. Borget, V. Traore-Ertiegne, C. Maurice, M. Nolan, J. N. Nkengasong, and L. Kestens. 2004. HIV-specific T helper responses and frequency of exposure among HIV-exposed seronegative female sex workers in Abidjan, Cote d'Ivoire. *J. Infect. Dis.* 189: 602–610.
23. Jennes, W., S. Sawadogo, S. Koblavi-Deme, B. Vuylsteke, C. Maurice, T. H. Roels, T. Chorba, J. N. Nkengasong, and L. Kestens. 2003. Cellular human immunodeficiency virus (HIV)-protective factors: a comparison of HIV-exposed seronegative female sex workers and female blood donors in Abidjan, Cote d'Ivoire. *J. Infect. Dis.* 187: 206–214.
24. Martinson, J. J., N. H. Chapman, D. C. Rees, Y. T. Liu, and J. B. Clegg. 1997. Global distribution of the CCR5 gene 32-basepair deletion. *Nat. Genet.* 16: 100–103.
25. Scott-Algara, D., L. X. Truong, P. Versmisse, A. David, T. T. Luong, N. V. Nguyen, I. Theodorou, F. Barre-Sinoussi, and G. Pancino. 2003. Cutting edge: increased NK cell activity in HIV-1-exposed but uninfected Vietnamese intravenous drug users. *J. Immunol.* 171: 5663–5667.
26. Lajoie, J., J. Hargrove, L. S. Zijenah, J. H. Humphrey, B. J. Ward, and M. Roger. 2006. Genetic variants in nonclassical major histocompatibility complex class I human leukocyte antigen (HLA)-E and HLA-G molecules are associated with susceptibility to heterosexual acquisition of HIV-1. *J. Infect. Dis.* 193: 298–301.