


NEW *InVivoSIM*[™]
Biosimilar Antibodies

For Research Use Only

DISCOVER BioCell



 *The Journal of
Immunology*

This information is current as
of June 16, 2021.

Comment on "Paroxetine Prevents Loss of Nigrostriatal Dopaminergic Neurons by Inhibiting Brain Inflammation and Oxidative Stress in an Experimental Model of Parkinson's Disease"

Jonas Hannestad

J Immunol 2010; 185:4966; ;

doi: 10.4049/jimmunol.1090095

<http://www.jimmunol.org/content/185/9/4966.1>

References This article **cites 1 articles**, 1 of which you can access for free at:
<http://www.jimmunol.org/content/185/9/4966.1.full#ref-list-1>

Why *The JI*? [Submit online.](#)

- **Rapid Reviews! 30 days*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

**average*

Subscription Information about subscribing to *The Journal of Immunology* is online at:
<http://jimmunol.org/subscription>

Permissions Submit copyright permission requests at:
<http://www.aai.org/About/Publications/JI/copyright.html>

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://jimmunol.org/alerts>

The Journal of Immunology is published twice each month by
The American Association of Immunologists, Inc.,
1451 Rockville Pike, Suite 650, Rockville, MD 20852
Copyright © 2010 by The American Association of
Immunologists, Inc. All rights reserved.
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



Comment on “Cutting Edge: FcR-Like 6 Is an MHC Class II Receptor”

The article published by Schreeder et al. (1) in the July 1, 2010 issue of *The Journal of Immunology* identifies MHC class II molecules as ligands for FcRL-6, an Ig-like molecule belonging to the FcR-like (FcRL) family. The function of FcRL molecules remains elusive because they share many features with classical FcR molecules, but they do not bind Igs [during the revision process, Wilson et al. (2) reported an intracellular association of FcRL-A and Igs]. Identification of MHC class II as a ligand for FcRL-6 renders this molecule strikingly similar to another member of the Ig superfamily, CD223. CD223 is not part of the FcRL family because of its distinct chromosomal localization and weak sequence homology, but it is a receptor for MHC class II (3). FcRL-6 and CD223 show a restricted cellular expression on NK, $\gamma\delta$, and Ag-experienced $\alpha\beta$ T cells, mostly CD8⁺ (4–6). In addition, a situation of chronic T cell activation, such as HIV infection (5, 7) and tumor development (6, 8), increases expression of these molecules. CD223 is an immunosuppressive molecule (9, 10), and it is likely that FcRL-6 also negatively regulates cell activation due to its ITIM present in the intracytoplasmic tail. Hence, several inhibitory receptors specific for MHC class II molecules may be present on the surface of NK and activated T cells. The function of these receptors is not easy to understand on NK cells, but CD223 and FcRL-6 may contribute to the exhaustion phenomenon frequently observed in chronically stimulated T cells.

Bertrand Huard

Hematology Unit, University Hospitals; and Department of Pathology-Immunology, Faculty of Medicine, University Medical Center, Geneva, Switzerland

References

1. Schreeder, D. M., J. P. Cannon, J. Wu, R. Li, M. A. Shakhmatov, and R. S. Davis. 2010. Cutting edge: FcR-like 6 is an MHC class II receptor. *J. Immunol.* 185: 23–27.
2. Wilson, T. J., S. Gilfillan, and M. Colonna. 2010. Fc receptor-like A associates with intracellular IgG and IgM but is dispensable for antigen-specific immune responses. *J. Immunol.* 185: 2960–2967.
3. Baixeras, E., B. Huard, C. Miossec, S. Jitsukawa, M. Martin, T. Hercend, C. Auffray, F. Triebel, and D. Piatier-Tonneau. 1992. Characterization of the lymphocyte activation gene 3-encoded protein. A new ligand for human leukocyte antigen class II antigens. *J. Exp. Med.* 176: 327–337.
4. Huard, B., P. Gaulard, F. Faure, T. Hercend, and F. Triebel. 1994. Cellular expression and tissue distribution of the human LAG-3-encoded protein, an MHC class II ligand. *Immunogenetics* 39: 213–217.
5. Wilson, T. J., R. M. Presti, I. Tassi, E. T. Overton, M. Cella, and M. Colonna. 2007. FcRL6, a new ITIM-bearing receptor on cytolytic cells, is broadly expressed by lymphocytes following HIV-1 infection. *Blood* 109: 3786–3793.

6. Schreeder, D. M., J. Pan, F. J. Li, E. Vivier, and R. S. Davis. 2008. FCRL6 distinguishes mature cytotoxic lymphocytes and is upregulated in patients with B-cell chronic lymphocytic leukemia. *Eur. J. Immunol.* 38: 3159–3166.
7. Li, Q., A. J. Smith, T. W. Schacker, J. V. Carlis, L. Duan, C. S. Reilly, and A. T. Haase. 2009. Microarray analysis of lymphatic tissue reveals stage-specific, gene expression signatures in HIV-1 infection. *J. Immunol.* 183: 1975–1982.
8. Demeure, C. E., J. Wolfers, N. Martin-Garcia, P. Gaulard, and F. Triebel. 2001. T Lymphocytes infiltrating various tumour types express the MHC class II ligand lymphocyte activation gene-3 (LAG-3): role of LAG-3/MHC class II interactions in cell-cell contacts. *Eur. J. Cancer* 37: 1709–1718.
9. Huard, B., P. Prigent, F. Pagès, D. Bruniquel, and F. Triebel. 1996. T cell major histocompatibility complex class II molecules down-regulate CD4+ T cell clone responses following LAG-3 binding. *Eur. J. Immunol.* 26: 1180–1186.
10. Workman, C. J., and D. A. Vignali. 2005. Negative regulation of T cell homeostasis by lymphocyte activation gene-3 (CD223). *J. Immunol.* 174: 688–695.

www.jimmunol.org/cgi/doi/10.4049/jimmunol.1090093

Response to Comment on “Cutting Edge: FcR-Like 6 Is an MHC Class II Receptor”

We thank Dr. Huard for his letter and appreciate the opportunity to comment on the interesting parallels between FcR-like 6 (FCRL6) and the CD223 receptor, lymphocyte activation gene 3. Indeed, both proteins bind MHC class II molecules (1, 2) and are expressed by cytotoxic NK and effector T cell subsets (3–5), but these receptors also possess several distinct features. Whereas FCRL6 is related to the classical FcRs for IgG and IgE and the more recently identified FCRL1–5 family that all derive from human chromosome 1q (6), the gene encoding CD223 is a CD4 paralog positioned in tandem on chromosome 12 (7). That FCRL6 and CD223, at least in humans, have evolved independently as T and NK cell surface receptors for MHC class II indicates the importance of these interactions in immunobiology. Interestingly, the *Fcrl6* ortholog in mice, which occupies a syntenic genomic location, has differences in its protein structure and expression, suggesting that its function may differ from its human counterpart (3, 8 and W.J. Won, R.P. Stephan, and R.S. Davis, unpublished observations). Another distinction is their modulation in response to cytokine stimulation. Whereas CD223 is induced following exposure to activating signals, such as IL-12 and IL-2 (9), FCRL6 is downregulated under similar conditions (3). This indicates that these molecules are regulated differently and implies that they may vary temporally in their interactions with MHC class II. Finally, although FCRL6 has a consensus ITIM and can recruit Src homology region 2 domain-containing phosphatase-2 (3), whether it functions as a bona fide inhibitory molecule and/or plays a role in T cell exhaustion similar to CD223 (10) are subjects of ongoing investigation.

Daniel M. Schreeder and Randall S. Davis

Department of Medicine, Department of Microbiology, and Department of Biochemistry and Molecular Genetics, University of Alabama at Birmingham, Birmingham, AL 35294

References

- Schreeder, D. M., J. P. Cannon, J. Wu, R. Li, M. A. Shakhmatov, and R. S. Davis. 2010. Cutting edge: FcR-like 6 is an MHC class II receptor. *J. Immunol.* 185: 23–27.
- Baixeras, E., B. Huard, C. Miossec, S. Jitsukawa, M. Martin, T. Hercend, C. Auffray, F. Triebel, and D. Piatier-Tonneau. 1992. Characterization of the lymphocyte activation gene 3-encoded protein. A new ligand for human leukocyte antigen class II antigens. *J. Exp. Med.* 176: 327–337.
- Wilson, T. J., R. M. Presti, I. Tassi, E. T. Overton, M. Cella, and M. Colonna. 2007. FcRL6, a new ITIM-bearing receptor on cytolytic cells, is broadly expressed by lymphocytes following HIV-1 infection. *Blood* 109: 3786–3793.
- Schreeder, D. M., J. Pan, F. J. Li, E. Vivier, and R. S. Davis. 2008. FCRL6 distinguishes mature cytotoxic lymphocytes and is upregulated in patients with B-cell chronic lymphocytic leukemia. *Eur. J. Immunol.* 38: 3159–3166.
- Huard, B., P. Gaulard, F. Faure, T. Hercend, and F. Triebel. 1994. Cellular expression and tissue distribution of the human LAG-3-encoded protein, an MHC class II ligand. *Immunogenetics* 39: 213–217.
- Davis, R. S. 2007. Fc receptor-like molecules. *Annu. Rev. Immunol.* 25: 525–560.
- Triebel, F., S. Jitsukawa, E. Baixeras, S. Roman-Roman, C. Genevee, E. Viegas-Pequignot, and T. Hercend. 1990. LAG-3, a novel lymphocyte activation gene closely related to CD4. *J. Exp. Med.* 171: 1393–1405.
- Davis, R. S., G. Dennis, Jr., M. R. Odom, A. W. Gibson, R. P. Kimberly, P. D. Burrows, and M. D. Cooper. 2002. Fc receptor homologs: newest members of a remarkably diverse Fc receptor gene family. *Immunol. Rev.* 190: 123–136.
- Bruniquel, D., N. Borie, S. Hannier, and F. Triebel. 1998. Regulation of expression of the human lymphocyte activation gene-3 (LAG-3) molecule, a ligand for MHC class II. *Immunogenetics* 48: 116–124.
- Blackburn, S. D., H. Shin, W. N. Haining, T. Zou, C. J. Workman, A. Polley, M. R. Betts, G. J. Freeman, D. A. Vignali, and E. J. Wherry. 2009. Coregulation of CD8+ T cell exhaustion by multiple inhibitory receptors during chronic viral infection. *Nat. Immunol.* 10: 29–37.

www.jimmunol.org/cgi/doi/10.4049/jimmunol.1090094

Comment on “Paroxetine Prevents Loss of Nigrostriatal Dopaminergic Neurons by Inhibiting Brain Inflammation and Oxidative Stress in an Experimental Model of Parkinson’s Disease”

I read with interest the article by Chung et al. (1) describing how pretreatment with the serotonin reuptake inhibitor paroxetine (10 mg/d for 6 d) prevented 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine–induced neuroinflammation (increased production of reactive oxygen species and proinflammatory cytokines by glial cells) and neuronal cell death in the substantia nigra in mice. The mechanism by which paroxetine inhibits inflammation is unknown. We have data from humans ($n = 9$) receiving i.v. LPS (0.8 ng/kg) that pretreatment with the serotonin reuptake inhibitor citalopram (20 mg/d for 5 d) did not reduce the innate immune response to LPS compared with placebo pretreatment. For instance, LPS administration caused IL-6 serum levels to increase from 1.3 ± 0.4 to 197.8 ± 47.9 pg/ml in subjects who received placebo pretreatment and from 0.9 ± 0.2 to 215.9 ± 54.2 in subjects who received citalopram pretreatment. Similarly, TNF levels increased from 2.0 ± 0.6 to 45.2 ± 10.8 in placebo-pretreated subjects and from 1.4 ± 0.3 to 38.3 ± 6.2 in citalopram-pretreated individuals. All p values were >0.8 for pairwise t tests. Our data indicate that the selective serotonin reuptake inhibitor citalopram does not have an anti-inflammatory effect in peripheral innate immune cells. This

may suggest that the anti-inflammatory effects of paroxetine observed by Chung et al. (1) is either specific to neuroinflammation (i.e., paroxetine has a differential effect on microglia compared with peripheral innate immune cells) or that the anti-inflammatory effect of paroxetine is mediated by increases in synaptic serotonin in the brain.

Jonas Hannestad

Department of Psychiatry, Yale University School of Medicine, New Haven, CT 06511

Reference

- Chung, Y. C., S. R. Kim, and B. K. Jin. 2010. Paroxetine prevents loss of nigrostriatal dopaminergic neurons by inhibiting brain inflammation and oxidative stress in an experimental model of Parkinson’s disease. *J. Immunol.* 185: 1230–1237.

www.jimmunol.org/cgi/doi/10.4049/jimmunol.1090095

Response to Comment on “Paroxetine Prevents Loss of Nigrostriatal Dopaminergic Neurons by Inhibiting Brain Inflammation and Oxidative Stress in an Experimental Model of Parkinson’s Disease”

We have evidence from 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-treated mice ($n = 3$) that citalopram (5 mg/kg/d for 6 d) did not inhibit microglial activation as evidenced by Mac-1 immunocytochemical staining. However, another serotonin uptake inhibitor (fluoxetine; 5 mg/kg/d for 6 d) inhibited MPTP-activated glial activation as assessed by Mac-1 and GFAP immunostaining. Fluoxetine was also found to prevent MPTP-induced neuroinflammation (increases in reactive oxygen species and proinflammatory cytokines, such as TNF- α , IL-1 β , and inducible NO synthase). In the LPS-injected rat substantia nigra, fluoxetine (10 mg/kg/twice a d for 6 d) dramatically reduced neuroinflammation (reactive oxygen species and proinflammatory cytokines). This is in line with recent findings that fluoxetine inhibits the expression of proinflammatory cytokines, such as TNF- α , and IL-1 β in the rat cerebral ischemia model of middle cerebral artery occlusion (1) and kainic acid-treated mouse hippocampus (2). Contrary to citalopram, Taler and his colleagues (3, 4) reported that paroxetine and sertraline, another serotonin uptake inhibitor, reduced secretion of TNF- α in human T lymphocytes and rat splenocytes. It is therefore likely that anti-inflammatory actions of paroxetine may not be specific to neuroinflammation, and not all of serotonin uptake inhibitors possess anti-inflammatory properties.

Striatal serotonin levels were measured in MPTP-treated mice to determine the relationship between anti-inflammatory effects of the fluoxetine and serotonin system. Results (MPTP:

14.6 ± 2.5 $\mu\text{g}/\text{mg}$ protein, $n = 6$; MPTP + fluoxetine: 15.3 ± 1.3 $\mu\text{g}/\text{mg}$ protein, $n = 6$) show that, whereas fluoxetine prevented MPTP-induced neuroinflammation, it failed to increase serotonin levels in the striatum. These results carefully suggest that the anti-inflammatory effect of paroxetine is not mediated by increases in synaptic serotonin, although serotonin levels are not provided.

Byung K. Jin

Department of Biochemistry and Molecular Biology, Neurodegeneration Control Research Center and Age-Related and Brain Diseases Research Center, School of Medicine, Kyung Hee University, Seoul, South Korea

References

1. Lim, C. M., S. W. Kim, J. Y. Park, C. Kim, S. H. Yoon, and J. K. Lee. 2009. Fluoxetine affords robust neuroprotection in the postischemic brain via its anti-inflammatory effect. *J. Neurosci. Res.* 87: 1037–1045.
2. Jin, Y., C. M. Lim, S. W. Kim, J. Y. Park, J. S. Seo, P. L. Han, S. H. Yoon, and J. K. Lee. 2009. Fluoxetine attenuates kainic acid-induced neuronal cell death in the mouse hippocampus. *Brain Res.* 1281: 108–116.
3. Taler, M., I. Gil-Ad, L. Lomnitski, I. Korob, E. Baharav, M. Bar, A. Zolokov, and A. Weizman. 2007. Immunomodulatory effect of selective serotonin reuptake inhibitors (SSRIs) on human T lymphocyte function and gene expression. *Eur. Neuropsychopharmacol.* 17: 774–780.
4. Taler, M., M. Bar, I. Korob, L. Lomnitski, E. Baharav, N. Grunbaum-Novak, A. Weizman, and I. Gil-Ad. 2008. Evidence for an inhibitory immunomodulatory effect of selected antidepressants on rat splenocytes: possible relevance to depression and hyperactive-immune disorders. *Int. Immunopharmacol.* 8: 526–533.

www.jimmunol.org/cgi/doi/10.4049/jimmunol.1090096