

## Spectral Flow Cytometry Webinar Series

Watch our webinar series and learn how the ID7000™ system builds on Sony's experience with spectral analysis and simplifies many operations to advance the field of flow cytometry.



Watch Now

SONY



## The Liver Kinase B1 Is a Central Regulator of T Cell Development, Activation, and Metabolism

This information is current as of March 6, 2022.

Nancie J. MacIver, Julianna Blagih, Donte C. Saucillo, Luciana Tonelli, Takla Griss, Jeffrey C. Rathmell and Russell G. Jones

*J Immunol* 2011; 187:4187-4198; Prepublished online 19 September 2011;  
doi: 10.4049/jimmunol.1100367  
<http://www.jimmunol.org/content/187/8/4187>

**Supplementary Material** <http://www.jimmunol.org/content/suppl/2011/09/16/jimmunol.1100367.DC1>

**References** This article **cites 63 articles**, 21 of which you can access for free at:  
<http://www.jimmunol.org/content/187/8/4187.full#ref-list-1>

**Why *The JI*? Submit online.**

- **Rapid Reviews! 30 days\*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

*\*average*

**Subscription** Information about subscribing to *The Journal of Immunology* is online at:  
<http://jimmunol.org/subscription>

**Permissions** Submit copyright permission requests at:  
<http://www.aai.org/About/Publications/JI/copyright.html>

**Email Alerts** Receive free email-alerts when new articles cite this article. Sign up at:  
<http://jimmunol.org/alerts>



# The Liver Kinase B1 Is a Central Regulator of T Cell Development, Activation, and Metabolism

Nancie J. MacIver,<sup>\*,1</sup> Julianna Blagih,<sup>†,‡,1</sup> Donte C. Saucillo,<sup>\*</sup> Luciana Tonelli,<sup>†,‡</sup> Takla Griss,<sup>†,‡</sup> Jeffrey C. Rathmell,<sup>§,¶,||</sup> and Russell G. Jones<sup>†,‡</sup>

T cell activation leads to engagement of cellular metabolic pathways necessary to support cell proliferation and function. However, our understanding of the signal transduction pathways that regulate metabolism and their impact on T cell function remains limited. The liver kinase B1 (LKB1) is a serine/threonine kinase that links cellular metabolism with cell growth and proliferation. In this study, we demonstrate that LKB1 is a critical regulator of T cell development, viability, activation, and metabolism. T cell-specific ablation of the gene that encodes LKB1 resulted in blocked thymocyte development and a reduction in peripheral T cells. LKB1-deficient T cells exhibited defects in cell proliferation and viability and altered glycolytic and lipid metabolism. Interestingly, loss of LKB1 promoted increased T cell activation and inflammatory cytokine production by both CD4<sup>+</sup> and CD8<sup>+</sup> T cells. Activation of the AMP-activated protein kinase (AMPK) was decreased in LKB1-deficient T cells. AMPK was found to mediate a subset of LKB1 functions in T lymphocytes, as mice lacking the  $\alpha 1$  subunit of AMPK displayed similar defects in T cell activation, metabolism, and inflammatory cytokine production, but normal T cell development and peripheral T cell homeostasis. LKB1- and AMPK $\alpha 1$ -deficient T cells each displayed elevated mammalian target of rapamycin complex 1 signaling and IFN- $\gamma$  production that could be reversed by rapamycin treatment. Our data highlight a central role for LKB1 in T cell activation, viability, and metabolism and suggest that LKB1-AMPK signaling negatively regulates T cell effector function through regulation of mammalian target of rapamycin activity. *The Journal of Immunology*, 2011, 187: 4187–4198.

**T** cells play a critical role in the generation of immunity and inflammatory processes. Upon TCR stimulation, T cells initiate a coordinated program of rapid proliferation and differentiation, leading to the development of activated T cells with specific effector functions tailored toward pathogen clearance or control. As part of their program of activation, T cells dramatically alter their metabolic activity to meet the increased metabolic

demands of cell growth, proliferation, and effector function (1). Activated T cells generate ATP in large part by increasing their rate of glucose uptake and glycolysis (2, 3). Glucose availability is essential for the survival, proliferation, and effector function of activated T cells (4, 5). In contrast, quiescent T cells and regulatory T cells rely primarily on mitochondrial oxidative metabolism fueled by lipids and other metabolic substrates to maintain bioenergetic homeostasis and suppressor function (6–8). In addition, a switch from aerobic glycolysis to mitochondrial oxidative pathways is required for the establishment of CD8 T cell memory (9, 10). Thus, metabolic regulation is a crucial factor in T cell function, differentiation, and fate (11, 12).

The signal transduction pathways that regulate T cell metabolism remain poorly characterized. The PI3K/Akt signal transduction pathway is a central regulator of glycolytic metabolism in the immune system (13, 14). TCR- and CD28-dependent Akt activation results in increased Glut1-dependent glucose uptake and stimulation of glycolytic activity (13, 15). Akt also stimulates the mammalian target of rapamycin (mTOR), which regulates multiple metabolic pathways including protein translation, lipid synthesis, autophagy, and mitochondrial biogenesis and function in response to growth factor signals (16). mTOR has emerged as an important regulator of T cell differentiation and effector function (17). mTOR activity is required for the full effector function of both CD4<sup>+</sup> (18) and CD8<sup>+</sup> (19) T cells, and inhibition of mTOR signaling blocks the differentiation of CD4<sup>+</sup> T cells toward inflammatory Th cell lineages (18, 20) while stimulating the development of regulatory T cells (18, 21, 22). Similarly, inhibition of mTOR by rapamycin can promote the development of CD8<sup>+</sup> T cell memory (9, 10). Identifying the signal transduction pathways that regulate T cell metabolism, and in particular modulators of mTOR activity, may help us understand how metabolic regulation impacts T cell function.

The liver kinase B1 (LKB1) is a serine/threonine kinase that functions as part of an evolutionarily conserved energy-sensing

<sup>\*</sup>Department of Pediatrics, Duke University Medical Center, Durham, NC 27710;

<sup>†</sup>Department of Physiology, McGill University, Montreal, Quebec H3G 1Y6, Canada;

<sup>‡</sup>Goodman Cancer Research Centre, McGill University, Montreal, Quebec H3G 1Y6, Canada; <sup>§</sup>Department of Pharmacology and Cancer Biology, Duke University Medical Center, Durham, NC 27710; <sup>¶</sup>Department of Immunology, Duke University Medical Center, Durham, NC 27710; and <sup>||</sup>Sarah W. Stedman Center for Nutrition and Metabolism, Duke University Medical Center, Durham, NC 27710

<sup>1</sup>N.J.M. and J.B. contributed equally to this work.

Received for publication February 3, 2011. Accepted for publication August 15, 2011.

N.J.M. was supported by the Lawson Wilkins Pediatric Endocrine Society, Grants K12-HD-043494 and K08-DK-087944. J.B. was supported by the McGill Integrated Cancer Research Training Program. J.C.R. was supported by National Institutes of Health Grants R01s AI063345 and HL108006, the American Asthma Foundation (J.C.R. is a Bernard Osher Fellow of the American Asthma Foundation), the Leukemia and Lymphoma Society, and the Lupus Research Institute. R.G.J. was supported by grants from the Canadian Institutes for Health Research (MOP-93799), the Canadian Cancer Society (2010-700586), and The Arthritis Society.

Address correspondence and reprint requests to Dr. Russell G. Jones or Dr. Jeffrey C. Rathmell, Department of Physiology, McGill University, 3655 Promenade Sir William Osler, Room 705, Montreal, QC, H3G 1Y6, Canada (R.G.J.) or Department of Pharmacology and Cancer Biology, Duke University, DUMC 3813, 308 Research Drive, Durham, NC 27710 (J.C.R.). E-mail addresses: russell.jones@mcgill.ca (R.G.J.) and jeff.rathmell@duke.edu (J.C.R.)

The online version of this article contains supplemental material.

Abbreviations used in this article: 7-AAD, 7-aminoactinomycin D; ACC, acetyl-CoA carboxylase; AICAR, aminoimidazole carboxamide ribonucleotide; AMPK, AMP-activated protein kinase; 2-DG, 2-deoxyglucose; DN, double-negative; ICS, intracellular cytokine staining; LKB1, liver kinase B1; mTOR, mammalian target of rapamycin; mTORC1, mammalian target of rapamycin complex 1; PI, propidium iodide; qPCR, quantitative PCR; S6K, S6 kinase.

Copyright © 2011 by The American Association of Immunologists, Inc. 0022-1767/11/\$16.00

pathway to couple cellular bioenergetics to metabolic pathways important for cell growth control, proliferation, polarity, and stress resistance (23). LKB1 was first identified as the tumor suppressor responsible for Peutz-Jeghers syndrome, an autosomal dominant disorder that leads to intestinal hamartomas, mucocutaneous lesions, and an increased risk of spontaneous epithelial carcinomas (24, 25). LKB1 was recently found to have prominent roles in hematopoietic stem cell viability and renewal (26–28), suggesting a prominent function for LKB1 in hematopoietic cells. One of the targets that link LKB1 to cellular metabolism is the serine/threonine kinase AMP-activated protein kinase (AMPK), which can act to suppress mTOR activity (29). LKB1-dependent phosphorylation of the  $\alpha$  subunit of AMPK at Thr<sup>172</sup> stimulates AMPK activity, and LKB1 is essential for AMPK activation under conditions of bioenergetic stress (30–32). From a metabolic standpoint, AMPK promotes ATP conservation by activating pathways of catabolic metabolism and inhibiting anabolic processes that consume ATP (33). AMPK itself has been linked to stress resistance in lymphocytes (34).

Lymphocyte metabolism is a highly regulated process that plays a key role in the determination of T cell fate and function, and LKB1 is an established regulator of cell metabolism. However, although LKB1 has been shown to be important for thymocyte development (35, 36), its role in peripheral T cell metabolism, effector function, and mTOR regulation is not well defined. In this study, we show that LKB1 is a key regulator of glucose and lipid metabolism in T cells and is essential for both normal T cell development and peripheral T cell function. T cells lacking LKB1 displayed defects in cellular proliferation and viability in response to metabolic stress, yet showed increased activation and inflammatory cytokine production upon TCR stimulation. Moreover, loss of LKB1 in the T cell compartment led to an accumulation of IFN- $\gamma$ -producing CD8<sup>+</sup> and CD4<sup>+</sup> T cells in peripheral lymphoid organs. AMPK mediated a subset of LKB1 functions in peripheral T cells; AMPK $\alpha$ 1-deficient mice displayed normal thymocyte development and peripheral T cell homeostasis, but AMPK $\alpha$ 1<sup>-/-</sup> CD8<sup>+</sup> T cells displayed an activated phenotype with alterations in metabolism and cytokine production similar to LKB1-deficient T cells. Moreover, mTOR complex 1 (mTORC1) activity was elevated in LKB1- and AMPK $\alpha$ 1-deficient T cells, and elevated IFN- $\gamma$  production by these T cells could be reversed by rapamycin. Together, these data demonstrate a critical role for LKB1 in the regulation of peripheral T cell function and metabolism through regulation of AMPK-dependent and -independent pathways and mTORC1 signaling.

## Materials and Methods

### Mice

LKB1 floxed (*stk11-fl/fl*) mice, which carry an LKB1 allele with LoxP sites flanking exons 3–6 (37), were obtained from the National Cancer Institute (Frederick, MD). Rag1<sup>-/-</sup> mice, Lck-Cre mice, CD45.1 congenic mice, and Ub-Cre-ER<sup>T2</sup> mice were purchased from The Jackson Laboratory (Bar Harbor, MA), and Bcl-xL transgenic mice have been described (4). Mice deficient for AMPK $\alpha$ 1 (34) or floxed for AMPK $\alpha$ 2 (38) have been described previously. Lck-Cre mice were crossed with LKB1 floxed mice to generate *LKB1-fl/fl* Lck-cre<sup>+</sup> mutants or *LKB1-fl/fl* Lck-cre<sup>+</sup>, *LKB1-fl/fl* Lck-cre<sup>-</sup>, and *LKB1-fl/fl* Lck-cre<sup>-</sup> controls. Bcl-xL transgenic mice were crossed to the Lck-Cre LKB1 strains to generate *LKB1-fl/fl* Lck-cre<sup>+</sup> mice that express the Bcl-xL transgene. Mice were bred and maintained under specific pathogen-free conditions at Duke University and McGill University under approved protocols. Experiments were performed using mice between 8 and 20 wk of age.

### T cell purification and culture

T cells were purified from spleen and mesenteric lymph nodes by negative selection (StemCell Technologies, Vancouver, BC, Canada), and cells were cultured in IMDM or RPMI 1640 (Mediatech, Washington, DC) supplemented

with 10% FBS (Gemini Bio-Products, West Sacramento, CA), L-glutamine (Invitrogen, Chicago, IL), penicillin-streptomycin (Invitrogen), and 2-ME (Sigma-Aldrich, St. Louis, MO). CD44<sup>lo</sup> cells were isolated by negative selection via the addition of biotinylated anti-CD44 added to the enrichment mixture. Where indicated, IL-7 was supplemented into media at a concentration of 10 ng/ml (eBioscience, San Diego, CA). T cell stimulation was induced by culturing purified T cells on anti-CD3 $\epsilon$  (clone 145-2C11) and anti-CD28 (clone 37.51)-coated plates (eBioscience) and supplemented with IL-2 (PeproTech, Rocky Hill, NJ). CD4<sup>+</sup> T cell differentiation was performed as described previously (7). To activate Cre-ER and excise the floxed LKB1 transgene, isolated T cells were treated with 4-hydroxytamoxifen (0.5  $\mu$ M in ethanol; Sigma-Aldrich). 2-deoxyglucose (2-DG), aminoimidazole carboxamide ribonucleotide (AICAR), and rapamycin were obtained from Sigma-Aldrich. T cell proliferation was measured by [<sup>3</sup>H]thymidine incorporation (1  $\mu$ Ci/ml) 48 h post-activation as previously described (39) or via flow cytometry using CFSE (Molecular Probes, Eugene, OR).

### Flow cytometry, viability, and cell size measurements

Single-cell suspensions were stained with fluorescently conjugated Abs against murine CD4, CD8, CD3, Thy1.2, B220, CD19, CD25, CD44, CD45.1, CD62L, or CD69 (BD Biosciences, San Jose, CA). Cell viability was assessed via 7-aminoactinomycin D (7-AAD; Sigma-Aldrich) or propidium iodide (PI; Molecular Probes) exclusion as previously described (39). Flow cytometry was performed on an FACScan or LSR II (BD Biosciences) or Gallios (Beckman Coulter, Fullerton, CA) flow cytometer and analyzed with FlowJo software (Tree Star, Ashland, OR). Cell size was determined using a Coulter Z2 particle counter (Beckman Coulter). Bax activation was measured by fixing cells for 5 min with 0.25% paraformaldehyde in PBS, followed by incubation with an Ab specific to aa 12–24 (clone 6A7; BD Biosciences) as previously described (40).

### Cytokine measurements

Intracellular cytokine staining (ICS) was performed using specific fluorochrome-labeled mAbs (eBioscience) and flow cytometry as described (41). In brief, single-cell suspensions of spleen or lymph node cells were stimulated immediately ex vivo with PMA and ionomycin (Sigma-Aldrich) for 4 h. Brefeldin A was added to cultures after 2 h of stimulation. ICS was carried out using a Cytofix/Cytoperm kit (BD Biosciences) following the manufacturer's protocols. Cytokines in T cell supernatants were measured by capture ELISAs (eBioscience) using paired mAbs and recombinant standards, as previously described (15).

### Quantitative real-time PCR

Total mRNA was isolated from mouse T cells using TRIzol (Invitrogen), and cDNA was synthesized from 100 ng total RNA using the Superscript VILO cDNA Synthesis Kit (Invitrogen). Quantitative PCR (qPCR) was performed using SYBR Green qPCR SuperMix (Invitrogen) and an Mx3005 qPCR machine (Agilent Technologies) using the following primers: *glut1* forward, 5'-CTG GAC CTC AAA CTT CAT TGT GGG-3'; *glut1* reverse, 5'-GGG TGT CTT GTC ACT TTG GCT GG-3'; *hk2* forward, 5'-CCG TGG TGG ACA AGA TAA GAG AGA ACC-3'; and *hk2* reverse, 5'-GGA CAC GTC ACA TTT CGG AGC CAG-3'. All samples were normalized to  $\beta$ -actin mRNA levels.

### Immunoblotting

Cell lysate preparation, SDS-PAGE, electrophoretic transfer, immunoblotting, and development using ECL were conducted as described previously (1). To probe for Glut1, cells were lysed for 1 h on ice in PBS plus 1% Triton X-100 and 0.1% SDS containing protease inhibitors (BD Pharmingen). Otherwise, cells were lysed in RIPA buffer or AMPK Lysis Buffer (42) supplemented with protease and phosphatase inhibitors (Sigma-Aldrich). Primary Abs to AMPK (pT172-specific and total), phospho-acetyl-CoA carboxylase (ACC; pS79), p70 S6 kinase (S6K; pT389-specific and total), S6 ribosomal protein (pS235/236-specific and total), 4E-BP1 (pT37/46-specific and total), Bcl-xL, HKII, LDHA, and actin, as well as anti-rabbit and mouse HRP secondary Abs, were obtained from Cell Signaling Technology (Danvers, MA). Anti-LKB1 mouse mAb (Ley 37D/G6) was obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Glut1 rabbit Ab was obtained from Abcam (Cambridge, MA).

### Metabolic assays

Assays for glucose uptake, glycolytic flux, and mitochondrial-dependent  $\beta$ -oxidation of fatty acids were performed as previously described (9, 43, 44).



### Statistical analysis

Data are presented as mean  $\pm$  SD and were analyzed using paired Student *t* test. Unless stated otherwise, a confidence level of  $p < 0.05$  was considered statistically significant for all data.

## Results

### *LKB1 is required for T cell development and homeostasis*


To understand the role of the serine/threonine kinase LKB1 in T cell function, animals lacking expression of LKB1 in the T cell lineage were generated by crossing mice with LKB1 floxed (*LKB1-fl/fl*) (37) with mice expressing a Cre recombinase transgene under the control of the proximal *Lck* promoter (*Lck-cre*). Cre-expressing *LKB1-fl/fl* thymocytes and peripheral T cells displayed decreased expression of LKB1 protein as demonstrated by Western blotting (Fig. 1A). *Lck-cre<sup>+</sup>, LKB1-fl/fl* mice displayed dramatic reductions in both thymocyte (Fig. 1B) and peripheral T cell (Fig. 1C) numbers compared with control animals that retained endogenous LKB1 expression. Targeted T cell deletion of LKB1 also resulted in an increased proportion of CD4<sup>+</sup>CD8<sup>+</sup> double-negative (DN) thymocytes with accumulation in the DN3 (CD25<sup>hi</sup>, CD44<sup>lo</sup>) stage and decreased numbers of double-positive (CD4<sup>+</sup>CD8<sup>+</sup>) and CD4<sup>+</sup> and CD8<sup>+</sup> single-positive thymocytes, suggesting incomplete thymic selection (Fig. 1D). *Lck-cre<sup>+</sup>, LKB1-fl/fl* LKB1-deficient mice displayed a decrease in both the proportion (Fig. 1E) and total number (Fig. 1F) of CD4<sup>+</sup> and


CD8<sup>+</sup> peripheral T cells. LKB1-deficient T cells also displayed an increase in T cell size relative to Cre-negative controls (Fig. 1G).

### *LKB1-deficient T cells display defects in T cell proliferation and viability*

Despite an overall defect in thymic development, limited numbers of peripheral CD4<sup>+</sup> and CD8<sup>+</sup> T cells were observed in T cell-specific LKB1 mutant animals. To assess their functional status, LKB1-deficient T cells were purified from spleen, activated in vitro using anti-CD3 and anti-CD28 Abs, and T cell proliferation rates measured by thymidine incorporation. As seen in Fig. 2A, T cells from *Lck-cre<sup>+</sup>, LKB1-fl/fl* mice displayed decreased proliferative capacity when activated with anti-CD3 Abs alone or in combination with CD28 costimulation. The addition of rIL-2 led to a partial rescue in proliferation of LKB1-deficient (*LKB1-fl/fl; Lck-Cre<sup>+</sup>*) T cells (Supplemental Fig. 1A), but did not fully rescue T cell proliferation. To assess whether enhanced apoptosis could underlie the defective proliferation of LKB1-deficient T cells, we assessed levels of viability dye exclusion using PI. CFSE dilution profiles demonstrated that the majority of viable LKB1-deficient cells remained CFSE<sup>bright</sup> (Fig. 2B). However, LKB1-null T cells also displayed reduced viability relative to control T cells following activation (Fig. 2C). Although we cannot exclude cell death that occurred at the time of cell division, these data indicate that viable LKB1-deficient T cells exhibited delays in initiating cell division in response to TCR stimulation.

**FIGURE 1.** LKB1 is required for T cell development and peripheral homeostasis. *A*, LKB1 protein expression in LKB1-expressing (*Lck-cre<sup>-</sup>*) or LKB1-deficient (*Lck-cre<sup>+</sup>*) thymocytes and T cells. Total thymocytes and purified peripheral T cells were lysed and immunoblotted with Abs against LKB1 and actin. *B* and *C*, Summary of thymocyte (*B*) and peripheral T cell (*C*) counts in control (*Lck-cre<sup>-</sup>*, black bars) and LKB1 mutant (*Lck-cre<sup>+</sup>*, open bars) mice. Data are presented as means  $\pm$  SEM for a minimum of 10 mice per genotype. *D*, Representative flow plots for expression of CD4 and CD8 on total thymocytes (*top panel*) and CD44 and CD25 expression on CD4/CD8 DN thymocytes (*lower panel*) from control (*LKB1-fl/fl; Lck-cre<sup>-</sup>*) or LKB1 mutant (*LKB1-fl/fl; Lck-cre<sup>+</sup>*) animals. Numbers represent the percentage of cells in the indicated quadrant. *E*, Representative flow plots for B220 versus Thy1.2 expression (*upper panel*) and CD4 versus CD8 expression on total splenocytes (*lower panel*) from animals of the indicated genotypes. *F*, Summary of peripheral CD4 and CD8 T cell numbers from control (black bars) or LKB1 mutant (open bars) animals. *G*, Resting T cell size as measured by Coulter counter (Beckman Coulter). Black bars represent control T cells; open bars represent LKB1-deficient T cells.





**FIGURE 2.** LKB1-deficient T cells display defects in proliferation and viability. *A–C*, Proliferation of LKB1-deficient T cells. *A*, Peripheral T cells from control (black bars) or LKB1 mutant (open bars) mice were cultured with various concentrations of anti-CD3 and anti-CD28 Abs and proliferation measured by thymidine incorporation after 48 h. Data represent the mean  $\pm$  SD for samples in triplicate. *B*, Total T cells were activated with anti-CD3 (5  $\mu$ g/ml) and anti-CD28 (5  $\mu$ g/ml) for 72 h, and proliferation was measured by CFSE dilution. *C*, Viability of T cells in *B* was determined by flow cytometry using PI exclusion. *D* and *E*, Measurements of T cell apoptosis induced by metabolic stress. Thymocytes (*D*) or peripheral T cells (*E*) from control (black bars) or LKB1-mutant (open bars) animals were cultured with 2-DG (10 mM) and cell viability measured 24 h later by 7-AAD exclusion. Data are expressed as mean  $\pm$  SD for triplicate samples.  $**p < 0.05$ . *F*, Activated T cells were cultured with 2-DG (10 mM) for 18 h, and Bax activation was measured by flow cytometry.

Because altered development may have influenced activation of peripheral LKB1-deficient T cells, we examined proliferation following inducible deletion of LKB1 in mature peripheral T cells. T cells from *LKB1-fl/fl*, *Ub-Cre-ER<sup>+</sup>* or *LKB1-fl/fl*, *Ub-Cre-ER<sup>-</sup>* animals were cultured with rIL-7 and 4-hydroxytamoxifen for 4 d to promote deletion of the floxed LKB1 alleles in *Ub-Cre-ER<sup>+</sup>* T cells, and then T cells were labeled with CFSE and stimulated with anti-CD3 and anti-CD28 Abs for 48 h. CFSE dilution profiles showed that T cells with acute deletion of LKB1 displayed reduced levels of cell division (Supplemental Fig. 1*B*).

LKB1 is an important regulator of cellular stress responses (45) and plays an essential role in the maintenance of hematopoietic stem cell viability (27). To test whether loss of LKB1 could influence T cell viability, LKB1-deficient or control thymocytes and peripheral T cells were cultured in the presence of the glycolytic inhibitor 2-DG, and cell viability was assessed. Although the viability of control cells was largely unaffected by 2-DG treatment (10 mM), LKB1-deficient thymocytes (Fig. 2*E*) and T cells (Fig. 2*F*) displayed increased levels of cell death. Withdrawal of IL-2, which reduces glucose metabolism similar to the direct inhibition of glycolysis (9), promoted increased rates of cell death of LKB1-deficient T cells compared with controls (Supplemental Fig. 1*C*). In addition, treatment of activated LKB1-null T cells with 2-DG (Fig. 2*G*) or withdrawal from IL-2 (Supplemental Fig. 1*D*) promoted increased activation of the proapoptotic Bax protein, demonstrating an increase in apoptosis following metabolic stress in LKB1-null T cells.

#### LKB1 regulates T cell metabolism


Activated effector T cells rely heavily on glucose metabolism to survive, proliferate, and support specific effector functions (4, 5, 46). To evaluate the influence of LKB1 loss on T cell metabolism, the rates of glucose uptake and glycolysis were assessed in LKB1-deficient T cells. T cells lacking LKB1 displayed increased rates

of glucose uptake (Fig. 3*A*) and glycolytic flux (Fig. 3*B*) relative to control T cells, consistent with findings suggesting that LKB1 is required to regulate glucose metabolism in metabolic tissues (47). LKB1-deficient T cells displayed elevated expression of both *glut1* and *hk2* mRNA (Fig. 3*C*), as well as elevated Glut1 and hexokinase 2 protein levels (Fig. 3*D*, 3*E*). Glut1 protein expression was increased both at baseline and following TCR stimulation in the LKB1-null T cells as compared with controls (Fig. 3*D*).

We have previously shown that lipid oxidation can function as an alternate metabolic pathway in resting and memory T cells when glucose uptake is low or reduced or when cytokine availability is limiting (9). Thus, we assessed whether IL-2-dependent control of lipid metabolism was altered in LKB1-deficient T cells. Following cytokine withdrawal, which leads to downregulation of the glucose transporter Glut1 and reduced glycolysis (44), control T cells displayed an increase in their rate of lipid oxidation; however, the induction of lipid oxidation induced by IL-2 withdrawal was impaired in LKB1-deficient T cells (Fig. 3*F*). Collectively, these data suggest that LKB1 functions to coordinate glucose and lipid metabolism in T cells, and loss of LKB1 leads to deregulated metabolic function.

#### Loss of LKB1 results in the accumulation of activated inflammatory T cells

Given the disrupted metabolic and proliferative capacity of LKB1-deficient T cells, we next examined the effect of LKB1 loss on T cell effector function. Despite possessing fewer T cells overall, *Lck-cre<sup>+</sup>*, *LKB1-fl/fl* animals displayed increased numbers of CD4<sup>+</sup> and CD8<sup>+</sup> T cells with an activated phenotype (CD44<sup>hi</sup> CD62L<sup>lo</sup>) (Fig. 4*A*). Both the CD4<sup>+</sup> and CD8<sup>+</sup> T cell populations from *Lck-cre<sup>+</sup>*, *LKB1-fl/fl* mice displayed a higher proportion of CD44-expressing cells (Supplemental Fig. 2*A*). Moreover, both conditional and inducible deletion of LKB1 resulted in a pronounced increase in the surface expression of activation markers



**FIGURE 3.** LKB1 regulates T cell metabolism. *A*, Measurement of LKB1-dependent glucose uptake by T cells. Peripheral T cells from LKB1-expressing (black bars) or LKB1-mutant (open bars) mice were isolated from spleen and lymph nodes and glucose uptake assessed by [ $^3$ H]-2-DG uptake. *B*, Glycolytic flux in LKB1-deficient T cells. Control (black bars) or LKB1-null (open bars) T cells were isolated and glycolytic flux determined by the generation of tritiated water by enolase. Data are expressed relative to the glycolytic rate (nmol glucose/ $10^6$  cells/h) of control T cells. *C*, Control (black bars) or LKB1-deficient (open bars) T cells were stimulated with anti-CD3 and anti-CD28 Abs for 24 h and Glut 1 and hexokinase 2 mRNA expression determined by qPCR. Data are expressed relative to actin mRNA levels and normalized to mRNA levels in unstimulated T cells. *D*, Purified T cells from LKB1-expressing (*LKB1-fl/fl*; *Lck-cre*<sup>-</sup>) or LKB1-mutant (*LKB1-fl/fl*; *Lck-cre*<sup>+</sup>) mice were activated with anti-CD3 (5  $\mu$ g/ml) and anti-CD28 (5  $\mu$ g/ml) Abs for 24 h. Lysates were resolved by SDS-PAGE and immunoblotted for Glut1 and actin. *E*, Western blot of hexokinase 2 (HKII) expression in control (*LKB1-fl/fl*; *Lck-cre*<sup>-</sup>) or LKB1-deficient (*LKB1-fl/fl*; *Lck-cre*<sup>+</sup>) T cells 24 h following anti-CD3 and anti-CD28 treatment. *F*, Measurement of lipid oxidation in LKB1-deficient T cells. Activated T cells were cultured in the presence (black bars) or absence (open bars) of rIL-2 for 14 h, and the liberation of tritiated water from [ $^3$ H] palmitate (cpm/ $10^6$  cells) was measured 6 h later. Data are presented as the mean  $\pm$  SD for samples in triplicate.

CD25, CD44, and CD69 following TCR ligation (Supplemental Fig. 2*B*, 2*C*).

Given the heightened activation state of LKB1-null T cells, we next examined cytokine production by T cells lacking LKB1. Peripheral T cells from *LKB1-fl/fl*; *Lck-Cre*<sup>+</sup> or *LKB1-fl/fl*; *Lck-Cre*<sup>-</sup> control animals were isolated, stimulated with anti-CD3 and anti-CD28 Abs, and IFN- $\gamma$  and IL-17A in culture supernatants was measured. Despite lower overall levels of proliferation (Fig. 2*A*), LKB1-deficient T cells displayed elevated IFN- $\gamma$  and IL-17A production relative to control T cells (Fig. 4*B*). We next isolated naive (CD44<sup>lo</sup>) CD4<sup>+</sup> and CD8<sup>+</sup> T cells from LKB1 mutant and control mice and examined TCR-induced cytokine production by individual T cell subsets. LKB1-deficient CD8<sup>+</sup> T cells produced increased amounts of IFN- $\gamma$  and IL-17A upon activation relative to LKB1-expressing T cells (Fig. 4*C*). LKB1-deficient CD4<sup>+</sup> T cells also displayed a slight enhancement in IFN- $\gamma$  production (Supplemental Fig. 2*D*). Finally, we assessed the capacity of naive CD4<sup>+</sup> LKB1-null T cells to differentiate into Th1, Th2, or Th17 effector cells in vitro. T cells lacking LKB1 displayed enhanced differentiation toward Th1 and Th17 CD4<sup>+</sup> T cell lineages (Fig. 4*D*).

We next analyzed whether LKB1 deficiency could alter the number of cytokine-producing T cells in vivo. Total splenocytes or lymph node cells were isolated from *Lck-cre*<sup>+</sup>, *LKB1-fl/fl* animals or littermate controls, stimulated immediately ex vivo with PMA and ionomycin, and the presence of IFN- $\gamma$  and IL-2-producing cells was measured by ICS. Peripheral CD8<sup>+</sup> (Fig. 4*E*) and CD4<sup>+</sup> (Fig. 4*F*) T cells from T cell-specific LKB1-null animals were enriched for IFN- $\gamma$ -producing cells as determined by ICS. In


particular, a significant increase in the level of IFN- $\gamma$ /IL-2 double-positive CD8<sup>+</sup> T cells was observed in LKB1 T cell-deficient animals (Fig. 4*E*), suggesting that LKB1 functions as a negative regulator of IFN- $\gamma$ -producing T cells in vivo.

#### *Bcl-xL* rescues decreased thymocyte and peripheral T lymphocyte cellularity caused by LKB1 deficiency

In principle, deletion of LKB1 at the DN stage of thymocyte development may promote low T lymphocyte cellularity (Fig. 1) due to either inefficient thymocyte differentiation or reduced viability. The antiapoptotic Bcl-2 family protein Bcl-xL is a key regulator of thymocyte viability (48), and reduced Bcl-xL levels have been linked to LKB1 deficiency (35). Indeed, thymocytes with deletion of both alleles of LKB1 gene displayed decreased levels of Bcl-xL protein (Fig. 5*A*). To assess whether restoring Bcl-xL levels could rescue the defect in T cell development in T cell LKB1-deficient mice, we generated *Lck-cre*<sup>+</sup>, *LKB1-fl/fl* mice that also expressed a T cell-specific Bcl-xL transgene and analyzed the central and peripheral T cell compartments of these mice by flow cytometry. Transgenic expression of Bcl-xL rescued total thymocyte cellularity in T cell-specific LKB1-deficient mice to wild-type levels and increased splenic T cell cellularity above wild-type levels regardless of LKB1 status (Fig. 5*B*). Expression of Bcl-xL also restored the positive selection of CD4<sup>+</sup> and CD8<sup>+</sup> T cells in the thymus (Fig. 5*C*) and circumvented the developmental block at the DN3 (CD25<sup>hi</sup>CD44<sup>lo</sup>) stage of thymocyte development caused by targeted deletion of LKB1 in thymocytes (Fig. 5*D*).



**FIGURE 4.** LKB1-deficient T cells display a hyper-activated phenotype. **A**, CD4<sup>+</sup> and CD8<sup>+</sup> splenocytes from LKB1-expressing (*LKB1-fl/fl*, *Lck-cre*<sup>-</sup>) or LKB1-deficient (*LKB1-fl/fl*, *Lck-cre*<sup>+</sup>) mice were analyzed for CD44 and CD62L expression by flow cytometry. **B**, IFN- $\gamma$  and IL-17A production by LKB1-null T cells. Control (black bars) or LKB1-deficient (open bars) T cells were cultured with anti-CD3 Abs in the presence or absence of CD28 costimulation for 24 h, and IFN- $\gamma$  and IL-17A in culture supernatants was measured by ELISA. **C**, IFN- $\gamma$  (left panel) and IL-17A (right panel) production by anti-CD3/CD28-activated CD44<sup>lo</sup>CD8<sup>+</sup> T cells from *LKB1-fl/fl*, *Lck-cre*<sup>-</sup> or *LKB1-fl/fl*, *Lck-cre*<sup>+</sup> mice as measured by ELISA. **D**, CD44<sup>lo</sup>CD4<sup>+</sup> T cells from LKB1-expressing (*LKB1-wt/wt*, *Lck-cre*<sup>-</sup>) or LKB1-deficient (*LKB1-fl/fl*, *Lck-cre*<sup>+</sup>) mice were cultured in Th1, Th2, or Th17 skewing conditions and examined for effector function by intracellular staining for IL-2, IFN- $\gamma$ , IL-4, and IL-17. **E** and **F**, Frequency of IFN- $\gamma$  and IL-2 producing T cells in LKB1-mutant mice. Splenocytes (Spleen) or lymph node (LN) cells from LKB1-expressing (*LKB1-fl/fl*, *Lck-cre*<sup>-</sup>) or LKB1-deficient (*LKB1-fl/fl*, *Lck-cre*<sup>+</sup>) animals were stimulated with PMA and ionomycin for 4 h, and cytokine-producing cells were identified by ICS and flow cytometry. CD8<sup>+</sup> T cells were analyzed for IFN- $\gamma$  versus IL-2 expression (**E**), whereas CD4<sup>+</sup> cells were analyzed for surface expression of CD44 and intracellular IFN- $\gamma$  (**F**). The gated population in **F** refers to the percentage of endogenous IFN- $\gamma$ -producing CD4<sup>+</sup> T cells.




One explanation for the accumulation of activated T cells in T cell-specific LKB1-null animals is increased homeostatic T cell proliferation brought on by the reduced T cell compartment in these mice. To test this possibility, CFSE-labeled CD45.1 congenic donor T cells were adoptively transferred into either LKB1-deficient hosts or various control mice, and the level of homeostatic proliferation was assessed by CFSE dye dilution 1 wk posttransfer. Minimal levels of homeostatic proliferation by the CD45.1 donor T cell population were detected in *LKB1-fl/fl*, *Lck-cre*<sup>+</sup>, *Bcl-xL-Tg* mice, with significant levels of T cell division observed only in *Rag1*<sup>-/-</sup> hosts (Supplemental Fig. 3A). Moreover, LKB1-dependent thymic cellularity was restored by expression of a Bcl-xL transgene (Fig. 5B), which permitted the direct examination of LKB1-deficient peripheral T cell function in the absence of lymphopenia. Despite equivalent cell numbers as control Bcl-xL transgenic animals, LKB1-deficient T cells expressing Bcl-xL continued to display elevated CD44 expression at baseline, as well as increased CD44, CD25, and CD69 expression upon stimulation (Supplemental Fig. 3B). Similarly, LKB1-deficient T cells expressing Bcl-xL displayed increased IFN- $\gamma$  production upon T cell activation relative to control Bcl-xL-transgenic T cells (Fig. 5E). Thus, transgenic expression of Bcl-xL reversed the developmental and homeostatic defects, but not the activated T cell phenotype, in T cell-specific LKB1-deficient mice (Fig. 5), suggesting a cell-intrinsic role for LKB1 in the regulation of T cell activation.

#### AMPK $\alpha$ 1 regulates T cell viability and metabolism, but not TCR-mediated proliferation

The most well-defined physiological target of LKB1 is the serine/threonine kinase AMPK (45). Phosphorylation of AMPK $\alpha$  at Thr<sup>172</sup> and phosphorylation of its downstream target ACC occurs rapidly following TCR stimulation (Fig. 6A). As seen in Fig. 6B, LKB1-deficient T cells (*LKB1-fl/fl*, *Lck-cre*<sup>+</sup>) displayed impaired AMPK $\alpha$  and ACC phosphorylation following stimulation with the AMPK agonist AICAR, in agreement with previous work suggesting that T cells lacking LKB1 are defective for AMPK $\alpha$  phosphorylation at Thr<sup>172</sup> (36). To determine if the phenotype of LKB1-deficient T cells could be attributed to the loss of AMPK signaling, we examined T lymphocyte development and function in animals lacking *prkaa1*, the gene that encodes AMPK $\alpha$ 1. AICAR-induced AMPK activation was completely abolished in thymocytes lacking the  $\alpha$ 1 subunit of AMPK (Supplemental Fig. 4A), suggesting that T cells, like other hematopoietic cells, predominantly express the  $\alpha$ 1 catalytic subunit of the trimeric AMPK complex (49). In agreement with previous work (34), AMPK $\alpha$ 1-deficient mice displayed no overt defects in thymocyte development or peripheral T cell subsets (Supplemental Fig. 4B) and displayed normal proliferative responses to TCR and CD28 stimulation (Fig. 6C).

We next examined the effects of AMPK $\alpha$ 1 loss on T cell viability and metabolism. Activated T cells from AMPK $\alpha$ 1-deficient

**FIGURE 5.** Transgenic expression of Bcl-xL rescues thymocyte development but not aberrant T cell activation and cytokine production in LKB1-mutant animals. **A**, Bcl-xL protein expression in LKB1-deficient thymocytes. Lysates from homozygous LKB1-deficient (KO), LKB1 heterozygous (HET), or wild-type (WT) thymocytes were resolved by SDS-PAGE and immunoblotted using Abs to Bcl-xL and actin. **B**, Thymocyte and T cell numbers from LKB1-expressing (*LKB1-fl/wt*, *Lck-cre*<sup>+</sup>) or LKB1-deficient (*LKB1-fl/fl*, *Lck-cre*<sup>+</sup>) mice with or without expression of a Bcl-xL transgene. Data are expressed as mean  $\pm$  SEM for a minimum of five animals per genotype. **C** and **D**, Expression profiling of thymocytes from LKB1-mutant animals expressing a Bcl-xL transgene. Thymocytes from LKB1-expressing (*LKB1-fl/wt*, *Lck-cre*<sup>+</sup>) or LKB1-deficient (*LKB1-fl/fl*, *Lck-cre*<sup>+</sup>) animals with (+) or without (–) expression of a Bcl-xL transgene were analyzed for CD4 versus CD8 expression on total thymocytes (**C**) or CD25 versus CD44 expression on CD4<sup>+</sup> CD8<sup>+</sup> DN thymocytes (**D**) by flow cytometry. **E**, IFN- $\gamma$  production of LKB1-null T cells overexpressing Bcl-xL. Bcl-xL–expressing T cells from LKB1-expressing (black bars) or LKB1-deficient (open bars) animals were stimulated with anti-CD3 and anti-CD28 Abs, and IFN- $\gamma$  in supernatants was measured at 48 h by ELISA. Data are expressed as the mean  $\pm$  SD for triplicate cultures.



or control animals were cultured in the absence of IL-2 (Fig. 6D) or subjected to metabolic stress by 2-DG treatment (Fig. 6E) and cell viability measured. Similar to LKB1 deficiency, AMPK $\alpha$ 1-deficient T cells displayed reduced viability relative to control cells in response to these stresses (Fig. 6D, 6E). However, the expression of Bcl-xL protein was equivalent in both AMPK $\alpha$ 1<sup>-/-</sup> and control T cells (Fig. 6F). Similar to LKB1-deficient T cells, AMPK $\alpha$ 1<sup>-/-</sup> T cells displayed a 3-fold increase in the basal glycolytic rate of resting T cells ex vivo (Fig. 6G) and increased levels of *glut1* and *hk2* mRNA (Fig. 6H). AMPK $\alpha$ 1-deficient T cells also displayed elevated expression of hexokinase 2 protein (Fig. 6I). Thus, loss of AMPK activity appears to phenocopy the effect of LKB1 deficiency on T cell metabolism (Fig. 3).

#### AMPK $\alpha$ 1 regulates CD8<sup>+</sup> T cell activation and cytokine production


To assess the effect of AMPK $\alpha$ 1 loss on T cell effector function, we measured the level of activation and cytokine production of AMPK $\alpha$ 1-deficient T cells following TCR stimulation. CD8<sup>+</sup> T cells from AMPK $\alpha$ 1<sup>-/-</sup> mice displayed elevated CD44 expression both prior to and following anti-CD3 and anti-CD28 stimulation (Fig. 7A). Similar to LKB1-deficient T cells, increased levels of secreted IFN- $\gamma$  and IL-17A (Fig. 7B) were detected in AMPK $\alpha$ 1<sup>-/-</sup> cultures following TCR stimulation. To ensure that these effects were not due to the presence of previously activated cells, naive (CD44<sup>lo</sup>) CD4<sup>+</sup> and CD8<sup>+</sup> T cells were

isolated and activated with anti-CD3 and anti-CD28 Abs, and supernatants were analyzed for cytokine production by ELISA. Interestingly, AMPK $\alpha$ 1<sup>-/-</sup> CD8<sup>+</sup> T cells displayed elevated IFN- $\gamma$  and IL-17 production (Fig. 7C), whereas IFN- $\gamma$  production by AMPK $\alpha$ 1<sup>-/-</sup> CD4<sup>+</sup> T cells did not differ significantly from control cells (Supplemental Fig. 4C). Elevated production of IFN- $\gamma$  by AMPK $\alpha$ 1<sup>-/-</sup> CD8<sup>+</sup> T cells was also detected by intracellular cytokine staining (Fig. 7D). We next induced naive AMPK $\alpha$ 1<sup>-/-</sup> CD4<sup>+</sup> T cells to differentiate into Th1, Th2, and Th17 effector cells and observed no discernible difference in cytokine production between AMPK $\alpha$ 1-deficient or control CD4<sup>+</sup> T cells (Fig. 7E). Altogether, these data suggest that, in contrast to LKB1 (Fig. 4), AMPK $\alpha$ 1 specifically regulates the production of inflammatory cytokines by CD8<sup>+</sup> but not CD4<sup>+</sup> T cells.

#### LKB1-AMPK $\alpha$ 1-dependent control of mTORC1 regulates IFN- $\gamma$ production

A major downstream effect of LKB1-AMPK signaling is the inactivation of the mTOR pathway (45). To determine the influence of reduced LKB1 signaling on mTORC1 activity, LKB1-null T cells were stimulated with anti-CD3 and anti-CD28 for 24 h, and phosphorylation of the downstream mTORC1 targets S6K, rS6, and 4EBP1 were assessed by Western blot. T cells lacking LKB1 expression displayed elevated S6K, rS6, and 4EBP1 phosphorylation following activation (Fig. 8A). Similar increases in S6K, rS6, and 4EBP1 phosphorylation were observed in T cells






**FIGURE 6.** AMPK regulates T cell viability and metabolism, but not proliferation. *A*, Time course of AMPK activation following TCR and CD28 stimulation. Purified T cells were activated with anti-CD3 and anti-CD28 Abs for the indicated times, and lysates resolved by SDS-PAGE were immunoblotted for phospho-AMPKα [pAMPK (T172)], phospho-ACCα [pACCα (S79)], and total AMPKα. *B*, Western blot of phospho-AMPKα (T172) and phospho-ACCα (S79) levels in control (Lck-Cre<sup>-</sup>) or LKB1-deficient (Lck-Cre<sup>+</sup>) T cells following AICAR treatment (1 mM, 1 h). *C*, Peripheral T cells from wild-type (*AMPKα1*<sup>+/+</sup>, black bars) or *AMPKα1*-deficient (*AMPKα1*<sup>-/-</sup>, open bars) mice were stimulated with anti-CD3 Abs (1 μg/ml) with or without anti-CD28 costimulation (0.5 μg/ml). Proliferation was measured by thymidine incorporation after 48 h. Data represent the mean ± SD for triplicate samples. *D* and *E*, Measurements of apoptosis resistance of *AMPKα1*-deficient T cells. Peripheral T cells from control (black bars) or *AMPKα1*<sup>-/-</sup> (open bars) animals were cultured in the presence (+) or absence (-) of rIL-2 (*D*) or treated with 10 mM 2-DG (*E*), and cell viability was measured 24 h later by 7-AAD exclusion. Data are expressed as mean ± SD for triplicate samples. *F*, Western blot of Bcl-xL expression in *AMPKα1*<sup>+/+</sup> and *AMPKα1*<sup>-/-</sup> thymocytes. *G*, Measurement of relative glycolytic flux in *AMPKα1*-deficient T cells. Control (black bars) or *AMPKα1*<sup>-/-</sup> (open bars) T cells were isolated by magnetic sorting and glycolytic flux determined as in Fig. 3. *H*, Relative Glut1 and hexokinase 2 mRNA levels in *AMPKα1*<sup>+/+</sup> (black bars) and *AMPKα1*<sup>-/-</sup> (open bars) T cells 24 h following anti-CD3/CD28 treatment as determined by qPCR. Data are expressed relative to actin mRNA levels, and normalized to mRNA levels in unstimulated T cells. *I*, Western blot of hexokinase 2 (HKII) expression in *AMPKα1*<sup>+/+</sup> and *AMPKα1*<sup>-/-</sup> T cells 24 h following anti-CD3/CD28 treatment.

lacking AMPKα1 (Fig. 8*B*). Baseline phosphorylation of rS6 and 4EBP1 was also elevated in unstimulated *AMPKα1*<sup>-/-</sup> T cells (Fig. 8*B*). Inhibition of mTORC1 signaling by the addition of rapamycin to T cell cultures ablated the effect of LKB1- or AMPKα1-deficiency on enhanced IFN-γ production (Fig. 8*C*, 8*D*). Thus, LKB1 and AMPKα1 appear to play a key role in suppressing mTORC1 signaling, and elevated mTORC1 activity appears to promote the enhanced cytokine production by LKB1 and AMPKα1-deficient T cells.

## Discussion

LKB1 functions at the center of a signaling axis that couples cellular bioenergetics and environmental stimuli with cell growth

control and metabolism. In this study, we demonstrate that LKB1 plays a critical role in regulating T cell development and peripheral T cell function. Animals lacking LKB1 in the T cell compartment displayed a block in thymocyte development at the DN3 stage, resulting in decreased thymic cellularity and a reduced peripheral T cell compartment. LKB1-deficient T cells displayed defective proliferation in response to TCR and costimulatory signals, were increasingly susceptible to apoptosis induced by metabolic stressors or cytokine withdrawal, and displayed perturbations in glucose and lipid metabolism. Surprisingly, LKB1-deficient T cells that did reach the periphery, either naturally or through forced expression of Bcl-xL to block aberrant cell death due to the lack of LKB1, displayed an activated phenotype characterized by increased ex-



**FIGURE 7.** Loss of AMPK $\alpha$ 1 promotes hyperactivation of CD8<sup>+</sup> T cells. **A**, CD44 surface expression on AMPK $\alpha$ 1-deficient T cells. T cells isolated from control (gray histogram) or AMPK $\alpha$ 1<sup>-/-</sup> (open histogram) mice were left unstimulated (*left panel*) or anti-CD3/CD28 treated (*right panel*) for 24 h. CD44 surface expression on CD8<sup>+</sup> T cells was analyzed by flow cytometry. **B**, Inflammatory cytokine production by activated AMPK $\alpha$ 1-deficient T cells. Wild-type (black bars) or AMPK $\alpha$ 1-null (white bars) T cells were activated with anti-CD3 Abs in the presence or absence of anti-CD28 costimulation (0.5  $\mu$ g/ml). Culture supernatants were harvested and analyzed for IFN- $\gamma$  or IL-17A levels by ELISA. **C**, IFN- $\gamma$  (*left panel*) and IL-17A (*right panel*) production by anti-CD3/CD28-activated CD44<sup>lo</sup>CD8<sup>+</sup> T cells from AMPK $\alpha$ 1<sup>+/+</sup> (black bars) or AMPK $\alpha$ 1<sup>-/-</sup> (white bars) mice. Cytokines were measured by ELISA 24 h poststimulation. **D**, IFN- $\gamma$  production by AMPK-deficient T cells. Splenocytes from AMPK $\alpha$ 1<sup>+/+</sup> or AMPK $\alpha$ 1<sup>-/-</sup> animals were activated with anti-CD3 Abs, and IFN- $\gamma$ -producing cells were identified by ICS and flow cytometry. Surface expression of CD44 versus intracellular IFN- $\gamma$  is displayed for CD8<sup>+</sup> cells. **E**, Th differentiation of AMPK $\alpha$ 1<sup>-/-</sup> T cells in vitro. AMPK $\alpha$ 1<sup>+/+</sup> or AMPK $\alpha$ 1<sup>-/-</sup> CD44<sup>lo</sup>CD4<sup>+</sup> T cells were cultured in Th1, Th2, or Th17 skewing conditions and examined for effector function after 4 d of culture by intracellular staining for IFN- $\gamma$ , IL-4, and IL-17.


pression of activation markers and elevated production of inflammatory cytokines. In particular, loss of LKB1 enhanced IFN- $\gamma$  and IL-17 production by CD8<sup>+</sup> T cells and promoted enhanced differentiation of CD4<sup>+</sup> T cells toward Th1 and Th17 effector subtypes. In contrast, loss of AMPK $\alpha$ 1, a prominent downstream target of LKB1, only partially recapitulated the phenotype of LKB1-deficient T cells. AMPK $\alpha$ 1<sup>-/-</sup> T cells displayed similar defects in T cell viability and metabolism, whereas T cell development and proliferative responses were normal. Although IFN- $\gamma$  and IL-17 production by AMPK $\alpha$ 1<sup>-/-</sup> T cells was also enhanced, this was largely due to deregulated CD8<sup>+</sup> T cell function in the absence of AMPK $\alpha$ 1. The mTORC1 pathway was hyperactivated in both LKB1- and AMPK $\alpha$ 1-deficient T cells, and inhibition of this pathway with rapamycin was sufficient to block elevated IFN- $\gamma$  production by these cells. Together our results establish the LKB1–AMPK signaling axis as a critical regulator of peripheral T cell viability and metabolism and a negative regulator of T cell inflammatory cytokine production through regulation of mTORC1 activity (Fig. 8E).

The present study, coupled with lines of evidence from other groups, indicates that LKB1 plays a multifaceted role in T cell development and peripheral function. Our data support previous work focused largely on the role of LKB1 in thymocyte development (36) and thymocyte survival (35) and suggest that LKB1 is a cell-autonomous factor essential for T cell development. We also demonstrate that LKB1 functions largely to

regulate thymocyte viability, as both thymocyte cellularity and the impaired DN to double-positive transition are restored when apoptosis is blocked in T cell-specific LKB1-null mice by transgenic overexpression of Bcl-xL. This suggests that LKB1 is not essential for the T cell differentiation process itself, but rather may function to support cell survival during thymocyte development. Indeed, thymocytes and T cells lacking LKB1 display increased susceptibility to apoptosis induced by metabolic stress or cytokine withdrawal. Collectively, our data suggest that loss of LKB1 in T cells decreases overall T cell fitness leading to decreased viability, similar to that observed in hematopoietic stem cells (26–28). It is important to note, however, that the LKB1 mouse strain used in this study, which targets exons 3–6 of the *stk11* gene for LKB1 (37), is different from the animals used by Tamas et al. (50), which targets exons 4–7 and results in the generation of a hypomorphic mutation that decreases LKB1 expression in the absence Cre recombinase. Consequently, one cannot rule out the possibility that hypomorphic LKB1 expression in other tissues, such as thymic epithelium, could contribute to the phenotypes previously described (36). Together, however, these data suggest that LKB1 functions to couple environmental cues to T cell survival either through the regulation of cellular metabolism and bioenergetics and/or Bcl-xL levels.

AMPK is one of the key physiological effectors downstream of LKB1. However, our data suggest that LKB1 exerts effects on T cell biology through both AMPK-dependent and -independent

**FIGURE 8.** mTORC1 activity is deregulated in LKB1- and AMPK $\alpha$ 1-deficient T cells. **A**, Examination of mTORC1 signaling in LKB1-deficient T cells by Western blot. T cells were isolated from control (*Lck-cre*<sup>-</sup>) or LKB1-deficient (*Lck-cre*<sup>+</sup>) mice, left unstimulated (-) or stimulated (+) with anti-CD3 and anti-CD28 Abs for 24 h, and lysates resolved by SDS-PAGE. Phosphorylated and total levels of S6K, rS6, and 4E-BP were measured by immunoblotting. Protein loading was assessed by measuring total LDHA levels. **B**, mTORC1 signaling in AMPK $\alpha$ 1-deficient T cells. T cells from *AMPK $\alpha$ 1*<sup>+/+</sup> or *AMPK $\alpha$ 1*<sup>-/-</sup> mice were activated and lysates analyzed for mTORC1 signaling as in **A**. **C** and **D**, Analysis of LKB1- and AMPK-dependent IFN- $\gamma$  production in the presence of rapamycin. LKB1-deficient (**C**) or AMPK $\alpha$ 1-deficient (**D**) T cells were stimulated in the presence (+) or absence (-) of rapamycin (Rapa; 25 nM), and IFN- $\gamma$  in culture supernatants was measured by ELISA at 24 h. Black bars represent control T cell cultures for each experiment. Data are presented as the mean  $\pm$  SD for samples in triplicate. **E**, Integrated model for the function of LKB1 in T cell biology.



pathways (Fig. 8E). Although AMPK $\alpha$ 1 was largely dispensable for thymocyte development and T cell proliferation, both LKB1 and AMPK $\alpha$ 1 mutant T cells displayed defects in glucose and lipid metabolism and stress-induced apoptosis, suggesting these biological processes are regulated through a common pathway. Similarly, LKB1 and AMPK $\alpha$ 1 both function as negative regulators of inflammatory cytokine production. However, AMPK appears only to regulate CD8<sup>+</sup> T cell activation and cytokine production, whereas LKB1 influences both CD4<sup>+</sup> and CD8<sup>+</sup> T cell function. AMPK $\alpha$ 1 is just 1 out of 13 enzymes of the AMPK-related family of kinases activated by LKB1 (51) and, as such, may only represent a fraction of LKB1-dependent processes in T cells.

One striking aspect of the activated T cell phenotype of LKB1 mutant mice is the appearance of enhanced cytokine-producing CD4<sup>+</sup> and CD8<sup>+</sup> T cells with diminished proliferative capacity. LKB1 has previously been shown to be a negative regulator of cell cycle entry (52), so loss of proliferative capacity with LKB1 loss in T cells runs counter to other cellular systems. One possible explanation for the increased activation of LKB1-null T cells could be homeostatic proliferation driven by lymphopenia due to the reduced T cell compartment in *Lck-Cre*<sup>+</sup>, *LKB1-*fl/fl** mice (53, 54). However, expression of a Bcl-xL transgene rescues T cell numbers in LKB1 mutant mice but does not affect the heightened

activation profile of or cytokine production by LKB1-deficient T cells. Moreover, the observation that AMPK $\alpha$ 1-deficient animals display normal T cell development but enhanced T cell activation and cytokine production upon stimulation suggests that lymphopenia may not drive the activated phenotype of LKB1-deficient cells. Rather, altered LKB1/AMPK-dependent processes such as cellular metabolism or deregulated signaling downstream of LKB1 (i.e., increased mTORC1 activity) may drive T cell activation and cytokine production. Again, the complete scope of LKB1-dependent processes in T cells may only partially depend on AMPK, with AMPK $\alpha$ 1 coupling LKB1 to the regulation of effector cytokine production while being dispensable for proliferation (Fig. 8E).

Our data suggest that regulation of mTORC1 signaling downstream of LKB1/AMPK is critical for inflammatory cytokine production. mTORC1 activity is elevated in LKB1- and AMPK $\alpha$ 1-deficient T cells and rapamycin can reverse the elevated IFN- $\gamma$  production by these T cells. Thus, LKB1-AMPK signaling represents a novel mechanism for regulation of mTOR-dependent effector functions in T cells. Increased mTORC1 activity appears to generally influence T cell viability and effector function. T cell-specific deletion of Tsc1, a tumor suppressor that lies downstream of LKB1-AMPK signaling, results in increased mTORC1 activity and is required for the maintenance of T cell



viability and quiescence (55–57). Whether LKB1–AMPK signaling influences T cell function through control of metabolism remains unclear. Increased glucose uptake, such as that observed in LKB1- and AMPK $\alpha$ 1-deficient T cells, can enhance T cell activation and cytokine production (5). Thus, the increase in glucose metabolism associated with loss of LKB1–AMPK signaling may drive a feed-forward loop that promotes the development of proinflammatory effector T cells, resulting in the accumulation of activated IFN- $\gamma$ -producing cells. The engagement of alternate metabolic pathways such as lipid oxidation (9), which is defective in LKB1- and AMPK $\alpha$ 1-deficient T cells, may also contribute to this process. The activation of mTORC1 signaling, either through PI3K/Akt-mediated signal transduction or suppression of LKB1–AMPK activity, as shown in this study, appears to favor the development of inflammatory effector T cells. Our data add to the growing body of evidence that AMPK signaling can influence inflammatory processes (58, 59).

The signal transduction pathways that regulate LKB1 and AMPK activity in T cells remain unclear. LKB1 is considered to have constitutive kinase activity and may be limited by the availability and/or localization of its substrates (23). More recent data have suggested that LKB1 is phosphorylated by Lck following TCR stimulation and associates with downstream TCR signaling components linker for activation of T cells and phospholipase C $\gamma$ 1 (60). AMPK itself can be activated by other kinases in addition to LKB1, including CaMKK2 (61, 62), which may promote AMPK activation downstream of TCR stimulation (49). TCR activation stimulates the PI3K/Akt/mTOR pathway to promote Glut1 cell-surface trafficking and glycolysis, but upregulation of this metabolic pathway following TCR ligation takes several hours (5, 63). AMPK is rapidly activated following TCR stimulation (49) and thus may function to promote rapid increases in ATP generation through alternative metabolic pathways prior to the induction of glycolysis. Alternatively, the increased metabolic demands of T cell activation may promote metabolic stress conditions that in turn activate AMPK via LKB1. Developing and resting T cells require extrinsic signals mediated through the TCR and IL-7R to maintain glucose metabolism (4, 64), and these receptors may signal through LKB1 to regulate basal T cell metabolism. Further work must be done to characterize the signaling pathways both upstream and downstream of LKB1.

Together, our data establish LKB1 as a critical regulator of T cell development and peripheral T cell function. Our data suggest that inhibition of the LKB1–AMPK signaling axis disrupts T cell metabolism and promotes the development of proinflammatory T cells through hyperactivation of mTORC1 signaling. Deficiency in LKB1 or AMPK $\alpha$ 1 leads to a reciprocal phenotype to mTOR deficiency (17), supporting the model that these pathways act in opposition as critical regulators of T cell metabolism and function. Thus, manipulation of LKB1–AMPK signaling may provide a new means by which to modulate inflammation and autoimmunity.

## Acknowledgments

We thank Benoit Viollet for the generous gift of AMPK $\alpha$ 1- and AMPK $\alpha$ 2-deficient mice, Erika Pearce and Pamela Ohashi for critical reading of the manuscript, and members of the Rathmell and Jones Laboratories for helpful discussion and feedback.

## Disclosures

The authors have no financial conflicts of interest.

## References

- Jones, R. G., and C. B. Thompson. 2007. Revving the engine: signal transduction fuels T cell activation. *Immunity* 27: 173–178.
- Roos, D., and J. A. Loos. 1973. Changes in the carbohydrate metabolism of mitogenically stimulated human peripheral lymphocytes. II. Relative importance of glycolysis and oxidative phosphorylation on phytohemagglutinin stimulation. *Exp. Cell Res.* 77: 127–135.
- Bental, M., and C. Deutsch. 1993. Metabolic changes in activated T cells: an NMR study of human peripheral blood lymphocytes. *Magn. Reson. Med.* 29: 317–326.
- Rathmell, J. C., M. G. Vander Heiden, M. H. Harris, K. A. Frauwirth, and C. B. Thompson. 2000. In the absence of extrinsic signals, nutrient utilization by lymphocytes is insufficient to maintain either cell size or viability. *Mol. Cell* 6: 683–692.
- Jacobs, S. R., C. E. Herman, N. J. Maciver, J. A. Wofford, H. L. Wieman, J. J. Hammen, and J. C. Rathmell. 2008. Glucose uptake is limiting in T cell activation and requires CD28-mediated Akt-dependent and independent pathways. *J. Immunol.* 180: 4476–4486.
- Krauss, S., M. D. Brand, and F. Buttgerit. 2001. Signaling takes a breath—new quantitative perspectives on bioenergetics and signal transduction. *Immunity* 15: 497–502.
- Michalek, R. D., V. A. Gerriets, S. R. Jacobs, A. N. Macintyre, N. J. MacIver, E. F. Mason, S. A. Sullivan, A. G. Nichols, and J. C. Rathmell. 2011. Cutting edge: distinct glycolytic and lipid oxidative metabolic programs are essential for effector and regulatory CD4<sup>+</sup> T cell subsets. *J. Immunol.* 186: 3299–3303.
- Shi, L. Z., R. Wang, G. Huang, P. Vogel, G. Neale, D. R. Green, and H. Chi. 2011. HIF1 $\alpha$ -dependent glycolytic pathway orchestrates a metabolic checkpoint for the differentiation of TH17 and Treg cells. *J. Exp. Med.* 208: 1367–1376.
- Pearce, E. L., M. C. Walsh, P. J. Cajas, G. M. Harms, H. Shen, L. S. Wang, R. G. Jones, and Y. Choi. 2009. Enhancing CD8 T-cell memory by modulating fatty acid metabolism. *Nature* 460: 103–107.
- Araki, K., A. P. Turner, V. O. Shaffer, S. Gangappa, S. A. Keller, M. F. Bachmann, C. P. Larsen, and R. Ahmed. 2009. mTOR regulates memory CD8 T-cell differentiation. *Nature* 460: 108–112.
- Michalek, R. D., and J. C. Rathmell. 2010. The metabolic life and times of a T-cell. *Immunol. Rev.* 236: 190–202.
- Finlay, D., and D. A. Cantrell. 2011. Metabolism, migration and memory in cytotoxic T cells. *Nat. Rev. Immunol.* 11: 109–117.
- Frauwirth, K. A., J. L. Riley, M. H. Harris, R. V. Parry, J. C. Rathmell, D. R. Plas, R. L. Elstrom, C. H. June, and C. B. Thompson. 2002. The CD28 signaling pathway regulates glucose metabolism. *Immunity* 16: 769–777.
- Krawczyk, C. M., T. Holowka, J. Sun, J. Blagih, E. Amiel, R. J. DeBerardinis, J. R. Cross, E. Jung, C. B. Thompson, R. G. Jones, and E. J. Pearce. 2010. Toll-like receptor-induced changes in glycolytic metabolism regulate dendritic cell activation. *Blood* 115: 4742–4749.
- Rathmell, J. C., R. L. Elstrom, R. M. Cinalli, and C. B. Thompson. 2003. Activated Akt promotes increased resting T cell size, CD28-independent T cell growth, and development of autoimmunity and lymphoma. *Eur. J. Immunol.* 33: 2223–2232.
- Laplanche, M., and D. M. Sabatini. 2009. mTOR signaling at a glance. *J. Cell Sci.* 122: 3589–3594.
- Powell, J. D., and G. M. Delgoffe. 2010. The mammalian target of rapamycin: linking T cell differentiation, function, and metabolism. *Immunity* 33: 301–311.
- Delgoffe, G. M., T. P. Kole, Y. Zheng, P. E. Zarek, K. L. Matthews, B. Xiao, P. F. Worley, S. C. Kozma, and J. D. Powell. 2009. The mTOR kinase differentially regulates effector and regulatory T cell lineage commitment. *Immunity* 30: 832–844.
- Rao, R. R., Q. Li, K. Odunsi, and P. A. Shrikant. 2010. The mTOR kinase determines effector versus memory CD8<sup>+</sup> T cell fate by regulating the expression of transcription factors T-bet and Eomesodermin. *Immunity* 32: 67–78.
- Lee, K., P. Gudapati, S. Dragovic, C. Spencer, S. Joyce, N. Killen, M. A. Magnuson, and M. Boothby. 2010. Mammalian target of rapamycin protein complex 2 regulates differentiation of Th1 and Th2 cell subsets via distinct signaling pathways. *Immunity* 32: 743–753.
- Battaglia, M., A. Stabilini, and M. G. Roncarolo. 2005. Rapamycin selectively expands CD4<sup>+</sup>CD25<sup>+</sup>FoxP3<sup>+</sup> regulatory T cells. *Blood* 105: 4743–4748.
- Kopf, H., G. M. de la Rosa, O. M. Howard, and X. Chen. 2007. Rapamycin inhibits differentiation of Th17 cells and promotes generation of FoxP3<sup>+</sup> T regulatory cells. *Int. Immunopharmacol.* 7: 1819–1824.
- Alessi, D. R., K. Sakamoto, and J. R. Bayascas. 2006. LKB1-dependent signaling pathways. *Annu. Rev. Biochem.* 75: 137–163.
- Hemminki, A., D. Markie, I. Tomlinson, E. Avizienyte, S. Roth, A. Loukola, G. Bignell, W. Warren, M. Aminoff, P. Höglund, et al. 1998. A serine/threonine kinase gene defective in Peutz-Jeghers syndrome. *Nature* 391: 184–187.
- Jenne, D. E., H. Reimann, J. Nezu, W. Friedel, S. Loff, R. Jeschke, O. Müller, W. Back, and M. Zimmer. 1998. Peutz-Jeghers syndrome is caused by mutations in a novel serine threonine kinase. *Nat. Genet.* 18: 38–43.
- Gan, B., J. Hu, S. Jiang, Y. Liu, E. Sahin, L. Zhuang, E. Fletcher-Sananikone, S. Colla, Y. A. Wang, L. Chin, and R. A. Depinho. 2010. Lkb1 regulates quiescence and metabolic homeostasis of haematopoietic stem cells. *Nature* 468: 701–704.
- Gurumurthy, S., S. Z. Xie, B. Alagesan, J. Kim, R. Z. Yusuf, B. Saez, A. Tzatsos, F. Ozsolak, P. Milos, F. Ferrari, et al. 2010. The Lkb1 metabolic sensor maintains haematopoietic stem cell survival. *Nature* 468: 659–663.
- Nakada, D., T. L. Saunders, and S. J. Morrison. 2010. Lkb1 regulates cell cycle and energy metabolism in haematopoietic stem cells. *Nature* 468: 653–658.
- Inoki, K., T. Zhu, and K. L. Guan. 2003. TSC2 mediates cellular energy response to control cell growth and survival. *Cell* 115: 577–590.

30. Shaw, R. J., M. Kosmatka, N. Bardeesy, R. L. Hurley, L. A. Witters, R. A. DePinho, and L. C. Cantley. 2004. The tumor suppressor LKB1 kinase directly activates AMP-activated kinase and regulates apoptosis in response to energy stress. *Proc. Natl. Acad. Sci. USA* 101: 3329–3335.
31. Hawley, S. A., J. Boudeau, J. L. Reid, K. J. Mustard, L. Udd, T. P. Mäkelä, D. R. Alessi, and D. G. Hardie. 2003. Complexes between the LKB1 tumor suppressor, STRAD  $\alpha/\beta$  and MO25  $\alpha/\beta$  are upstream kinases in the AMP-activated protein kinase cascade. *J. Biol.* 2: 28.
32. Woods, A., S. R. Johnstone, K. Dickerson, F. C. Leiper, L. G. Fryer, D. Neumann, U. Schlattner, T. Wallimann, M. Carlson, and D. Carling. 2003. LKB1 is the upstream kinase in the AMP-activated protein kinase cascade. *Curr. Biol.* 13: 2004–2008.
33. Hardie, D. G. 2007. AMP-activated/SNFI protein kinases: conserved guardians of cellular energy. *Nat. Rev. Mol. Cell Biol.* 8: 774–785.
34. Mayer, A., S. Denanglaire, B. Viollet, O. Leo, and F. Andris. 2008. AMP-activated protein kinase regulates lymphocyte responses to metabolic stress but is largely dispensable for immune cell development and function. *Eur. J. Immunol.* 38: 948–956.
35. Cao, Y., H. Li, H. Liu, C. Zheng, H. Ji, and X. Liu. 2010. The serine/threonine kinase LKB1 controls thymocyte survival through regulation of AMPK activation and Bcl-XL expression. *Cell Res.* <http://www.ncbi.nlm.nih.gov/pubmed/2002938920>: 99–108.
36. Tamás, P., A. Macintyre, D. Finlay, R. Clarke, C. Feijoo-Carnero, A. Ashworth, and D. Cantrell. 2010. LKB1 is essential for the proliferation of T-cell progenitors and mature peripheral T cells. *Eur. J. Immunol.* 40: 242–253.
37. Bardeesy, N., M. Sinha, A. F. Hezel, S. Signoretti, N. A. Hathaway, N. E. Sharpless, M. Loda, D. R. Carrasco, and R. A. DePinho. 2002. Loss of the Lkb1 tumour suppressor provokes intestinal polyposis but resistance to transformation. *Nature* 419: 162–167.
38. Viollet, B., F. Andreelli, S. B. Jørgensen, C. Perrin, A. Geloën, D. Flamez, J. Mu, C. Lenzner, O. Baud, M. Bennoun, et al. 2003. The AMP-activated protein kinase  $\alpha 2$  catalytic subunit controls whole-body insulin sensitivity. *J. Clin. Invest.* 111: 91–98.
39. Jones, R. G., T. Bui, C. White, M. Madesh, C. M. Krawczyk, T. Lindsten, B. J. Hawkins, S. Kubek, K. A. Frauwirth, Y. L. Wang, et al. 2007. The proapoptotic factors Bax and Bak regulate T Cell proliferation through control of endoplasmic reticulum  $\text{Ca}^{2+}$  homeostasis. *Immunity* 27: 268–280.
40. Rathmell, J. C., C. J. Fox, D. R. Plas, P. S. Hammerman, R. M. Cinalli, and C. B. Thompson. 2003. Akt-directed glucose metabolism can prevent Bax conformation change and promote growth factor-independent survival. *Mol. Cell Biol.* 23: 7315–7328.
41. Krawczyk, C. M., H. Shen, and E. J. Pearce. 2007. Functional plasticity in memory T helper cell responses. *J. Immunol.* 178: 4080–4088.
42. Bungard, D., B. J. Fuerth, P. Y. Zeng, B. Faubert, N. L. Maas, B. Viollet, D. Carling, C. B. Thompson, R. G. Jones, and S. L. Berger. 2010. Signaling kinase AMPK activates stress-promoted transcription via histone H2B phosphorylation. *Science* 329: 1201–1205.
43. Wieman, H. L., J. A. Wofford, and J. C. Rathmell. 2007. Cytokine stimulation promotes glucose uptake via phosphatidylinositol-3 kinase/Akt regulation of Glut1 activity and trafficking. *Mol. Biol. Cell* 18: 1437–1446.
44. Plas, D. R., S. Talapatra, A. L. Edinger, J. C. Rathmell, and C. B. Thompson. 2001. Akt and Bcl-xL promote growth factor-independent survival through distinct effects on mitochondrial physiology. *J. Biol. Chem.* 276: 12041–12048.
45. Shackelford, D. B., and R. J. Shaw. 2009. The LKB1-AMPK pathway: metabolism and growth control in tumour suppression. *Nat. Rev. Cancer* 9: 563–575.
46. Cham, C. M., G. Driessens, J. P. O’Keefe, and T. F. Gajewski. 2008. Glucose deprivation inhibits multiple key gene expression events and effector functions in CD8<sup>+</sup> T cells. *Eur. J. Immunol.* 38: 2438–2450.
47. Shaw, R. J., K. A. Lamia, D. Vasquez, S. H. Koo, N. Bardeesy, R. A. Depinho, M. Montminy, and L. C. Cantley. 2005. The kinase LKB1 mediates glucose homeostasis in liver and therapeutic effects of metformin. *Science* 310: 1642–1646.
48. Motoyama, N., F. Wang, K. A. Roth, H. Sawa, K. Nakayama, K. Nakayama, I. Negishi, S. Senju, Q. Zhang, S. Fujii, et al. 1995. Massive cell death of immature hematopoietic cells and neurons in Bcl-x-deficient mice. *Science* 267: 1506–1510.
49. Tamás, P., S. A. Hawley, R. G. Clarke, K. J. Mustard, K. Green, D. G. Hardie, and D. A. Cantrell. 2006. Regulation of the energy sensor AMP-activated protein kinase by antigen receptor and  $\text{Ca}^{2+}$  in T lymphocytes. *J. Exp. Med.* 203: 1665–1670.
50. Huang, X., S. Wullschlegel, N. Shpiro, V. A. McGuire, K. Sakamoto, Y. L. Woods, W. McBurnie, S. Fleming, and D. R. Alessi. 2008. Important role of the LKB1-AMPK pathway in suppressing tumorigenesis in PTEN-deficient mice. *Biochem. J.* 412: 211–221.
51. Lizcano, J. M., O. Göransson, R. Toth, M. Deak, N. A. Morrice, J. Boudeau, S. A. Hawley, L. Udd, T. P. Mäkelä, D. G. Hardie, and D. R. Alessi. 2004. LKB1 is a master kinase that activates 13 kinases of the AMPK subfamily, including MARK/PAR-1. *EMBO J.* 23: 833–843.
52. Tiaainen, M., K. Vaahtomeri, A. Ylikorkala, and T. P. Mäkelä. 2002. Growth arrest by the LKB1 tumor suppressor: induction of p21(WAF1/CIP1). *Hum. Mol. Genet.* 11: 1497–1504.
53. Boyman, O., S. Létourneau, C. Krieg, and J. Sprent. 2009. Homeostatic proliferation and survival of naïve and memory T cells. *Eur. J. Immunol.* 39: 2088–2094.
54. Surh, C. D., and J. Sprent. 2008. Homeostasis of naïve and memory T cells. *Immunity* 29: 848–862.
55. Yang, K., G. Neale, D. R. Green, W. He, and H. Chi. 2011. The tumor suppressor Tsc1 enforces quiescence of naïve T cells to promote immune homeostasis and function. *Nat. Immunol.* 12: 888–897.
56. Wu, Q., Y. Liu, C. Chen, T. Ikenoue, Y. Qiao, C. S. Li, W. Li, K. L. Guan, and P. Zheng. 2011. The tuberous sclerosis complex-Mammalian target of rapamycin pathway maintains the quiescence and survival of naïve T cells. *J. Immunol.* 187: 1106–1112.
57. O’Brien, T. F., B. K. Gorenflo, D. Xie, S. Srivatsan, I. X. McLeod, Y. W. He, and X. P. Zhong. 2011. Regulation of T cell survival and mitochondrial homeostasis by TSC1. *Eur. J. Immunol.* DOI: 10.1002/eji.201141411.
58. Sag, D., D. Carling, R. D. Stout, and J. Suttles. 2008. Adenosine 5'-monophosphate-activated protein kinase promotes macrophage polarization to an anti-inflammatory functional phenotype. *J. Immunol.* 181: 8633–8641.
59. Nath, N., S. Giri, R. Prasad, M. L. Salem, A. K. Singh, and I. Singh. 2005. 5-aminimidazole-4-carboxamide ribonucleoside: a novel immunomodulator with therapeutic efficacy in experimental autoimmune encephalomyelitis. *J. Immunol.* 175: 566–574.
60. Cao, Y., H. Li, H. Liu, M. Zhang, Z. Hua, H. Ji, and X. Liu. 2011. LKB1 regulates TCR-mediated PLC $\gamma$ 1 activation and thymocyte positive selection. *EMBO J.* 30: 2083–2093.
61. Hawley, S. A., D. A. Pan, K. J. Mustard, L. Ross, J. Bain, A. M. Edelman, B. G. Frenguelli, and D. G. Hardie. 2005. Calmodulin-dependent protein kinase kinase-beta is an alternative upstream kinase for AMP-activated protein kinase. *Cell Metab.* 2: 9–19.
62. Woods, A., K. Dickerson, R. Heath, S. P. Hong, M. Momcilovic, S. R. Johnstone, M. Carlson, and D. Carling. 2005.  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase kinase-beta acts upstream of AMP-activated protein kinase in mammalian cells. *Cell Metab.* 2: 21–33.
63. Maciver, N. J., S. R. Jacobs, H. L. Wieman, J. A. Wofford, J. L. Colloff, and J. C. Rathmell. 2008. Glucose metabolism in lymphocytes is a regulated process with significant effects on immune cell function and survival. *J. Leukoc. Biol.* 84: 949–957.
64. Jacobs, S. R., R. D. Michalek, and J. C. Rathmell. 2010. IL-7 is essential for homeostatic control of T cell metabolism in vivo. *J. Immunol.* 184: 3461–3469.