

**BULK ANTIBODIES**  
for *in vivo*  
RESEARCH

**α-CD4** **α-CD8** **α-CD25** **α-NK1.1** **α-Ly6G**

Discover More

BioCell



## Correction: Exosomes Derived from Burkitt's Lymphoma Cell Lines Induce Proliferation, Differentiation, and Class-Switch Recombination in B Cells

This information is current as of September 20, 2019.

Cindy Gutzeit, Noemi Nagy, Maurizio Gentile, Katarina Lyberg, Janine Gumz, Helen Vallhov, Irene Puga, Eva Klein, Susanne Gabrielsson, Andrea Cerutti and Annika Scheynius

*J Immunol* published online 21 June 2019  
<http://www.jimmunol.org/content/early/2019/06/21/jimmunol.1900638.citation>

Why *The JI*? [Submit online.](#)

- **Rapid Reviews! 30 days\*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

*\*average*

- Subscription** Information about subscribing to *The Journal of Immunology* is online at: <http://jimmunol.org/subscription>
- Permissions** Submit copyright permission requests at: <http://www.aai.org/About/Publications/JI/copyright.html>
- Email Alerts** Receive free email-alerts when new articles cite this article. Sign up at: <http://jimmunol.org/alerts>


*The Journal of Immunology* is published twice each month by The American Association of Immunologists, Inc., 1451 Rockville Pike, Suite 650, Rockville, MD 20852  
 Copyright © 2019 by The American Association of Immunologists, Inc. All rights reserved.  
 Print ISSN: 0022-1767 Online ISSN: 1550-6606.



## Corrections

Gutzeit, C., N. Nagy, M. Gentile, K. Lyberg, J. Gumz, H. Vallhov, I. Puga, E. Klein, S. Gabrielsson, A. Cerutti, and A. Scheynius. 2014. Exosomes derived from Burkitt's lymphoma cell lines induce proliferation, differentiation, and class-switch recombination in B cells. *J. Immunol.* 192: 5852–5862.

In the published version of Fig. 6A, the FACS dotplot shown for the stimulation of DG75-COex at 25  $\mu$ g in the presence of IL-21 was inadvertently duplicated from the 5  $\mu$ g/IL-21 panel. The indicated frequency of 5% of CD38<sup>high</sup>CD20<sup>low</sup> B cells, however, is correct. This correct value of 5% was used in Fig. 6B, so the error in Fig. 6A did not influence the interpretation of the results and conclusions of this work. Fig. 6A has now been corrected with the correct dotplot for stimulation of DG75-COex at 25  $\mu$ g in the presence of IL-21. The corrected version of Fig. 6 is shown below. The figure legend was correct as published and is shown below for reference. Fig. 6 has also been corrected in the online version of the article, which now differs from the print version originally published.



**FIGURE 6.** DG75-LMP1ex induces differentiation into a CD38<sup>high</sup>CD20<sup>low</sup> plasmablast-like B cell population. CFSE-labeled B cells ( $1.5 \times 10^5$  cells/200  $\mu$ l) were either left unstimulated (co) or stimulated with IL-21, CD40L, or IL-21 + CD40L or with 5 or 25  $\mu$ g DG75 exosomes/well, with or without IL-21. After 5 d of culture, B cells were stained for CD19, CD20, and CD38 and assessed by flow cytometry. **(A)** Representative plots from one donor are shown, and the percentage of CD38<sup>high</sup>CD20<sup>low</sup> B cells is indicated. **(B)** Combined results from four blood donors (mean  $\pm$  SEM). \* $p \leq 0.05$ , Mann-Whitney  $U$  test.